

Using RAMPAGE to identify and annotate regulatory elements in insect genomes

R. Taylor Raborn^{*1,2} and Volker P. Brendel^{1,2}

¹Department of Biology, Indiana University

²School of Informatics and Computing, Indiana University

Department of Biology and School of Informatics and Computing,
Indiana University

212 S. Hawthorne Drive 205 Simon Hall, Bloomington, IN 47401, USA
<http://www.brendelgroup.org>

Abstract. Application of Transcription Start Site (TSS) profiling technologies, coupled with large-scale next-generation sequencing (NGS) has yielded valuable insights into the location, structure and activity of promoters across diverse metazoan model systems. In insects, TSS profiling has been used to characterize the promoter architecture of *D. melanogaster*, and, shortly thereafter, to reveal widespread transposon-driven alternative promoter usage.

In this chapter we highlight the utility of one TSS profiling method, RAMPAGE (RNA annotation and mapping of promoters for analysis of gene expression), for the precise, quantitative identification of promoters in insect genomes. We demonstrate this using our tools GoRAMPAGE and TSSrchitect, providing details instructions with the aim of taking the user from raw reads to processed results.

Keywords: *cis*-regulatory regions, promoter architecture, transcription initiation, transcription start sites (TSSs)

1 Introduction

1.1 TSS Profiling Identifies Promoters at Genome-Scale

The promoter, defined in eukaryotes as the genomic region bound by RNA Polymerase II immediately prior to transcription initiation [1], is the site where regulatory signals unite to direct gene expression. The identification of promoter regions is a valuable step for understanding the *cis*-regulatory signals that are present in an organism, and is important for genome annotation. However, despite the rapid accumulation of genome sequences across metazoan and arthropod diversity, accurate annotation of promoter regions remains sparse. This is because—empirical mapping of TSSs—precisely identifying sequence motifs that demarcate the promoter is unreliable. In contrast with current *in*

* Correspondence: rtraborn@indiana.edu

silico approaches, direct mapping of TSSs identifies the location of the core promoter. Cap Analysis of Gene Expression (CAGE) [2], one of the first methods devised to identify 5'-ends of mRNAs at large-scale, involves selective capture of 5'-capped transcripts, first-strand reverse-transcription and ligation of a short oligonucleotide (CAGE tag). CAGE was initially utilized by the FANTOM (Functional Annotation of the Mammalian Genome) consortium to identify promoter architecture in human and mouse [3], providing the first glimpse of the global landscape of transcription initiation. At the onset of the NGS era, CAGE was coupled with massively-parallel sequencing to generate 5'-ends of mRNAs at substantially higher scale. This advance provided more extensive coverage of the expressed transcriptome, and provided increased sensitivity for quantitative measurements *i.e.* measurement of promoter activity.

1.2 Promoter Architecture of *Drosophila melanogaster*

Hoskins and colleagues [4] performed CAGE in *D. melanogaster* as part of the modENCODE consortium, identifying promoters at large-scale and characterizing the promoter architecture of an insect genome for the first time. Hoskins [4] indicated that TSS distributions at *Drosophila* promoters exhibit a range of shapes that can be generally grouped into two major classifications: *peaked* and *broad*. Peaked promoters have a single, major TSS position occupying a narrow genomic region, whereas broad promoters lack a single, major TSS and contain TSSs across a wider region [5][6]. The authors also showed a strong association between promoter class and motif composition (consistent with previous findings [5, 7]). Peaked promoters were associated with positionally-enriched *cis*-regulatory motifs including TATA, Initiator (Inr) and DPE, while broad promoters contained an enrichment of less-well characterized motifs, including *Ohler6* and *Ohler7* [8]. The existence of two promoter classes appears to be conserved among metazoans, and has been reported (using TSS profiling methodologies) in insects, cladocerans [9], fish [10] and mammals [11, 6].

1.3 Promoter Structure of Insects

Beyond *D. melanogaster*, few investigations have utilized TSS profiling in insect genomes. As a consequence, what is known about promoter architecture in insects is largely restricted to the *Drosophila* genus. As part of the modENCODE effort, CAGE was performed in multiple tissues and developmental stages of the *Drosophila pseudoobscura*. TSSs were found to be highly similar between species: more than 80% of TSSs (81%) of aligned, CAGE-identified TSSs from *D. pseudoobscura* were positioned within 20nt of their counterparts in *D. melanogaster*. An enrichment of the CA dinucleotide was detected at the TSS ([-1, +1]), and the motifs corresponding to TATA, Inr and DPE were positioned at the same locations relative to the TSS in both species. The one other insect species for which TSS profiling has been applied is the Tsetse fly (*Glossina morsitans morsitans*) [12]. Using TSS-seq (specifically Oligo-capping; for details on this method see [13]), the authors identified 3134 mapping to 1424 genes. The authors found

a preference for CA and AA dinucleotides at the TSS, and observe the major core promoter elements observed in *Drosophila*: TATA, Inr, DPE, in addition to MTE (Motif Ten Element). As in *D. melanogaster*, peaked promoters were more likely to contain TATA and Inr than broad promoters. While the taxonomic sampling of species for TSS profiling has been limited, the existing studies are sufficient to provide a general picture of insect promoter architecture. A major demarcation between the promoter architecture of insects and mammals appears to be the large fraction of mammalian promoters found in CpG islands [12]. CpG island promoters (CPIs) form the largest class of promoter in mammals [14]; by contrast, CPIs are not known to exist as a class in invertebrates.

1.4 Paired-end TSS Profiling with RAMPAGE

The most recent major methodological advance in TSS Profiling is RAMPAGE (RNA Annotation and Mapping of Promoters for the Analysis of Gene Expression) . RAMPAGE is a protocol for 5'-cDNA sequencing that combines cap trapping and template-switching with paired-end sequence information. A key advantage of generating paired-end sequence is transcript connectivity, which provides a direct link between a given 5'-end and its associated mRNA molecule. Because short or spurious RNAs are found within the transcriptome, transcript connectivity allows the TSSs (and thus promoters) of full-length mRNAs to be unambiguously identified, which benefits genome annotation. Batut and colleagues generated libraries from total RNA isolated from 36 stages across the life cycle of *D. melanogaster* providing a comprehensive gene expression and promoter atlas for fruit fly and in the process demonstrating the utility of RAMPAGE. RAMPAGE is currently being applied as part of the latest iteration of ENCODE to identify promoters in human, but as of this writing it has not been applied to any non-*Drosophila* insect species. In anticipation of the future application of TSS profiling into other insect model systems here we provide a documented protocol for the computational processing RAMPAGE data, using selected libraries from Batut *et al.*. This method will consist of two parts: first, we will process, filter and align the sequenced RAMPAGE libraries to the *D. melanogaster* genome. Second, we will identify TSSs and promoters from the aligned sequences and associate them with coding regions. In closing, we will consider further applications of this data and discuss the utility of reproducible workflows in bioinformatic analysis.

2 Materials

The analyses described herein require a workstation capable for modern bioinformatics. An intermediate understanding of the Linux/Unix command line will be extremely useful, although we make efforts to explain the procedures with clarity. In addition, it will likely be necessary for the participant to have superuser privileges on the machine. If you do not have a machine (or access to one) that meets

these requirements, it is recommended that you consider cloud-based cyberinfrastructure, including Amazon Web Services (AWS; <https://aws.amazon.com/>) or CyVerse (<http://www.cyverse.org/>). The former is a well-known pay-per-use solution, while the latter is an NSF-funded resource that is made freely available to the public.

2.1 Hardware Requirements

- x86-64 compatible processors
- At least 8GB RAM
- 30GB+ hard disk space

2.2 Software Requirements

- Operating system: 64 bit Linux (preferred) or Mac OS X (with Command Line Tools from XCode)
- R (version 3.4)
- Bioconductor (version 3.5)
- FASTX-Toolit (version 0.0.13)
- Samtools (version 1.3 or above)
- SRA Toolkit (version 2.3.4-2 or above)
- STAR aligner (version 2.4 or above)
- TagDust (version 2.33)

2.3 Installation of R packages

For installation of the software listed above, please follow the instructions provided by each respective package. Part of our analysis will require the use of R packages found in the Bioconductor suite. To install Bioconductor, please type the following from an R console:

```
source("https://bioconductor.org/biocLite.R")
biocLite()
```

We will use the R package *TSRchitect* to identify promoters from aligned RAMAPGE libraries. First, we will need to install a series of prerequisite packages to *TSRchitect* from Bioconductor. Please install these packages as follows (as before, from an R console):

```
source("https://bioconductor.org/biocLite.R")
biocLite(c("AnnotationHub", "BiocGenerics", "BiocParallel",
"ENCODEExplorer", "GenomicAlignments", "GenomeInfoDb",
"GenomicRanges", "IRanges", "methods",
"Rsamtools", "rtracklayer", "S4Vectors",
"SummarizedExperiment"))
```

To install *TSRchitect*, please type the following from an R console:

```

131 source("https://bioconductor.org/biocLite.R")
132 biocLite("TSRchitect")

```

133 Finally, please confirm that TSRchitect has been installed correctly by load-
 134 ing it from your R console as follows:

```

135 library(TSRchitect)

```

136 3 Methods

137 3.1 Retrieving the RAMPAGE sequence data from NCBI's 138 Gene Expression Omnibus (GEO)

139 To begin our analysis, we must download the RAMPAGE data to our work-
 140 station. We will utilize tools provided by the SRA Toolkit, which should
 141 already be installed on your machine (see **Materials**). The command *fastq-*
 142 *dump* allows one to directly retrieve data from the GEO database using
 143 the appropriate identifier(s). While there are 36 RAMPAGE libraries in the
 144 Batut *et al.* dataset, we will select a subset of these to analyze here. We will
 145 compare samples from selected embryonic (E01h-E04h; E22h-E24h) tissues,
 146 representing the beginning and end of embryonic development. For more
 147 information about the experiment and the available RAMPAGE libraries,
 148 please see the following link: <https://www.ncbi.nlm.nih.gov/Traces/study/?acc=SRP011193>
 149 First, let's proceed with the libraries from early embryonic tissues. Note
 150 that since these fastq files are paired-end, we use the argument *-split-files*
 151 to generate separate files for each read pair.

```

152 fastq-dump --split-files SRR424683
153 fastq-dump --split-files SRR424684
154 fastq-dump --split-files SRR424685
155 fastq-dump --split-files SRR424686

```

156 We continue with downloading the RAMPAGE libraries from late embryonic
 157 tissues:

```

158 fastq-dump --split-files SRR424704
159 fastq-dump --split-files SRR424705
160 fastq-dump --split-files SRR424706

```

161 4 Notes

162 Acknowledgments

163 Disclosure Declaration

164 The authors declare that they have no competing interests.

5 Figures

For \LaTeX users, we recommend using the *graphics* or *graphicx* package and the `\includegraphics` command.

Please check that the lines in line drawings are not interrupted and are of a constant width. Grids and details within the figures must be clearly legible and may not be written one on top of the other. Line drawings should have a resolution of at least 800 dpi (preferably 1200 dpi). The lettering in figures should have a height of 2 mm (10-point type). Figures should be numbered and should have a caption which should always be positioned *under* the figures, in contrast to the caption belonging to a table, which should always appear *above* the table; this is simply achieved as matter of sequence in your source.

Please center the figures or your tabular material by using the `\centering` declaration. Short captions are centered by default between the margins and typeset in 9-point type (Fig. 1 shows an example). The distance between text and figure is preset to be about 8 mm, the distance between figure and caption about 6 mm.

To ensure that the reproduction of your illustrations is of a reasonable quality, we advise against the use of shading. The contrast should be as pronounced as possible.

If screenshots are necessary, please make sure that you are happy with the print quality before you send the files.




Fig. 1. One kernel at x_s (*dotted kernel*) or two kernels at x_i and x_j (*left and right*) lead to the same summed estimate at x_s . This shows a figure consisting of different types of lines. Elements of the figure described in the caption should be set in italics, in parentheses, as shown in this sample caption.

186 Please define figures (and tables) as floating objects. Please avoid using op-
 187 tional location parameters like “[h]” for “here”.

188 5.1 Formulas

189 Displayed equations or formulas are centered and set on a separate line (with an
 190 extra line or halfline space above and below). Displayed expressions should be
 191 numbered for reference. The numbers should be consecutive within each section
 192 or within the contribution, with numbers enclosed in parentheses and set on the
 193 right margin – which is the default if you use the *equation* environment, e.g.,

$$\psi(u) = \int_o^T \left[\frac{1}{2} (A_o^{-1}u, u) + N^*(-u) \right] dt . \quad (1)$$

194 Equations should be punctuated in the same way as ordinary text but with
 195 a small space before the end punctuation mark.

196 5.2 Footnotes

197 The superscript numeral used to refer to a footnote appears in the text either
 198 directly after the word to be discussed or – in relation to a phrase or a sentence –
 199 following the punctuation sign (comma, semicolon, or period). Footnotes should
 200 appear at the bottom of the normal text area, with a line of about 2 cm set
 201 immediately above them.¹

202 5.3 Program Code

203 Program listings or program commands in the text are normally set in typewriter
 204 font, e.g., CMTT10 or Courier.

205 *Example of a Computer Program*

```
206 program Inflation (Output)
207   {Assuming annual inflation rates of 7%, 8%, and 10%,...
208   years};
209   const
210     MaxYears = 10;
211   var
212     Year: 0..MaxYears;
213     Factor1, Factor2, Factor3: Real;
214   begin
215     Year := 0;
216     Factor1 := 1.0; Factor2 := 1.0; Factor3 := 1.0;
217     WriteLn('Year 7% 8% 10%'); WriteLn;
```

¹ The footnote numeral is set flush left and the text follows with the usual word spacing.

```

218     repeat
219         Year := Year + 1;
220         Factor1 := Factor1 * 1.07;
221         Factor2 := Factor2 * 1.08;
222         Factor3 := Factor3 * 1.10;
223         WriteLn(Year:5,Factor1:7:3,Factor2:7:3,Factor3:7:3)
224     until Year = MaxYears
225 end.

```

226 (Example from Jensen K., Wirth N. (1991) Pascal user manual and report. Springer,
 227 New York)

228 5.4 Citations

229 For citations in the text please use square brackets and consecutive numbers:
 230 [?], [?], [?] – provided automatically by L^AT_EX's `\cite ... \bibitem` mechanism.

231 5.5 Page Numbering and Running Heads

232 There is no need to include page numbers. If your paper title is too long to serve
 233 as a running head, it will be shortened. Your suggestion as to how to shorten it
 234 would be most welcome.

235 6 References

References

1. J. T. Kadonaga, "Perspectives on the RNA polymerase II core promoter." *Wiley Interdisciplinary Reviews: Developmental Biology*, vol. 1, no. 1, pp. 40–51, Jan. 2012.
2. R. Kodzius, M. Kojima, H. Nishiyori, M. Nakamura, S. Fukuda, M. Tagami, D. Sasaki, K. Imamura, C. Kai, M. Harbers, Y. Hayashizaki, and P. Carninci, "CAGE: cap analysis of gene expression." *Nature Methods*, vol. 3, no. 3, pp. 211–222, Mar. 2006.
3. P. Carninci, T. Kasukawa, S. Katayama, J. Gough, M. C. Frith, N. Maeda, R. Oyama, T. Ravasi, B. Lenhard, C. Wells, R. Kodzius, K. Shimokawa, V. B. Bajic, S. E. Brenner, S. Batalov, A. R. R. Forrest, M. Zavolan, M. J. Davis, L. G. Wilming, V. Aidinis, J. E. Allen, A. Ambesi-Impimbato, R. Apweiler, R. N. Aturaliya, T. L. Bailey, M. Bansal, L. Baxter, K. W. Beisel, T. Bersano, H. Bono, A. M. Chalk, K. P. Chiu, V. Choudhary, A. Christoffels, D. R. Clutterbuck, M. L. Crowe, E. Dalla, B. P. Dalrymple, B. de Bono, G. Della Gatta, D. di Bernardo, T. Down, P. Engstrom, M. Fagiolini, G. Faulkner, C. F. Fletcher, T. Fukushima, M. Furuno, S. Futaki, M. Gariboldi, P. Georgii-Hemming, T. R. Gingeras, T. Gojobori, R. E. Green, S. Gustincich, M. Harbers, Y. Hayashi, T. K. Hensch, N. Hirokawa, D. Hill, L. Huminiecki, M. Iacono, K. Ikeo, A. Iwama, T. Ishikawa, M. Jakt, A. Kanapin, M. Katoh, Y. Kawasawa, J. Kelso, H. Kitamura, H. Kitano, G. Kollias, S. P. T. Krishnan, A. Kruger, S. K. Kummerfeld, I. V. Kurochkin, L. F. Lareau, D. Lazarevic, L. Lipovich, J. Liu, S. Liuni, S. McWilliam, M. Madan Babu, M. Madera, L. Marchionni, H. Matsuda, S. Matsuzawa, H. Miki, F. Mignone, S. Miyake, K. Morris, S. Mottagui-Tabar, N. Mulder, N. Nakano, H. Nakauchi, P. Ng, R. Nilsson, S. Nishiguchi, S. Nishikawa, F. Nori, O. Ohara, Y. Okazaki, V. Orlando, K. C. Pang, W. J. Pavan, G. Pavesi, G. Pesole, N. Petrovsky, S. Piazza, J. Reed, J. F. Reid, B. Z. Ring, M. Ringwald, B. Rost, Y. Ruan, S. L. Salzberg, A. Sandelin, C. Schneider, C. Schönbach, K. Sekiguchi, C. A. M. Semple, S. Seno, L. Sessa, Y. Sheng, Y. Shibata, H. Shimada, K. Shimada, D. Silva, B. Sinclair, S. Sperling, E. Stupka, K. Sugiura, R. Sultana, Y. Takenaka, K. Taki, K. Tammoja, S. L. Tan, S. Tang, M. S. Taylor, J. Tegner, S. A. Teichmann, H. R. Ueda, E. van Nimwegen, R. Verardo, C. L. Wei, K. Yagi, H. Yamanishi, E. Zabarovsky, S. Zhu, A. Zimmer, W. Hide, C. Bult, S. M. Grimmond, R. D. Teasdale, E. T. Liu, V. Brusic, J. Quackenbush, C. Wahlestedt, J. S. Mattick, D. A. Hume, C. Kai, D. Sasaki, Y. Tomaru, S. Fukuda, M. Kanamori-Katayama, M. Suzuki, J. Aoki, T. Arakawa, J. Iida, K. Imamura, M. Itoh, T. Kato, H. Kawaji, N. Kawagashira, T. Kawashima, M. Kojima, S. Kondo, H. Konno, K. Nakano, N. Ninomiya, T. Nishio, M. Okada, C. Plessy, K. Shibata, T. Shiraki, S. Suzuki, M. Tagami, K. Waki, A. Watahiki, Y. Okamura-Oho, H. Suzuki, J. Kawai, Y. Hayashizaki, F. Consortium, R. G. E. R. Group, and G. S. G. G. N. P. C. Group, "The transcriptional landscape of the mammalian genome," *Science (New York, NY)*, vol. 309, no. 5740, pp. 1559–1563, Sep. 2005.
4. R. A. Hoskins, R. A. Hoskins, J. M. Landolin, J. M. Landolin, J. B. Brown, J. B. Brown, J. E. Sandler, J. E. Sandler, H. Takahashi, H. Takahashi, T. Lassmann, T. Lassmann, C. Yu, C. Yu, B. W. Booth, B. W. Booth, D. Zhang, D. Zhang, K. H. Wan, K. H. Wan, L. Yang, L. Yang, N. Boley, N. Boley, J. Andrews, J. Andrews, T. C. Kaufman, T. C. Kaufman, B. R. Graveley, B. R. Graveley, P. J. Bickel, P. J. Bickel, P. Carninci, J. W. Carlson, J. W. Carlson, S. E. Celniker,

- 284 and S. E. Celniker, "Genome-wide analysis of promoter architecture in *Drosophila*
285 *melanogaster*." *Genome Research*, vol. 21, no. 2, pp. 182–192, Feb. 2011.
- 286 5. E. A. Rach, H.-Y. Yuan, W. H. Majoros, P. Tomancak, and U. Ohler, "Motif
287 composition, conservation and condition-specificity of single and alternative tran-
288 scription start sites in the *Drosophila* genome." *Genome Biology*, vol. 10, no. 7, p.
289 R73, 2009.
- 290 6. B. Lenhard, A. Sandelin, and P. Carninci, "Metazoan promoters: emerging char-
291 acteristics and insights into transcriptional regulation." *Nature Reviews Genetics*,
292 vol. 13, no. 4, pp. 233–245, Apr. 2012.
- 293 7. T. Ni, D. L. Corcoran, E. A. Rach, S. Song, E. P. Spana, Y. Gao, U. Ohler,
294 and J. Zhu, "A paired-end sequencing strategy to map the complex landscape of
295 transcription initiation." *Nature Methods*, vol. 7, no. 7, pp. 521–527, Jul. 2010.
- 296 8. U. Ohler, G.-c. Liao, H. Niemann, and G. M. Rubin, "Computational analysis of
297 core promoters in the *Drosophila* genome." *Genome Biology*, vol. 3, no. 12, pp.
298 research0087.1–0087.12, 2002.
- 299 9. R. T. Raborn, K. Spitze, V. P. Brendel, and M. Lynch, "Promoter Architecture
300 and Sex-Specific Gene Expression in *Daphnia pulex*." *Genetics*, vol. 204, no. 2, pp.
301 593–612, Aug. 2016.
- 302 10. C. Nepal, Y. Hadzhiev, C. Previti, V. Haberle, N. Li, H. Takahashi, A. M. M.
303 Suzuki, Y. Sheng, R. F. Abdelhamid, S. Anand, J. Gehrig, A. Akalin, C. E. M.
304 Kockx, A. A. J. van der Sloot, W. F. J. van IJcken, O. Armant, S. Rastegar,
305 C. Watson, U. Strahle, E. Stupka, P. Carninci, B. Lenhard, and F. Muller, "Dy-
306 namic regulation of the transcription initiation landscape at single nucleotide res-
307 olution during vertebrate embryogenesis," *Genome Research*, vol. 23, no. 11, pp.
308 1938–1950, Nov. 2013.
- 309 11. P. Carninci, A. Sandelin, B. Lenhard, S. Katayama, K. Shimokawa, J. Ponjavic,
310 C. A. M. Semple, M. S. Taylor, P. G. Engström, M. C. Frith, A. R. R. For-
311 rest, W. B. Alkema, S. L. Tan, C. Plessy, R. Kodzius, T. Ravasi, T. Kasukawa,
312 S. Fukuda, M. Kanamori-Katayama, Y. Kitazume, H. Kawaji, C. Kai, M. Naka-
313 mura, H. Konno, K. Nakano, S. Mottagui-Tabar, P. Arner, A. Chesi, S. Gustincich,
314 F. Persichetti, H. Suzuki, S. M. Grimmond, C. A. Wells, V. Orlando, C. Wahlest-
315 edt, E. T. Liu, M. Harbers, J. Kawai, V. B. Bajic, D. A. Hume, and Y. Hayashizaki,
316 "Genome-wide analysis of mammalian promoter architecture and evolution," *Na-
317 ture Genetics*, vol. 38, no. 6, pp. 626–635, Apr. 2006.
- 318 12. S. Mwangi, G. Attardo, Y. Suzuki, S. Aksoy, and A. Christoffels, "TSS seq based
319 core promoter architecture in blood feeding Tsetse fly (*Glossina morsitans mor-
320 sitans*) vector of Trypanosomiasis," *BMC Genomics*, vol. 16, no. 1, p. 722, Sep.
321 2015.
- 322 13. K. Tsuchihara, Y. Suzuki, H. Wakaguri, T. Irie, K. Tanimoto, S.-i. Hashimoto,
323 K. Matsushima, J. Mizushima-Sugano, R. Yamashita, K. Nakai, D. Bentley, H. Es-
324 umi, and S. Sugano, "Massive transcriptional start site analysis of human genes in
325 hypoxia cells," *Nucleic Acids Research*, vol. 37, no. 7, pp. 2249–2263, Apr. 2009.
- 326 14. N. Cvetesic and B. Lenhard, "Core promoters across the genome," *Nature Biotech-
327 nology*, vol. 35, no. 2, pp. 123–124, Feb. 2017.

328 In order to permit cross referencing within LNCS-Online, and eventually
329 between different publishers and their online databases, LNCS will, from now
330 on, be standardizing the format of the references. This new feature will increase
331 the visibility of publications and facilitate academic research considerably. Please
332 base your references on the examples below. References that don't adhere to this

333 style will be reformatted by Springer. You should therefore check your references
 334 thoroughly when you receive the final pdf of your paper. The reference section
 335 must be complete. You may not omit references. Instructions as to where to find
 336 a fuller version of the references are not permissible.

337 We only accept references written using the latin alphabet. If the title of the
 338 book you are referring to is in Russian or Chinese, then please write (in Russian)
 339 or (in Chinese) at the end of the transcript or translation of the title.

340 The following section shows a sample reference list with entries for journal
 341 articles [?], an LNCS chapter [?], a book [?], proceedings without editors [?] and
 342 [?], as well as a URL [?]. Please note that proceedings published in LNCS are
 343 not cited with their full titles, but with their acronyms!

344 7 Checklist of Items to be Sent to Volume Editors

345 Here is a checklist of everything the volume editor requires from you:

- 346 ☐ The final L^AT_EX source files
- 347 ☐ A final PDF file
- 348 ☐ A copyright form, signed by one author on behalf of all of the authors of the
 349 paper.
- 350 ☐ A readme giving the name and email address of the corresponding author.