

1 Degree Correlation Coefficient

For all the computations, please see `Problem 6-1.ipynb`

1.1 Degree Correlation Matrix

$$E = \begin{bmatrix} 0 & 0 & 1/8 \\ 0 & 1/4 & 1/4 \\ 1/8 & 1/4 & 0 \end{bmatrix}$$

1.2 Probabilities q_k

$$\begin{aligned} q_1 &= 0 + 0 + 1/8 = 1/8 \\ q_2 &= 0 + 1/4 + 1/4 = 1/2 \\ q_3 &= 1/8 + 1/4 + 0 = 3/8 \end{aligned}$$

1.3 Degree Correlation Coefficient

$$r = -0.7142857142857143$$

Based on r , the given network is disassortative ($r < 0$)

2 Degree Correlations in Random Graphs

2.1

$$p(a_{ij} = 1) = \frac{L}{\frac{N(N-1)}{2}} = \frac{2L}{N(N-1)}$$

2.2

If link a_{xy} exists, then $p(a_{ij})(i, j \neq x, y)$ goes down, since we know that one link is already present and the final number of links is fixed. Similarly, if the link doesn't exist, the probability of all other links goes up.

$$\begin{aligned} p(a_{ij} = 1 | a_{xy} = 1) &= \frac{L-1}{\frac{N(N-1)}{2} - 1} \\ &= 2 \frac{L-1}{N(N-1) - 2} \\ &= \frac{2L-2}{N^2 - N - 2} \end{aligned}$$

$$\begin{aligned} p(a_{ij} = 1 | a_{xy} = 0) &= \frac{L}{\frac{N(N-1)}{2} - 1} \\ &= 2 \frac{L}{N(N-1) - 2} \\ &= \frac{2L}{N^2 - N - 2} \end{aligned}$$

2.3

$$\begin{aligned} \frac{P(a_{ij} = 1 | a_{xy} = 1)}{P(a_{ij} = 0)} &= \frac{L-1}{\frac{N(N-1)}{2} - 1} \cdot \frac{\frac{N(N-1)}{2}}{L} \\ &= \frac{L-1}{\frac{N(N-1)-2}{2}} \cdot \frac{N(N-1)}{2L} \\ &= \frac{N(L-1)(N-1)}{\frac{2LN(N-1)-2}{2}} \\ &= \frac{N(L-1)(N-1)}{LN(N-1) - 2} \end{aligned}$$

$$\lim_{N \rightarrow \infty} \frac{N(L-1)(N-1)}{LN(N-1) - 2} = \frac{L-1}{L}$$

$$\begin{aligned}\frac{P(a_{ij} = 1 | a_{xy} = 0)}{P(a_{ij} = 0)} &= \frac{L}{\frac{N(N-1)}{2} - 1} \cdot \frac{\frac{N(N-1)}{2}}{L} \\ &= \frac{2}{N(N-1) - 2} \cdot \frac{N(N-1)}{2} \\ &= \frac{N(N-1)}{N(N-1) - 2}\end{aligned}$$

$$\lim_{N \rightarrow \infty} \frac{N(N-1)}{N(N-1) - 2} = 1$$

2.4

For $G(N, p)$ model, $r'_0 = r'_1 = 1$, in other words, the conditional probabilities don't change and are always the same as $p(a_{ij} = 1)$. This is because the probability p that an edge exists in the $G(N, p)$ model is fixed, unlike in the $G(N, L)$ model.

2.5

3 Degree Correlations and Assortativity




Figure 1: Degree correlation heatmaps




Figure 2: Scatter plots