


# MolProbity Ramachandran analysis

6abu.B99990001.pdb, model 1



85.0% (1140/1332) of all residues were in the favored (90%) regions.  
10.0% (133/1332) of all residues were in the allowed (5.0%) regions.

There were 60 outliers (4.5%).

70.0% (105/150) of all residues were in the favored (90%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.

10.0% (15/150) of all residues were in the allowed (5.0%) regions.