


MolProbity Ramachandran analysis

1yq2.B99990022.pdb, model 1



RM (1993) (1993) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)

1000 (1000) of all residues were in favored (98%) regions.

Of the 1000 (1000) of all residues were in allowed (100%) regions.

There were 11 outliers (98% good)