

This is a testfile for vscode

Ali-loner

2020.08.02

摘要

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Contents

1 This is a section

First, we present a lemma on the Lyapunov function: . Let a system $\dot{x}(t) = f(x(t), x(t-h(t)))$ with $f(0,0) = 0$. Assume the Lyapunov function $F : G \rightarrow \mathbb{R}$ exists with $x, y \in G$, $F(y) < F(x)$ implies

$$\left(\dot{F}(x) f(x, y) \right) \left(\ddot{F}(x) f(x, y) \right) \leq 0. \quad (1)$$

Then the solution $x(t) \equiv 0$ is stable.

Suppose there exists a Lyapunov function $F : \mathbb{R}^n \rightarrow \mathbb{R}$. Then define functional $V : \mathcal{C} \rightarrow \mathbb{R}$ as follows:

$$V(\phi) := \max_{-h \leq \theta \leq 0} F(\phi(\theta)), (\forall \phi \in \mathcal{C}). \quad (2)$$

By definition, the following conditions hold:

$$\dot{V}(\phi) \begin{cases} \leq 0, & \text{if } F(\phi(0)) < V(\phi), \\ = \max \left(\dot{F}(\phi(0)), f(\phi(0), \phi(-h(t))), 0 \right), & \text{if } F(\phi(0)) = V(\phi), \end{cases} \quad (3)$$

where $f(\phi(0), \phi(-h(t))) = \Psi\phi(0) + \Psi_a\phi(-h(t))$.

Thus $\dot{V}(\phi) > 0$ holds if and only if the following condition holds:

$$F(\phi(0)) = \max_{-h \leq \theta \leq 0} F(\phi(\theta)) \text{ and } (\dot{F}(\phi(0)), f(\phi(0), \phi(-h(t)))) > 0. \quad (4)$$

The function F can be defined in some neighborhood $G \subset \mathbb{R}^n$. Moreover, the functional V is then defined for $\phi \in \mathcal{C}$ with values in G .

Suppose Equation (??) holds for some functions $\phi \in \mathcal{C}$, then we can obtain the inequality $F(\phi(-h(t))) < F(\phi(0))$ making ϕ arbitrarily small. Thus the second condition in Equation (??) still holds but conflicts with Lemma ???. Therefore $\dot{V}(\phi) \leq 0$ holds constantly for all ϕ .




图 1: this is Ali

123123
123123
12312
123123
123123
123123
123123
12312
123123
123123
123123
123123
123123
12312
123123

这句话是测试能否进行引用及支持中文^[?]。