

Assignment 4

Rubeena Aafreen

Gradient Descent

Abstract—This document contains the solution to find the maximum / minimum value of given function by gradient descent method

Download all python codes from

<https://github.com/rubeenaafreen20/EE5600AI-ML/tree/master/Assignment4/Code>

Download latex-tikz codes from

<https://github.com/rubeenaafreen20/EE5600AI-ML/tree/master/Assignment4>

minimization part of gradient descent.

Assume,

$$\lambda_0 = 1 \quad (2.0.4)$$

$$\mu = 0.001 \quad (2.0.5)$$

$$precision = 0.00000001 \quad (2.0.6)$$

$$\Rightarrow \lambda_1 = 1 - 0.001(2 \times 9 \times 1 + 12) \quad (2.0.7)$$

$$\Rightarrow \lambda_1 = 1 - 0.03 \quad (2.0.8)$$

$$= 0.97 \quad (2.0.9)$$

following the above method, we keep doing iterations until $\lambda_{n+1} - \lambda_n$ becomes less than the value of precision we have chosen.

3 RESULTS

Using python, the results are:

- 1) The local minimum occurs at - 0.666666130125316.
- 2) The value of $f(x)$ at minima is - 1.999999999974087

Figure 1 shows plot of parabola obtained from python code:




Fig. 1: Plot obtained from python code

1 PROBLEM

Find the maximum and minimum values, if any, of the following function

$$f(x) = 9x^2 + 12x + 2 \quad (1.0.1)$$

2 SOLUTION

From figure 1, the line segment between any two points A and B on the graph of the real-valued function lies above the graph between the two points.

Therefore, the function is convex.

For a general quadratic equation,

$$f(x) = ax^2 + bx + c \quad (2.0.1)$$

The update equation for gradient descent to find minimum of a function is given by:

$$\lambda_{n+1} = \lambda_n - \mu f'(\lambda_n) \quad (2.0.2)$$

$$= \lambda_n - \mu(2a\lambda_n + b) \quad (2.0.3)$$

In equation (2.0.2) λ_0 is an initial guess and μ is a variable parameter, known as step size λ_{n+1} is the next position. The minus sign refers to the