

Fungal and Bacterial Infections Discrimination in ICU Patients based on serum molecular fingerprint

Rúben Araújo^{1,2}, Luís Ramalheira^{3,4}, Tiago Fonseca², Cristiana Von Rekowski², Luís Bento^{1,5,6}, Cecília R.C. Calado²

¹ Comprehensive Health Research Centre, NOVA Medical School, Lisboa, Portugal.

² Laboratory of Health and Engineering, Instituto Superior de Engenharia de Lisboa, Portugal

³ Blood and Transplantation Center of Lisbon, Instituto Português do Sangue e da Transplantação, Alameda das Linhas de Torres, nº 117, 1769 – 001 Lisboa, Portugal;

⁴ iNOVA4Health - Advancing Precision Medicine, RG11: Reno-Vascular Diseases Group, NOVA Medical School, Faculdade de Ciências Médicas, Universidade NOVA de Lisboa, 1169-056 Lisbon, Portugal;

⁵ Intensive Care Department, Centro Hospitalar Universitário de Lisboa Central, CHULC; Rua José António Serrano, 1150-199 Lisboa, Portugal.

⁶ Integrated Pathophysiological Mechanisms, CHRC, NOVA Medical School, Faculdade de Ciências Médicas, NMS, FCM, Universidade NOVA de Lisboa; Campo Mártires da Pátria, 130, 1169-056 Lisboa, Portugal.

ABSTRACT

The rapid discrimination between fungal and bacterial infections in intensive care unit (ICU) patients is crucial for an efficient antimicrobial therapy. This work aims to develop a predictive model capable of discriminating fungal and bacterial infections among these patients based on a rapid serum analysis. Serum from 46 patients at ICU, with COVID-19, were analyzed by Fourier-Transform infrared (FTIR) spectroscopy: 17 did not present any other infection; 12 presented bacterial infections (including *Enterobacter aerogenes* (2), *Escherichia coli* (4), *Enterococcus faecalis* (1), *Haemophilus influenzae* (1) and *Serratia marcescens* (1), *Klebsiella pneumoniae* (3)), isolated from blood, tracheobronchial aspiration, bronchoalveolar lavage and urine; 17 with fungal infections (including *Candida albicans* (14), *C. tropicalis* (1), *C. parapsilosis* (1) and *C. glabrata* (1), isolated from bronchoalveolar lavage, urine and urethral exudate. The impact of a Fast Correlation Based Filter (FCBF) of normalized second derivatized spectra (between 406-1800 and 2800-3992 cm^{-1}) was evaluated on a t-distributed Stochastic Neighbor Embedding (t-SNE), and a Naïve-Bayes model. The predictive models were based on 10 random iterations, with 80% of samples

used for training and the remaining 20% for validation. The impact of FCBF was evaluated on t-SNE to select the spectral bands with high significance in detecting the infection and in discriminating the bacterial from the fungus infection. Optimized Naïve-Bayes models enabled to detect the infection with a sensitivity of 83% and a specificity of 81%, and the discrimination of the bacterial and fungal infection with a sensitivity and specificity of 86% and 96%, respectively. Serum analysis, based on FTIR spectroscopy associated to machine learning algorithms presents a high potential to detect in a rapid and economic mode either the infection either the discrimination between bacterial or fungal infection.

Keywords: FTIR spectroscopy, Intensive Care Unit, fungal infection, bacterial infection, diagnostics

Acknowledgements: This research was funded by grant DSAIPA/DS/0117/2020, *Fundação para a Ciência e a Tecnologia*.