

Lab 2 732A97 Multivariate Statistical Methods

Raymond Sseguya

2019-11-28

Inference about mean vectors

Question 1: Test of outliers

Consider again the data set from the T1-9.dat file, National track records for women. In the first assignment we studied different distance measures between an observation and the sample average vector. The most common multivariate residual is the Mahalanobis distance and we computed this distance for all observations.

a) The Mahalanobis distance is approximately chi-square distributed, if the data comes from a multivariate normal distribution and the number of observations is large. Use this chi-square approximation for testing each observation at the 0.1% significance level and conclude which countries can be regarded as outliers. Should you use a multiple testing correction procedure? Compare the results with and without one. Why is (or maybe is not) 0.1% a sensible significance level for this task?

```
trackrcs <- read.table("T1-9.dat",
  col.names = c("countries", "x100m", "x200m",
    "x400m", "x800m", "x1500m", "x3000m", "marathon"))

trackrcs2 <- (trackrcs)[,-1]
rownames(trackrcs2) <- trackrcs[,1]

C <- cov((trackrcs)[,-1])
x_bar = apply(trackrcs2,1,mean)
d0 = as.matrix(trackrcs2-x_bar)
deviation = sqrt( d0%*%t(d0) )

d_sq_m <- d0%*%solve(C)%*%t(d0)
diagonal_vector3 <- diag(d_sq_m)
deviation_countries3 <-
  cbind.data.frame(countries = as.vector(trackrcs[,1]),diagonal_vector3)
deviation_countries_ordered3 <-
  deviation_countries3[order(-deviation_countries3$diagonal_vector3), ]

named_Mahalanobis <- as.vector(deviation_countries_ordered3[,2])
names(named_Mahalanobis) <- rownames(deviation_countries_ordered3)
ch_s <- combn(x=named_Mahalanobis, m=2,
  FUN = function(c){
```

```

sg <- 0.1/100
pv <- chisq.test(x=c, p=rep(sg,2), rescale.p = TRUE)$p.value
pvname <- paste0(names(c)[1], " ", names(c)[2])
assign(pvname, pv)
return(list(pv, pvname, xx=names(c)[1]))
})

outliers_unsorted <- unlist(ch_s[3, which(ch_s[1,] == 0)])
summary(as.data.frame.character(outliers_unsorted), maxsum = 10)

```

```

## outliers_unsorted
## COK      : 52
## PNG      : 52
## SAM      : 51
## GUA      : 49
## BER      : 48
## MRI      : 44
## CRC      : 43
## DOM      : 43
## MAS      : 43
## (Other) : 632

```

```

print(deviation_countries_ordered3[1:5,])

```

```

##      countries diagonal_vector3
## COK      COK      2094451
## PNG      PNG      2083382
## SAM      SAM      1793652
## GUA      GUA      1485602
## BER      BER      1472580

```

We can see from the summary that “COK”, “PNG”, “SAM”, “GUA” and “BER” remain the top outliers even with using the chi-square test.

possibly using a multiple-testing correction procedure

```

chisq.test(deviation_countries_ordered3$diagonal_vector3,
           p=rep(0.1, nrow(deviation_countries_ordered3)), rescale.p = TRUE )

```

```

##
## Chi-squared test for given probabilities
##
## data: deviation_countries_ordered3$diagonal_vector3
## X-squared = 2034800, df = 53, p-value < 2.2e-16

```

We can see that a multiple-testing correction procedure is clearly a bad idea as it does not actually pinpoint the actual outliers but it tells us only that there are indeed some outliers.

b) One outlier is North Korea. This country is not an outlier with the Euclidean distance. Try to explain these seemingly contradictory results.

It seems in our results, North Korea is an outlier throughout.

Question 2: Test, confidence region and confidence intervals for a mean vector

Look at the bird data in file T5-12.dat and solve Exercise 5:20 of Johnson, Wichern. Do not use any extra R package or built{in test but code all required matrix calculations. You MAY NOT use loops!