

## Reconhecimento de Padrões

### Feature Extraction

Prof. Antonio de Padua Braga

Departamento de Eng. Eletrônica - UFMG

5 de maio de 2011

## Summary

- 1 Introduction
- 2 Principal Component Analysis - PCA
  - Projections
- 3 PCA Examples
- 4 Examples

## Outline

- 1 Introduction
- 2 Principal Component Analysis - PCA
  - Projections
- 3 PCA Examples
- 4 Examples


## Introduction to Feature Extraction/Selection

- A pattern recognition system involves:
- Data aquisition/sensing;
  - Data set conditioning/Pre-processing;
  - Feature extraction;
  - Feature selection;
  - Classification  $\Rightarrow$  Model selection;
  - Post-processing;
  - Decision.
- Input variables (features) for classification should characterize the problem by a set of properties that provide discrimination [ev69].

## Introduction to Feature Extraction/Selection

- Feature Extraction
- Suppose, for instance that, in a face recognition system, our transducer provides color images to the classification system.
  - The goal of feature extraction in this case is to extract relevant features (properties) from the images so that classification can be accomplished.
  - Features to be extracted are not explicitly represented in the images themselves and feature extraction methods are required.
  - Such features could be: face color, hair texture and color, face geometry and so on.

## Introduction to Feature Extraction/Selection




## Introduction to Feature Extraction/Selection

- Feature Extraction
- Suppose, for instance that, in a face recognition system, our transducer provides color images to the classification system.
  - The goal of feature extraction in this case is to extract relevant features (properties) from the images so that classification can be accomplished.
  - Features to be extracted are not explicitly represented in the images themselves and feature extraction methods are required.
  - Such features could be: face color, hair texture and color, face geometry and so on.

## Introduction to Feature Extraction/Selection

### Feature Extraction


Figure below shows an example of two poorly selected features, since they do not provide discrimination:



## Introduction to Feature Extraction/Selection

### Feature Extraction Example


Figure below shows an example of two relevant features, since they provide discrimination:



## Introduction to Feature Extraction/Selection

### Feature Extraction Example

Figure below shows an example of two poorly selected features, since they do not provide discrimination:



## Introduction to Feature Extraction/Selection

Consider the heart diagnosis problem with the following (extracted) features:

- age
  - sex
  - chest pain type (4 values)
  - resting blood pressure
  - serum cholesterol in mg/dl
  - fasting blood sugar > 120 mg/dl
  - resting electrocardiographic results (values 0,1,2)
  - maximum heart rate achieved
  - exercise induced angina
  - oldpeak = ST depression induced by exercise relative to rest
  - the slope of the peak exercise ST segment
  - number of major vessels (0-3) colored by flourosopy
  - thal: 3 = normal; 6 = fixed defect; 7 = reversible defect
- Prof. António da Piedade Braga (Departamento de Engenharia Civil)
- Reconhecimento de Padrões | 5 de maio de 2011 | 10 / 45


## Introduction to Feature Extraction/Selection

### Introduction to Feature Extraction/Selection

- Objective is to classify patients into two classes: heart disease (class 2) and no heart disease (class 1);
- There are 150 examples of class 2 and 120 examples of class 1;
- Feature selection may also identify the most relevant factors associated to heart disease in this particular dataset.









## Projections through $\mathbf{x}_m$

- Figure 3 shows the projections of  $\tau$  on an arbitrary line passing through  $\mathbf{x}_m$ .



## Projections through $\mathbf{x}_m$

- Figure 3 shows the projections of  $\tau$  on an arbitrary line passing through  $\mathbf{x}_m$ .



## Optimal Projection


- By substituting Equation 2 into 1, by obtaining the partial derivative in relation to  $a_i$ , and by setting the result to zero, the expression for the optimal projections  $a_i^*$  are obtained and presented in Equation 3.
- The interpretation of Equation 3 is that the optimal projection  $a_i^*$  corresponds to the projection of the difference vector  $\mathbf{x}_i - \mathbf{x}_m$  on the line passing through  $\mathbf{x}_m$ .
- We need to know now **what is the direction of  $\mathbf{u}$** .

$$a_i^* = \mathbf{u}^T (\mathbf{x}_i - \mathbf{x}_m) \quad (3)$$


- The interpretation of Equation 3 is that the optimal projection  $a_i^*$  corresponds to the projection of the difference vector  $\mathbf{x}_i - \mathbf{x}_m$  on the line passing through  $\mathbf{x}_m$ .
- We need to know now **what is the direction of  $\mathbf{u}$** .

## Removing the mean vector


This is equivalent to solving the problem with  $\mathbf{x}_m$  removed from the dataset:



Which is the line that maximizes the projections?




Which is the line that maximizes the maximum variance of the projections?



The third line yields the maximum variance of the projections.

## Finding the optimal $\mathbf{u}$ through $\mathbf{x}_m$



The third line yields the maximum variance of the projections.

## Finding the optimal $\mathbf{u}$ through $\mathbf{x}_m$



The third line yields the maximum variance of the projections.

The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.



The third line yields the maximum variance of the projections.




The third line yields the maximum variance of the projections.



## Optimal line

Optimal projection:



- By substituting Equation 3 into 1, Equation 4 is obtained.

$$J(u) = -\mathbf{u}^T \mathbf{S} \mathbf{u} + \sum_{i=1}^N (\mathbf{x}_i - \mathbf{x}_m)^2 \quad (4)$$

where  $\mathbf{S}$  is the  $n \times n$  covariance matrix;

$\mathbf{S}$  is  $n \times n$ , where  $n$  in the vector's space dimension;

- We can conclude also that  $\mathbf{u}^T \mathbf{S} \mathbf{u} = \lambda \mathbf{u}^T \mathbf{u} = \lambda$ . So, in order to **maximize**  $\mathbf{u}^T \mathbf{S} \mathbf{u}$  we need to **maximize**  $\lambda$  or, in other words, choose  $\mathbf{u}$  in the direction of the eigenvector with the largest  $\lambda$ .

## Optimal Projection

- By substituting Equation 3 into 1, Equation 4 is obtained.

$$Q(u) = \mathbf{u}^T \mathbf{S} \mathbf{u} - \lambda(\mathbf{u}^T \mathbf{u} - 1) \quad (5)$$

$$\mathbf{S} \mathbf{u} = \lambda \mathbf{u} \quad (6)$$

what implies that  $\mathbf{u}$  is an eigenvector of  $\mathbf{S}$ .

- We can conclude also that  $\mathbf{u}^T \mathbf{S} \mathbf{u} = \lambda \mathbf{u}^T \mathbf{u} = \lambda$ . So, in order to **maximize**  $\mathbf{u}^T \mathbf{S} \mathbf{u}$  we need to **maximize**  $\lambda$  or, in other words, choose  $\mathbf{u}$  in the direction of the eigenvector with the largest  $\lambda$ .

## Principal Components

- The vector  $\mathbf{u}$  that minimizes 4 is the one that maximizes  $\mathbf{u}^T \mathbf{S} \mathbf{u}$ , so the Lagrange Multipliers' solution of Equation 5 is given in Equation 6.

$$Q(u) = \mathbf{u}^T \mathbf{S} \mathbf{u} - \lambda(\mathbf{u}^T \mathbf{u} - 1) \quad (5)$$


$$\mathbf{S} \mathbf{u} = \lambda \mathbf{u} \quad (6)$$

what implies that  $\mathbf{u}$  is an eigenvector of  $\mathbf{S}$ .

- We can conclude also that  $\mathbf{u}^T \mathbf{S} \mathbf{u} = \lambda \mathbf{u}^T \mathbf{u} = \lambda$ . So, in order to **maximize**  $\mathbf{u}^T \mathbf{S} \mathbf{u}$  we need to **maximize**  $\lambda$  or, in other words, choose  $\mathbf{u}$  in the direction of the eigenvector with the largest  $\lambda$ .


## Meaning of PCA

- Data matrix  $\mathbf{X}$  yields the covariance matrix  $\mathbf{S}$  after the mean vector  $\mathbf{x}_m$  is removed from all vectors  $\mathbf{x}_i$  of the dataset.
- The elements  $s_{ij} \in \mathbf{S}$  are obtained as the dot-product between rows  $i$  and  $j$  of the data matrix  $\mathbf{X}$ .




## Meaning of PCA

- Data matrix  $\mathbf{X}$  yields the covariance matrix  $\mathbf{S}$  after the mean vector  $\mathbf{x}_m$  is removed from all vectors  $\mathbf{x}_i$  of the dataset.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.




## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.



## Meaning of PCA

- PCA is an **eigenvalue problem** on the covariance matrix  $\mathbf{S}$ .
  - Covariance matrix  $\mathbf{S}$  contains information about how variables are related.



## Outline

- 1 Introduction
- 2 Principal Component Analysis - PCA
  - Projections
- 3 PCA Examples
  - Examples

Prof. António de Padua Braga (Departamento de Matemática da Universidade de Braga) Reconhecimento de Padrões 5 de maio de 2011 37 / 46

## Outline

- 1 Introduction
- 2 Principal Component Analysis - PCA
  - Projections
- 3 PCA Examples
  - Examples

Prof. António de Padua Braga (Departamento de Matemática da Universidade de Braga) Reconhecimento de Padrões 5 de maio de 2011 38 / 46

## Stepwise Example

Prof. António de Padua Braga (Departamento de Matemática da Universidade de Braga) Reconhecimento de Padrões 5 de maio de 2011 39 / 46

## Stepwise Example

The PCA projected data  $\mathbf{V}\mathbf{X}$ , where  $\mathbf{V}$  is the eigenvector's matrix of  $\mathbf{S}$ , is:

$$\mathbf{V}\mathbf{X} = \begin{bmatrix} 0.1567 & -0.0961 & 0.0842 & 0.2956 & -0.3971 & 0.0992 \\ 3.00442 & 3.0039 & 2.8307 & 6.3712 & 5.6500 & 5.5893 \end{bmatrix}$$

Prof. António de Padua Braga (Departamento de Matemática da Universidade de Braga) Reconhecimento de Padrões 5 de maio de 2011 40 / 46

## Stepwise Example

By removing  $\mathbf{x}_m$  from  $\mathbf{X}$  the following matrix is obtained:

$$\mathbf{X} - [\mathbf{x}_m] = \begin{bmatrix} -1.0417 & -0.8917 & -1.1417 & 1.1583 & 1.1583 & 0.7583 \\ -0.9000 & -1.1000 & -1.1000 & 1.6000 & 0.6000 & 0.9000 \end{bmatrix}$$

that results on the following (non-normalized) matrix  $\mathbf{S}$ :

$$\mathbf{S} = \begin{bmatrix} 6.4421 & 6.4050 \\ 6.4050 & 6.9600 \end{bmatrix}$$

that has the following eigenvalues:

$$\lambda = \begin{bmatrix} 0.2908 \\ 13.1113 \end{bmatrix}$$

indicating that the second axis contains most of the variance of the data.

Prof. António de Padua Braga (Departamento de Matemática da Universidade de Braga) Reconhecimento de Padrões 5 de maio de 2011 41 / 46

## Stepwise Example

The data matrix is:

$$\mathbf{X} = \begin{bmatrix} 2.0000 & 2.1500 & 1.9000 & 4.2000 & 4.2000 & 3.8000 \\ 2.3000 & 2.1000 & 2.1000 & 4.8000 & 3.8000 & 4.1000 \end{bmatrix}$$

and the corresponding mean vector is:

$$\mathbf{x}_m = \begin{bmatrix} 3.0117 \\ 3.2000 \end{bmatrix}$$

Prof. António de Padua Braga (Departamento de Matemática da Universidade de Braga) Reconhecimento de Padrões 5 de maio de 2011 42 / 46

## Stepwise Example

### Breast Cancer Data


- The 2D PCA projections obtained for the CAPES-COFFECUB dataset are presented below [BHN<sup>+</sup>08].



The dataset can now be represented by the largest eigenvector projection

$$\begin{bmatrix} 3.0442 & 3.0039 & 2.8307 & \mathbf{6.3712} & 5.6500 & 5.5893 \end{bmatrix}$$

by removing one of the projections.



**P. Baldi and K. Hornik.**  
Neural networks and principal components analysis: learning from examples without local minima.  
*Neural Networks*, 2:53–58, 1989.

**A. P. Braga, E. G. Horta, R. Natowicz, R. Rouzier, R. Incitti, T. S. Rodrigues, M. A. Costa, C. D. M. Patraro, and A. Cela.**  
Bayesian classifiers for predicting the outcome of breast cancer preoperative chemotherapy.  
In *ANNPR*, pages 263–266, 2008.

**M.D. Levine.**  
Feature extraction: a survey.  
*Proceedings of the IEEE*, 57(9):1391–1407, 1969.

**B Schölkopf, A Smola, and KR Müller.**  
Nonlinear component analysis as a kernel eigenvalue problem.  
*Neural computation*, 1998.