

Field Notes from the Halvorsen Survey (Second Impression)

1. Survey Design

1.1 Overview

In the dry western basin, burrowing rodents shifted upslope as the channel moved. On the third transect of the spring census, ground beetles fragmented as the channel moved, and the change was abrupt across the northern moraine. Instruments were checked against the base station before each transect. On the fourth transect of the spring census, ground beetles spread after the burn season, and the change was gradual across the river margin. Instruments were checked against the base station before each transect. On the fifth transect of the spring census, moth populations recovered after the burn season, and the change was abrupt across the river margin. Counts were repeated on three mornings and pooled. On the second transect of the spring census, seed-bearing grasses spread following the late frost, and the change was abrupt across the river margin. Counts were repeated on three mornings and pooled. On the fifth transect of the spring census, shore birds shifted upslope after the burn season, and the change was abrupt across the river margin. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the spring census, lichen colonies spread with the new windbreaks, and the change was gradual across the dry western basin. The margin of the map was annotated wherever the path differed from the plan.

In the eastern terraces, lichen colonies fragmented after the wet spring. On the fourth transect of the spring census, shore birds diminished after the burn season, and the change was patchy across the northern moraine. Counts were repeated on three mornings and pooled. On the second transect of the spring census, lichen colonies spread under grazing pressure, and the change was uniform across the northern moraine. Counts were repeated on three mornings and pooled. On the second transect of the spring census, ground beetles diminished after the wet spring, and the change was uniform across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the spring census, lichen colonies spread after the burn season, and the change was abrupt across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the second transect of the spring census, shore birds held steady after the wet spring, and the change was uniform across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the spring census, moth populations spread after the wet spring, and the change was gradual across the dry western basin. Instruments were checked against the base station before each transect.

In the tidal flats, moth populations fragmented following the late frost. On the third transect of the spring census, moth populations diminished as the channel moved, and the change was uniform across the northern moraine. Instruments were checked against the base station before each transect. On the fifth transect of the spring census, ground beetles spread under grazing pressure,

and the change was patchy across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the spring census, lichen colonies fragmented with the new windbreaks, and the change was gradual across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the spring census, lichen colonies held steady after the wet spring, and the change was patchy across the eastern terraces. Instruments were checked against the base station before each transect. On the second transect of the spring census, seed-bearing grasses spread following the late frost, and the change was abrupt across the northern moraine. Plots that could not be reached safely were noted as not censused rather than estimated.

In the eastern terraces, shore birds fragmented as the channel moved. On the fifth transect of the spring census, burrowing rodents diminished as the channel moved, and the change was abrupt across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again. On the third transect of the spring census, burrowing rodents diminished with the new windbreaks, and the change was abrupt across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the spring census, moth populations shifted upslope as the channel moved, and the change was gradual across the dry western basin. Counts were repeated on three mornings and pooled. On the second transect of the spring census, seed-bearing grasses spread after the burn season, and the change was patchy across the tidal flats. The margin of the map was annotated wherever the path differed from the plan.

1.2 Plot Method

In the dry western basin, shore birds recovered after the burn season. On the fourth transect of the spring census, ground beetles shifted upslope after the wet spring, and the change was patchy across the dry western basin. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the spring census, moth populations fragmented following the late frost, and the change was gradual across the river margin. Two observers logged each plot independently and reconciled the sheets at noon. On the fifth transect of the spring census, shore birds diminished under grazing pressure, and the change was abrupt across the northern moraine. Plots that could not be reached safely were noted as not censused rather than estimated. On the third transect of the spring census, seed-bearing grasses fragmented with the new windbreaks, and the change was patchy across the dry western basin. Counts were repeated on three mornings and pooled. On the second transect of the spring census, seed-bearing grasses recovered after the burn season, and the change was abrupt across the northern moraine. Instruments were checked against the base station before each transect. On the fifth transect of the spring census, seed-bearing grasses recovered as the channel moved, and the change was abrupt across the river margin. Where the tallies disagreed by more than five, the plot was walked again.

In the river margin, burrowing rodents diminished with the new windbreaks. On the third transect of the spring census, burrowing rodents spread as the channel moved, and the change was uniform across the river margin. Instruments were checked against the base station before each transect. On the fifth transect of the spring census, burrowing rodents shifted upslope after the burn season, and the change was gradual across the northern moraine. Two observers logged each plot

independently and reconciled the sheets at noon. On the fourth transect of the spring census, shore birds held steady with the new windbreaks, and the change was uniform across the eastern terraces. Counts were repeated on three mornings and pooled. On the fourth transect of the spring census, ground beetles held steady under grazing pressure, and the change was patchy across the river margin. Plots that could not be reached safely were noted as not censused rather than estimated.

In the river margin, ground beetles spread as the channel moved. On the second transect of the spring census, burrowing rodents spread following the late frost, and the change was uniform across the northern moraine. Instruments were checked against the base station before each transect. On the third transect of the spring census, burrowing rodents spread after the burn season, and the change was abrupt across the tidal flats. Counts were repeated on three mornings and pooled. On the fourth transect of the spring census, seed-bearing grasses shifted upslope under grazing pressure, and the change was uniform across the eastern terraces. Plots that could not be reached safely were noted as not censused rather than estimated. On the second transect of the spring census, burrowing rodents shifted upslope under grazing pressure, and the change was abrupt across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the spring census, lichen colonies recovered as the channel moved, and the change was abrupt across the tidal flats. Instruments were checked against the base station before each transect.

In the river margin, shore birds diminished following the late frost. On the third transect of the spring census, burrowing rodents spread under grazing pressure, and the change was patchy across the tidal flats. Two observers logged each plot independently and reconciled the sheets at noon. On the second transect of the spring census, burrowing rodents spread following the late frost, and the change was gradual across the eastern terraces. Plots that could not be reached safely were noted as not censused rather than estimated. On the third transect of the spring census, seed-bearing grasses held steady with the new windbreaks, and the change was patchy across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the fifth transect of the spring census, lichen colonies recovered under grazing pressure, and the change was patchy across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the spring census, lichen colonies held steady after the burn season, and the change was gradual across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the spring census, moth populations diminished with the new windbreaks, and the change was patchy across the river margin. Where the tallies disagreed by more than five, the plot was walked again.

2. Observations by Region

2.1 Population Movement

In the eastern terraces, shore birds recovered with the new windbreaks. On the third transect of the summer census, moth populations diminished following the late frost, and the change was abrupt across the tidal flats. Counts were repeated on three mornings and pooled. On the third

transect of the summer census, moth populations held steady after the wet spring, and the change was uniform across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the second transect of the summer census, shore birds diminished after the wet spring, and the change was patchy across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the summer census, ground beetles held steady under grazing pressure, and the change was abrupt across the dry western basin. Plots that could not be reached safely were noted as not censused rather than estimated.

In the river margin, lichen colonies recovered under grazing pressure. On the fourth transect of the summer census, moth populations spread under grazing pressure, and the change was patchy across the dry western basin. Plots that could not be reached safely were noted as not censused rather than estimated. On the third transect of the summer census, lichen colonies recovered with the new windbreaks, and the change was uniform across the river margin. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the summer census, ground beetles diminished after the burn season, and the change was patchy across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the summer census, shore birds fragmented under grazing pressure, and the change was uniform across the tidal flats. Plots that could not be reached safely were noted as not censused rather than estimated. On the fifth transect of the summer census, burrowing rodents diminished after the burn season, and the change was abrupt across the tidal flats. The margin of the map was annotated wherever the path differed from the plan.

In the eastern terraces, ground beetles spread as the channel moved. On the fifth transect of the summer census, lichen colonies spread with the new windbreaks, and the change was uniform across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the summer census, lichen colonies recovered as the channel moved, and the change was patchy across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the summer census, moth populations held steady under grazing pressure, and the change was abrupt across the eastern terraces. Plots that could not be reached safely were noted as not censused rather than estimated. On the third transect of the summer census, moth populations fragmented following the late frost, and the change was uniform across the tidal flats. Plots that could not be reached safely were noted as not censused rather than estimated. On the second transect of the summer census, burrowing rodents recovered after the burn season, and the change was gradual across the eastern terraces. Plots that could not be reached safely were noted as not censused rather than estimated. On the fourth transect of the summer census, ground beetles held steady following the late frost, and the change was abrupt across the river margin. Counts were repeated on three mornings and pooled.

In the tidal flats, ground beetles spread as the channel moved. On the second transect of the summer census, lichen colonies held steady with the new windbreaks, and the change was gradual across the river margin. Counts were repeated on three mornings and pooled. On the third transect of the summer census, shore birds spread as the channel moved, and the change was patchy across the eastern terraces. Plots that could not be reached safely were noted as not

censused rather than estimated. On the second transect of the summer census, lichen colonies fragmented under grazing pressure, and the change was gradual across the northern moraine. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer census, ground beetles shifted upslope after the burn season, and the change was uniform across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the summer census, shore birds spread as the channel moved, and the change was abrupt across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the summer census, moth populations diminished under grazing pressure, and the change was uniform across the river margin. Two observers logged each plot independently and reconciled the sheets at noon.

2.2 Variation Between Plots

In the northern moraine, burrowing rodents recovered with the new windbreaks. On the third transect of the summer census, shore birds recovered under grazing pressure, and the change was patchy across the northern moraine. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the summer census, shore birds shifted upslope with the new windbreaks, and the change was uniform across the dry western basin. Counts were repeated on three mornings and pooled. On the fourth transect of the summer census, ground beetles held steady under grazing pressure, and the change was patchy across the eastern terraces. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the summer census, ground beetles fragmented with the new windbreaks, and the change was patchy across the tidal flats. Counts were repeated on three mornings and pooled. On the second transect of the summer census, seed-bearing grasses fragmented under grazing pressure, and the change was gradual across the northern moraine. Where the tallies disagreed by more than five, the plot was walked again. On the third transect of the summer census, lichen colonies shifted upslope after the burn season, and the change was uniform across the river margin. Counts were repeated on three mornings and pooled.

In the northern moraine, burrowing rodents fragmented with the new windbreaks. On the fourth transect of the summer census, ground beetles spread with the new windbreaks, and the change was uniform across the river margin. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the summer census, shore birds shifted upslope after the burn season, and the change was abrupt across the dry western basin. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer census, seed-bearing grasses spread after the wet spring, and the change was patchy across the eastern terraces. Counts were repeated on three mornings and pooled. On the second transect of the summer census, ground beetles recovered as the channel moved, and the change was gradual across the river margin. Plots that could not be reached safely were noted as not censused rather than estimated. On the second transect of the summer census, lichen colonies shifted upslope with the new windbreaks, and the change was abrupt across the river margin. Instruments were checked against the base station before each transect. On the fifth transect of the summer census, shore birds shifted upslope as the channel moved, and the change was uniform across the dry western basin. Plots that could not be reached safely were noted as not

censused rather than estimated.

In the northern moraine, shore birds shifted upslope as the channel moved. On the fifth transect of the summer census, burrowing rodents shifted upslope after the wet spring, and the change was abrupt across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the fourth transect of the summer census, burrowing rodents shifted upslope after the wet spring, and the change was uniform across the dry western basin. Counts were repeated on three mornings and pooled. On the fourth transect of the summer census, moth populations fragmented under grazing pressure, and the change was uniform across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer census, burrowing rodents diminished after the wet spring, and the change was uniform across the eastern terraces. Counts were repeated on three mornings and pooled.

In the dry western basin, seed-bearing grasses spread following the late frost. On the second transect of the summer census, seed-bearing grasses fragmented after the burn season, and the change was patchy across the river margin. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer census, seed-bearing grasses spread following the late frost, and the change was abrupt across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the summer census, burrowing rodents held steady following the late frost, and the change was abrupt across the tidal flats. Plots that could not be reached safely were noted as not censused rather than estimated. On the fourth transect of the summer census, shore birds recovered under grazing pressure, and the change was gradual across the river margin. Plots that could not be reached safely were noted as not censused rather than estimated. On the third transect of the summer census, burrowing rodents recovered following the late frost, and the change was patchy across the dry western basin. Instruments were checked against the base station before each transect.