

Field Notes from the Halvorsen Survey

1. Survey Design

1.1 Overview

In the dry western basin, burrowing rodents shifted upslope as the channel moved. On the third transect of the spring survey, ground beetles fragmented as the channel moved, and the change was abrupt across the northern moraine. Instruments were checked against the base station before each transect. On the fourth transect of the spring survey, ground beetles expanded after the burn season, and the change was gradual across the river margin. Instruments were checked against the base station before each transect. On the fifth transect of the spring survey, moth populations recovered after the burn season, and the change was abrupt across the river margin. Counts were repeated on three mornings and averaged. On the second transect of the spring survey, seed-bearing grasses expanded following the late frost, and the change was abrupt across the river margin. Counts were repeated on three mornings and averaged. On the fifth transect of the spring survey, shore birds shifted upslope after the burn season, and the change was abrupt across the river margin. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the spring survey, lichen colonies expanded with the new windbreaks, and the change was gradual across the dry western basin. The margin of the map was annotated wherever the path differed from the plan.

In the eastern terraces, lichen colonies fragmented after the wet spring. On the fourth transect of the spring survey, shore birds declined after the burn season, and the change was patchy across the northern moraine. Counts were repeated on three mornings and averaged. On the second transect of the spring survey, lichen colonies expanded under grazing pressure, and the change was uniform across the northern moraine. Counts were repeated on three mornings and averaged. On the second transect of the spring survey, ground beetles declined after the wet spring, and the change was uniform across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the spring survey, lichen colonies expanded after the burn season, and the change was abrupt across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the second transect of the spring survey, shore birds held steady after the wet spring, and the change was uniform across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the spring survey, moth populations expanded after the wet spring, and the change was gradual across the dry western basin. Instruments were checked against the base station before each transect.

In the tidal flats, moth populations fragmented following the late frost. On the third transect of the spring survey, moth populations declined as the channel moved, and the change was uniform across the northern moraine. Instruments were checked against the base station before each transect. On the fifth transect of the spring survey, ground beetles expanded under grazing pressure, and the change was patchy across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the spring survey,

lichen colonies fragmented with the new windbreaks, and the change was gradual across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the spring survey, lichen colonies held steady after the wet spring, and the change was patchy across the eastern terraces. Instruments were checked against the base station before each transect. On the second transect of the spring survey, seed-bearing grasses expanded following the late frost, and the change was abrupt across the northern moraine. Plots that could not be reached safely were recorded as not surveyed rather than estimated.

In the eastern terraces, shore birds fragmented as the channel moved. On the fifth transect of the spring survey, burrowing rodents declined as the channel moved, and the change was abrupt across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again. On the third transect of the spring survey, burrowing rodents declined with the new windbreaks, and the change was abrupt across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the spring survey, moth populations shifted upslope as the channel moved, and the change was gradual across the dry western basin. Counts were repeated on three mornings and averaged. On the second transect of the spring survey, seed-bearing grasses expanded after the burn season, and the change was patchy across the tidal flats. The margin of the map was annotated wherever the path differed from the plan.




Figure 1: Illustrative population counts by region.

1.2 Plot Method

In the dry western basin, shore birds recovered after the burn season. On the fourth transect of the spring survey, ground beetles shifted upslope after the wet spring, and the change was patchy across the dry western basin. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the spring survey, moth populations fragmented following the late frost, and the change was gradual across the river margin. Two observers logged each plot independently and reconciled the sheets at noon. On the fifth transect of the spring survey, shore birds declined under grazing pressure, and the change was abrupt across the northern moraine. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the third transect of the spring survey, seed-bearing grasses fragmented with the new windbreaks, and the change was patchy across the dry western basin. Counts were repeated on three mornings and averaged. On the second transect of the spring survey, seed-bearing grasses recovered after the burn season, and the change was abrupt across the northern moraine.

Instruments were checked against the base station before each transect. On the fifth transect of the spring survey, seed-bearing grasses recovered as the channel moved, and the change was abrupt across the river margin. Where the tallies disagreed by more than five, the plot was walked again.

In the river margin, burrowing rodents declined with the new windbreaks. On the third transect of the spring survey, burrowing rodents expanded as the channel moved, and the change was uniform across the river margin. Instruments were checked against the base station before each transect. On the fifth transect of the spring survey, burrowing rodents shifted upslope after the burn season, and the change was gradual across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the spring survey, shore birds held steady with the new windbreaks, and the change was uniform across the eastern terraces. Counts were repeated on three mornings and averaged. On the fourth transect of the spring survey, ground beetles held steady under grazing pressure, and the change was patchy across the river margin. Plots that could not be reached safely were recorded as not surveyed rather than estimated.

In the river margin, ground beetles expanded as the channel moved. On the second transect of the spring survey, burrowing rodents expanded following the late frost, and the change was uniform across the northern moraine. Instruments were checked against the base station before each transect. On the third transect of the spring survey, burrowing rodents expanded after the burn season, and the change was abrupt across the tidal flats. Counts were repeated on three mornings and averaged. On the fourth transect of the spring survey, seed-bearing grasses shifted upslope under grazing pressure, and the change was uniform across the eastern terraces. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the second transect of the spring survey, burrowing rodents shifted upslope under grazing pressure, and the change was abrupt across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the spring survey, lichen colonies recovered as the channel moved, and the change was abrupt across the tidal flats. Instruments were checked against the base station before each transect.

In the river margin, shore birds declined following the late frost. On the third transect of the spring survey, burrowing rodents expanded under grazing pressure, and the change was patchy across the tidal flats. Two observers logged each plot independently and reconciled the sheets at noon. On the second transect of the spring survey, burrowing rodents expanded following the late frost, and the change was gradual across the eastern terraces. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the third transect of the spring survey, seed-bearing grasses held steady with the new windbreaks, and the change was patchy across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the fifth transect of the spring survey, lichen colonies recovered under grazing pressure, and the change was patchy across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the spring survey, lichen colonies held steady after the burn season, and the change was gradual across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the spring survey, moth populations declined with the new windbreaks, and the change was

patchy across the river margin. Where the tallies disagreed by more than five, the plot was walked again.

2. Observations by Region

2.1 Population Movement

In the eastern terraces, shore birds recovered with the new windbreaks. On the third transect of the summer survey, moth populations declined following the late frost, and the change was abrupt across the tidal flats. Counts were repeated on three mornings and averaged. On the third transect of the summer survey, moth populations held steady after the wet spring, and the change was uniform across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the second transect of the summer survey, shore birds declined after the wet spring, and the change was patchy across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the summer survey, ground beetles held steady under grazing pressure, and the change was abrupt across the dry western basin. Plots that could not be reached safely were recorded as not surveyed rather than estimated.

In the river margin, lichen colonies recovered under grazing pressure. On the fourth transect of the summer survey, moth populations expanded under grazing pressure, and the change was patchy across the dry western basin. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the third transect of the summer survey, lichen colonies recovered with the new windbreaks, and the change was uniform across the river margin. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the summer survey, ground beetles declined after the burn season, and the change was patchy across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the summer survey, shore birds fragmented under grazing pressure, and the change was uniform across the tidal flats. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the fifth transect of the summer survey, burrowing rodents declined after the burn season, and the change was abrupt across the tidal flats. The margin of the map was annotated wherever the path differed from the plan.

In the eastern terraces, ground beetles expanded as the channel moved. On the fifth transect of the summer survey, lichen colonies expanded with the new windbreaks, and the change was uniform across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the summer survey, lichen colonies recovered as the channel moved, and the change was patchy across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the summer survey, moth populations held steady under grazing pressure, and the change was abrupt across the eastern terraces. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the third transect of the summer survey, moth populations fragmented following the late frost, and the change was uniform across the tidal flats. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the second transect of

the summer survey, burrowing rodents recovered after the burn season, and the change was gradual across the eastern terraces. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the fourth transect of the summer survey, ground beetles held steady following the late frost, and the change was abrupt across the river margin. Counts were repeated on three mornings and averaged.

In the tidal flats, ground beetles expanded as the channel moved. On the second transect of the summer survey, lichen colonies held steady with the new windbreaks, and the change was gradual across the river margin. Counts were repeated on three mornings and averaged. On the third transect of the summer survey, shore birds expanded as the channel moved, and the change was patchy across the eastern terraces. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the second transect of the summer survey, lichen colonies fragmented under grazing pressure, and the change was gradual across the northern moraine. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer survey, ground beetles shifted upslope after the burn season, and the change was uniform across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the summer survey, shore birds expanded as the channel moved, and the change was abrupt across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again. On the fifth transect of the summer survey, moth populations declined under grazing pressure, and the change was uniform across the river margin. Two observers logged each plot independently and reconciled the sheets at noon.

$$\sum_{i=1}^n w_i \cdot \partial f / \partial x_i \approx \int \phi(x) \, dx, \text{ with } \alpha, \beta, \Delta \geq 0 \text{ and } \forall x \in \Omega: |x_i| \leq \varepsilon.$$

2.2 Variation Between Plots

In the northern moraine, burrowing rodents recovered with the new windbreaks. On the third transect of the summer survey, shore birds recovered under grazing pressure, and the change was patchy across the northern moraine. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the summer survey, shore birds shifted upslope with the new windbreaks, and the change was uniform across the dry western basin. Counts were repeated on three mornings and averaged. On the fourth transect of the summer survey, ground beetles held steady under grazing pressure, and the change was patchy across the eastern terraces. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the summer survey, ground beetles fragmented with the new windbreaks, and the change was patchy across the tidal flats. Counts were repeated on three mornings and averaged. On the second transect of the summer survey, seed-bearing grasses fragmented under grazing pressure, and the change was gradual across the northern moraine. Where the tallies disagreed by more than five, the plot was walked again. On the third transect of the summer survey, lichen colonies shifted upslope after the burn season, and the change was uniform across the river margin. Counts were repeated on three mornings and averaged.

In the northern moraine, burrowing rodents fragmented with the new windbreaks. On the fourth transect of the summer survey, ground beetles expanded with the new windbreaks, and the change was uniform across the river margin. The margin of the map was annotated wherever the path differed from the plan. On the third transect of the summer survey, shore birds shifted

upslope after the burn season, and the change was abrupt across the dry western basin. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer survey, seed-bearing grasses expanded after the wet spring, and the change was patchy across the eastern terraces. Counts were repeated on three mornings and averaged. On the second transect of the summer survey, ground beetles recovered as the channel moved, and the change was gradual across the river margin. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the second transect of the summer survey, lichen colonies shifted upslope with the new windbreaks, and the change was abrupt across the river margin. Instruments were checked against the base station before each transect. On the fifth transect of the summer survey, shore birds shifted upslope as the channel moved, and the change was uniform across the dry western basin. Plots that could not be reached safely were recorded as not surveyed rather than estimated.

In the northern moraine, shore birds shifted upslope as the channel moved. On the fifth transect of the summer survey, burrowing rodents shifted upslope after the wet spring, and the change was abrupt across the dry western basin. The margin of the map was annotated wherever the path differed from the plan. On the fourth transect of the summer survey, burrowing rodents shifted upslope after the wet spring, and the change was uniform across the dry western basin. Counts were repeated on three mornings and averaged. On the fourth transect of the summer survey, moth populations fragmented under grazing pressure, and the change was uniform across the eastern terraces. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer survey, burrowing rodents declined after the wet spring, and the change was uniform across the eastern terraces. Counts were repeated on three mornings and averaged.

In the dry western basin, seed-bearing grasses expanded following the late frost. On the second transect of the summer survey, seed-bearing grasses fragmented after the burn season, and the change was patchy across the river margin. Where the tallies disagreed by more than five, the plot was walked again. On the fourth transect of the summer survey, seed-bearing grasses expanded following the late frost, and the change was abrupt across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the summer survey, burrowing rodents held steady following the late frost, and the change was abrupt across the tidal flats. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the fourth transect of the summer survey, shore birds recovered under grazing pressure, and the change was gradual across the river margin. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the third transect of the summer survey, burrowing rodents recovered following the late frost, and the change was patchy across the dry western basin. Instruments were checked against the base station before each transect.

3. Sources of Error

In the tidal flats, seed-bearing grasses fragmented with the new windbreaks. On the third transect of the autumn survey, ground beetles expanded after the wet spring, and the change was abrupt across the eastern terraces. Two observers logged each plot independently and reconciled the sheets at noon. On the second transect of the autumn survey, shore birds shifted upslope with the

new windbreaks, and the change was patchy across the eastern terraces. Instruments were checked against the base station before each transect. On the fourth transect of the autumn survey, shore birds declined following the late frost, and the change was patchy across the eastern terraces. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the autumn survey, moth populations fragmented as the channel moved, and the change was abrupt across the river margin. Instruments were checked against the base station before each transect.

In the dry western basin, lichen colonies expanded as the channel moved. On the fourth transect of the autumn survey, lichen colonies fragmented under grazing pressure, and the change was uniform across the dry western basin. Counts were repeated on three mornings and averaged. On the fifth transect of the autumn survey, seed-bearing grasses declined following the late frost, and the change was abrupt across the river margin. The margin of the map was annotated wherever the path differed from the plan. On the second transect of the autumn survey, shore birds fragmented as the channel moved, and the change was abrupt across the northern moraine. Counts were repeated on three mornings and averaged. On the fifth transect of the autumn survey, seed-bearing grasses expanded as the channel moved, and the change was patchy across the dry western basin. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the third transect of the autumn survey, ground beetles fragmented after the burn season, and the change was patchy across the northern moraine. Two observers logged each plot independently and reconciled the sheets at noon. On the fourth transect of the autumn survey, shore birds declined with the new windbreaks, and the change was patchy across the dry western basin. Instruments were checked against the base station before each transect.

In the eastern terraces, lichen colonies expanded under grazing pressure. On the third transect of the autumn survey, seed-bearing grasses recovered as the channel moved, and the change was uniform across the dry western basin. Counts were repeated on three mornings and averaged. On the fourth transect of the autumn survey, moth populations expanded under grazing pressure, and the change was gradual across the northern moraine. Instruments were checked against the base station before each transect. On the second transect of the autumn survey, shore birds declined as the channel moved, and the change was uniform across the dry western basin. Counts were repeated on three mornings and averaged. On the second transect of the autumn survey, lichen colonies expanded after the burn season, and the change was uniform across the tidal flats. Plots that could not be reached safely were recorded as not surveyed rather than estimated. On the fifth transect of the autumn survey, moth populations recovered under grazing pressure, and the change was gradual across the river margin. Counts were repeated on three mornings and averaged.

In the northern moraine, shore birds recovered as the channel moved. On the fourth transect of the autumn survey, seed-bearing grasses fragmented with the new windbreaks, and the change was patchy across the river margin. Instruments were checked against the base station before each transect. On the fourth transect of the autumn survey, burrowing rodents expanded as the channel moved, and the change was uniform across the northern moraine. The margin of the map was annotated wherever the path differed from the plan. On the fourth transect of the autumn

survey, burrowing rodents shifted upslope after the burn season, and the change was gradual across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again. On the third transect of the autumn survey, lichen colonies fragmented under grazing pressure, and the change was patchy across the tidal flats. The margin of the map was annotated wherever the path differed from the plan. On the fourth transect of the autumn survey, lichen colonies held steady after the wet spring, and the change was patchy across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again.

In the northern moraine, ground beetles recovered after the burn season. On the second transect of the autumn survey, moth populations fragmented as the channel moved, and the change was patchy across the tidal flats. Where the tallies disagreed by more than five, the plot was walked again. On the second transect of the autumn survey, shore birds held steady as the channel moved, and the change was uniform across the river margin. Counts were repeated on three mornings and averaged. On the fifth transect of the autumn survey, burrowing rodents expanded after the wet spring, and the change was abrupt across the dry western basin. Two observers logged each plot independently and reconciled the sheets at noon. On the third transect of the autumn survey, burrowing rodents fragmented after the burn season, and the change was patchy across the eastern terraces. Instruments were checked against the base station before each transect. On the third transect of the autumn survey, burrowing rodents recovered after the burn season, and the change was gradual across the river margin. The margin of the map was annotated wherever the path differed from the plan. On the fifth transect of the autumn survey, burrowing rodents declined with the new windbreaks, and the change was patchy across the dry western basin. Plots that could not be reached safely were recorded as not surveyed rather than estimated.

Table 1: Comparative observations.

Region	Species observed	Variants
North	128	17
South	94	9
East	142	21
West	63	5