

IIT MATHEMATICS

SEQUENCE & SERIES

WORKBOOK



RESULTS OF BEST MENTORSHIP BY THE NUCLEUS TEAM



AIR-1
CHITRAANG MURDIA
Gen. Category - 2014



AIR-3
GOVIND LAHOTI
Gen. Category - 2014



AIR-6
NISHIT AGARWAL
Gen. Category - 2012



AIR-8
AMEY GUPTA
Gen. Category - 2014

JEE MAIN RESULTS 2020 (January Attempt) OF NUCLEUS EDUCATION



100 Percentile
(Physics)
DAKSH KHANDELWAL
2020



100 Percentile
(Maths & Physics)
VAIBHAV SAHA
2020



100 Percentile
(Physics)
ANISH MOHAN
2020



100 Percentile
(Maths)
ARCHIT PATNAIK
2020




100 Percentile
(Maths)
SWAPNIL YASAVI
2020



100 Percentile
(Maths)
PARSHANT ARORA
2020


JEE ADVANCED RESULTS OF NUCLEUS EDUCATION




AIR-10
LAKSHAY SHARMA
Gen. Category - 2017




AIR-12
YATEESH AGRAWAL
Gen. Category - 2017




AIR-23
ABHEY GOYAL
Gen. Category - 2017




AIR-24
TUSHAR GAUTAM
Gen. Category - 2017




AIR-37
PIYUSH TIBAREWAL
Gen. Category - 2017




AIR-42
SATVIK
Gen. Category - 2017



AIR-66
MAYANK DUBEY
Gen. Category - 2017



AIR-98
HRITHIK
Gen. Category - 2017



AIR-20
SHASHANK AGRAWAL
Gen. Category - 2018



AIR-27
RAAGHAV RAAJ
Gen. Category - 2018




AIR-32
SHREYA PATHAK
Gen. Category - 2018



AIR-61
SIDDHANT CHOURAHY
Gen. Category - 2018



AIR-67
ANISWAR S KRISHNAN
Gen. Category - 2018
DLP



AIR-78
AAYUSH KADAM
Gen. Category - 2018




AIR-61
SARTHAK BEHERA
Gen. Category - 2018
*SDCCP



AIR-91
ANDREWS G. VARGHESE
Gen. Category - 2018
*SDCCP



AIR-2
HIMANSHU GAURAV SINGH
Gen. Category - 2019
*SDCCP



AIR-19
VIBHAV AGGARWAL
Gen. Category - 2019



AIR-33
S. PRAJEETH
Gen. Category - 2019
*SDCCP



AIR-48
SOHAM MISTRI
Gen. Category - 2019
DLP



AIR-51
SAYANTAN DHAR
Gen. Category - 2019
DLP



AIR-53
GAURAV KRISHAN GUPTA
Gen. Category - 2019
DLP



AIR-86
SATVIK JAIN
Gen. Category - 2019

SEQUENCE & SERIES

1. DEFINITION :

Sequence : A succession of terms a_1, a_2, a_3, \dots formed according to some rule or law.

Examples are : 1, 4, 9, 16, 25
 $-1, 1, -1, 1, \dots$

$$\frac{x}{1!}, \frac{x^2}{2!}, \frac{x^3}{3!}, \frac{x^4}{4!}, \dots$$

A finite sequence has a finite (i.e. limited) number of terms, as in the first example above. An infinite sequence has an unlimited number of terms, i.e. there is no last term, as in the second and third examples.

Series : The indicated sum of the terms of a sequence. In the case of a finite sequence $a_1, a_2, a_3, \dots, a_n$, the corresponding series is $a_1 + a_2 + a_3 + \dots + a_n = \sum_{k=1}^n a_k$. This series has a

finite or limited number of terms and is called a finite series.

2. ARITHMETIC PROGRESSION (A.P.) :

A.P. is a sequence whose terms differ by a fixed number. This fixed number is called the common difference. If a is the first term & d the common difference, then A.P. can be written as

$a, a+d, a+2d, \dots, a+(n-1)d, \dots$

(a) n^{th} term of AP $T_n = a + (n-1)d$, where $d = t_n - t_{n-1}$

(b) The sum of the first n terms : $S_n = \frac{n}{2}$

$[a+\ell] = \frac{n}{2} [2a + (n-1)d]$ where ℓ is n^{th} term.

Note :

(i) n^{th} term of an A.P. is of the form $An + B$ i.e. a linear expression in ' n ', in such a case the coefficient of n is the common difference of the A.P. i.e. A .

- (ii) Sum of first ' n ' terms of an A.P. is of the form $An^2 + Bn$ i.e. a quadratic expression in ' n ', in such case the common difference is twice the coefficient of n^2 . i.e. $2A$
- (iii) Also n^{th} term $T_n = S_n - S_{n-1}$

Illustration 1 :

If $(x+1), 3x$ and $(4x+2)$ are first three terms of an A.P. then its 5^{th} term is –

- (A) 14 (B) 19
 (C) 24 (D) 28

Solution :

$$\begin{aligned} & (x+1), 3x, (4x+2) \text{ are in AP} \\ \Rightarrow & 3x - (x+1) = (4x+2) - 3x \\ \Rightarrow & x = 3 \\ \therefore & a = 4, d = 9 - 4 = 5 \\ \Rightarrow & T_5 = 4 + (4)5 = 24 \quad \text{Ans. (C)} \end{aligned}$$

Illustration 2 :

The sum of first four terms of an A.P. is 56 and the sum of its last four terms is 112. If its first term is 11 then find the number of terms in the A.P.

Solution :

$$\begin{aligned} & a + a + d + a + 2d + a + 3d = 56 \\ \Rightarrow & 4a + 6d = 56 \\ \Rightarrow & 44 + 6d = 56 \quad (\text{as } a = 11) \\ \Rightarrow & 6d = 12 \\ \Rightarrow & \text{hence } d = 2 \\ & \text{Let total number of terms} = n \\ & \text{Now sum of last four terms.} \\ & a + (n-1)d + a + (n-2)d + a + (n-3)d \\ & + a + (n-4)d = 112 \\ \Rightarrow & 4a + (4n-10)d = 112 \\ \Rightarrow & 44 + (4n-10)2 = 112 \\ \Rightarrow & 4n - 10 = 34 \\ \Rightarrow & n = 11 \quad \text{Ans.} \end{aligned}$$

Illustration 3 :

The sum of first n terms of two A.P.s. are in ratio $\frac{7n+1}{4n+27}$. Find the ratio of their 11^{th} terms.

Solution :

Let a_1 and a_2 be the first terms and d_1 and d_2 be the common differences of two A.P.s respectively then

$$\frac{\frac{n}{2}[2a_1 + (n-1)d_1]}{\frac{n}{2}[2a_2 + (n-1)d_2]} = \frac{7n+1}{4n+27}$$

$$\Rightarrow \frac{a_1 + \left(\frac{n-1}{2}\right)d_1}{a_2 + \left(\frac{n-1}{2}\right)d_2} = \frac{7n+1}{4n+27}$$

For ratio of 11th terms

$$\frac{n-1}{2} = 10 \Rightarrow n = 21$$

so ratio of 11th terms is

$$\frac{7(21)+1}{4(21)+27} = \frac{148}{111} = \frac{4}{3} \quad \text{Ans.}$$

Do yourself - 1 :

- (i) Write down the sequence whose n th terms is :

(a) $\frac{2^n}{n}$ (b) $\frac{3+(-1)^n}{3^n}$

- (ii) For an A.P, show that $t_m + t_{2n+m} = 2t_{m+n}$
 (iii) If the sum of p terms of an A.P. is q and the sum of its q terms is p , then find the sum of its $(p+q)$ term.

3. PROPERTIES OF A.P. :

- (a) If each term of an A.P. is increased, decreased, multiplied or divided by the some nonzero number, then the resulting sequence is also an A.P.

(b) Three numbers in A.P. :

$$a-d, a, a+d$$

Four numbers in A.P. :

$$a-3d, a-d, a+d, a+3d$$

Five numbers in A.P. :

$$a-2d, a-d, a, a+d, a+2d$$

Six numbers in A.P. :

$$a-5d, a-3d, a-d, a+d, a+3d, a+5d \text{ etc.}$$

- (c) The common difference can be zero, positive or negative.
- (d) k^{th} term from the last = $(n-k+1)^{\text{th}}$ term from the beginning (If total number of terms = n).
- (e) The sum of the two terms of an AP equidistant from the beginning & end is constant and equal to the sum of first & last terms. $\Rightarrow T_k + T_{n-k+1} = \text{constant} = a + \ell$.
- (f) Any term of an AP (except the first) is equal to half the sum of terms which are equidistant from it. $a_n = (1/2)(a_{n-k} + a_{n+k})$, $k < n$
 For $k = 1$, $a_n = (1/2)(a_{n-k} + a_{n+k})$; For $k = 2$, $a_n = (1/2)(a_{n-2} + a_{n+2})$ and so on.
- (g) If a, b, c are in AP, then $2b = a + c$.

Illustration 4 :

Four numbers are in A.P. If their sum is 20 and the sum of their squares is 120, then the middle terms are -

- (A) 2, 4 (B) 4, 6
 (C) 6, 8 (D) 8, 10

Solution :

Let the numbers are $a-3d, a-d, a+d, a+3d$

given, $a-3d + a-d + a+d + a+3d = 20$

$$\begin{aligned} \Rightarrow 4a = 20 &\Rightarrow a = 5 \\ \text{and } (a-3d)^2 + (a-d)^2 + (a+d)^2 + (a+3d)^2 &= 120 \\ \Rightarrow 4a^2 + 20d^2 &= 120 \\ \Rightarrow 4 \times 5^2 + 20d^2 &= 120 \\ \Rightarrow d^2 &= 1 \Rightarrow d = \pm 1 \\ \text{Hence numbers are} \\ 2, 4, 6, 8 \text{ or } 8, 6, 4, 2 &\quad \text{Ans. (B)} \end{aligned}$$

Illustration 5 :

If $a_1, a_2, a_3, \dots, a_n$ are in A.P. where $a_i > 0$ for all i , show that :

$$\begin{aligned} \frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \\ \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} &= \frac{(n-1)}{\sqrt{a_1} + \sqrt{a_n}} \end{aligned}$$

Solution :

$$\begin{aligned}
 \text{L.H.S.} &= \frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \\
 &\dots\dots\dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} \\
 &= \frac{1}{\sqrt{a_2} + \sqrt{a_1}} + \frac{1}{\sqrt{a_3} + \sqrt{a_2}} + \dots\dots\dots + \\
 &\frac{1}{\sqrt{a_n} + \sqrt{a_{n-1}}} \\
 &= \frac{\sqrt{a_2} - \sqrt{a_1}}{(a_2 - a_1)} + \frac{\sqrt{a_3} - \sqrt{a_2}}{(a_3 - a_2)} + \dots\dots\dots + \\
 &\frac{\sqrt{a_n} - \sqrt{a_{n-1}}}{a_n - a_{n-1}}
 \end{aligned}$$

Let 'd' is the common difference of this A.P.

then $a_2 - a_1 = a_3 - a_2 = \dots\dots\dots = a_n - a_{n-1} = d$

Now L.H.S.

$$\begin{aligned}
 &= \frac{1}{d} \left\{ \sqrt{a_2} - \sqrt{a_1} + \sqrt{a_3} - \sqrt{a_2} + \dots\dots\dots + \right. \\
 &\quad \left. \sqrt{a_{n-1}} - \sqrt{a_{n-2}} + \sqrt{a_n} - \sqrt{a_{n-1}} \right\} = \frac{1}{d} \\
 &\quad \quad \quad \left\{ \sqrt{a_n} - \sqrt{a_1} \right\} \\
 &= \frac{a_n - a_1}{d(\sqrt{a_n} + \sqrt{a_1})} = \frac{a_1 + (n-1)d - a_1}{d(\sqrt{a_n} + \sqrt{a_1})} \\
 &= \frac{1}{d} \frac{(n-1)d}{d(\sqrt{a_n} + \sqrt{a_1})} = \frac{n-1}{\sqrt{a_n} + \sqrt{a_1}} \\
 &= \text{R.H.S.}
 \end{aligned}$$

Do yourself - 2 :

- (i) Find the sum of first 24 terms of the A.P. a_1, a_2, a_3, \dots , if it is known that $a_1 + a_5 + a_{10} + a_{15} + a_{20} + a_{24} = 225$.
- (ii) Find the number of terms common to the two A.P.'s $3, 7, 11, \dots, 407$ and $2, 9, 16, \dots, 709$

4. GEOMETRIC PROGRESSION (G.P.) :

G.P. is a sequence of non zero numbers each of the succeeding term is equal to the preceding term multiplied by a constant. Thus in a GP the ratio of successive terms is constant. This constant factor is called the COMMON RATIO of the sequence & is obtained by dividing any term by the immediately previous term. Therefore $a, ar, ar^2, ar^3, ar^4, \dots$ is a GP with ' a ' as the first term & ' r ' as common ratio.

n^{th} term ; $T_n = a r^{n-1}$

(a) Sum of the first n terms; $S_n = \frac{a(r^n - 1)}{r - 1}$,

if $r \neq 1$

(c) Sum of infinite G.P. , $S_\infty = \frac{a}{1-r}$; $0 < |r| < 1$

5. PROPERTIES OF GP :

(a) If each term of a G.P. be multiplied or divided by the some non-zero quantity, then the resulting sequence is also a G.P.

(b) Three consecutive terms of a GP :

$a/r, a, ar$;

Four consecutive terms of a GP : $a/r^3, a/r, ar, ar^3$ & so on.

(c) If a, b, c are in G.P. then $b^2 = ac$.

(d) If in a G.P, the product of two terms which are equidistant from the first and the last term, is constant and is equal to the product of first and last term.

$\Rightarrow T_k \cdot T_{n-k+1} = \text{constant} = a \cdot l$

(e) If each term of a G.P. be raised to the same power, then resulting sequence is also a G.P.

(f) In a G.P., $T_r^2 = T_{r-k} \cdot T_{r+k}$, $k < r, r \neq 1$

(g) If the terms of a given G.P. are chosen at regular intervals, then the new sequence is also a G.P.

- (h) If $a_1, a_2, a_3, \dots, a_n$ is a G.P. of positive terms, then $\log a_1, \log a_2, \dots, \log a_n$ is an A.P. and vice-versa.
- (i) If a_1, a_2, a_3, \dots and b_1, b_2, b_3, \dots are two G.P.'s then $a_1b_1, a_2b_2, a_3b_3, \dots$ & $\frac{a_1}{b_1}, \frac{a_2}{b_2}, \frac{a_3}{b_3}, \dots$ is also in G.P.

Illustration 6 :

If a, b, c, d and p are distinct real numbers such that

$$(a^2 + b^2 + c^2)p^2 - 2p(ab + bc + cd) + (b^2 + c^2 + d^2) \leq 0 \text{ then } a, b, c, d \text{ are in}$$

- (A) A.P. (B) G.P.
 (C) H.P. (D) none of these

Solution :

Here, the given condition $(a^2 + b^2 + c^2)p^2 - 2p(ab + bc + ca) + b^2 + c^2 + d^2 \leq 0$

$$\Rightarrow (ap - b)^2 + (bp - c)^2 + (cp - d)^2 \leq 0$$

\therefore a square can not be negative

$$\therefore ap - b = 0, bp - c = 0, cp - d = 0$$

$$\Rightarrow p = \frac{b}{a} = \frac{c}{b} = \frac{d}{c} \Rightarrow a, b, c, d \text{ are in G.P.}$$

Ans. (B)

Illustration 7 :

If a, b, c are in G.P., then the equations $ax^2 + 2bx + c = 0$ and $dx^2 + 2ex + f = 0$

have a common root if $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in -

- (A) A.P. (B) G.P.
 (C) H.P. (D) none of these

Solution :

a, b, c are in G.P. $\Rightarrow b^2 = ac$

Now the equation $ax^2 + 2bx + c = 0$ can be rewritten as $ax^2 + 2\sqrt{ac}x + c = 0$

$$\Rightarrow (\sqrt{ax} + \sqrt{c})^2 = 0 \Rightarrow -\sqrt{\frac{c}{a}}, -\sqrt{\frac{c}{a}}$$

If the two given equations have a common root, then this root must be -

$$\sqrt{\frac{c}{a}}.$$

$$\text{Thus } d \frac{c}{a} - 2e \sqrt{\frac{c}{a}} + f = 0$$

$$\Rightarrow \frac{d}{a} + \frac{f}{c} = \frac{2e}{c} \sqrt{\frac{c}{a}} = \frac{2e}{\sqrt{ac}} = \frac{2e}{b}$$

$$\Rightarrow \frac{d}{a}, \frac{e}{b}, \frac{f}{c} \text{ are in A.P. } \text{Ans. (A)}$$

Illustration 8 :

A number consists of three digits which are in G.P. the sum of the right hand and left hand digits exceeds twice the middle digit by 1 and the sum of the left hand and middle digits is two third of the sum of the middle and right hand digits. Find the numbers.

Solution :

Let the three digits be a, ar and ar^2 then number is

$$100a + 10ar + ar^2 \quad \dots \text{(i)}$$

$$\text{Given, } a + ar^2 = 2ar + 1$$

$$\text{or } a(r^2 - 2r + 1) = 1 \quad \dots \text{(ii)}$$

$$\text{or } a(r - 1)^2 = 1 \quad \dots \text{(ii)}$$

$$\text{Also given } a + ar = \frac{2}{3}(ar + ar^2)$$

$$\Rightarrow 3 + 3r = 2r + 2r^2$$

$$\Rightarrow 2r^2 - r - 3 = 0 \Rightarrow (r + 1)(2r - 3) = 0$$

$$\therefore r = -1, 3/2$$

$$\text{for } r = -1, \quad a = \frac{1}{(r-1)^2} = \frac{1}{4} \notin I$$

$$\therefore r \neq -1$$

$$\text{for } r = 3/2,$$

$$a = \frac{1}{\left(\frac{3}{2}-1\right)^2} = 4 \quad \{ \text{from (ii)} \}$$

From (i), number is

$$400 + 10.4 \cdot \frac{3}{2} + 4 \cdot \frac{9}{4} = 469 \quad \text{Ans.}$$

Illustration 9 :

Find the value of $0.\overline{3258}$

Solution :

$$\text{Let } R = 0.32 \overline{58}$$

$$\Rightarrow R = 0.32585858\dots \quad \dots \text{(i)}$$

Here number of figures which are not recurring is 2 and number of figures which are recurring is also 2.

then $100R = 32.585858\dots\dots\dots$ (ii)
and $10000R = 3258.5858\dots\dots\dots$ (iii)

Subtracting (ii) from (iii), we get

$$9900R = 3226 \Rightarrow R = \frac{1613}{4950}$$

Aliter Method : $R = .32 + .0058 + .0058 + .000058 + \dots\dots\dots$

$$= .32 + \frac{58}{10^4} \left(1 + \frac{1}{10} + \frac{1}{10^4} + \dots\dots\dots \infty \right)$$

$$= .32 + \frac{58}{10^4} \left(\frac{1}{1 - \frac{1}{100}} \right)$$

$$= \frac{32}{100} + \frac{58}{9900} = \frac{3168+58}{9900} = \frac{3226}{9900}$$

$$= \frac{1613}{4950}$$

Do yourself - 3 :

- (i) Find a three digit number whose consecutive digits form a G.P. If we subtract 792 from this number, we get a number consisting of the same digits written in the reverse order. Now, if we increase the second digit of the required number by 2, then the resulting digits will form an A.P.
- (ii) If the third term of G.P. is 4, then find the product of first five terms.
- (iii) If a, b, c are respectively the p^{th} , q^{th} and r^{th} terms of the given G.P., then show that $(q-r)\log a + (r-p)\log b + (p-q)\log c = 0$, where $a, b, c > 0$.
- (iv) Find three numbers in G.P., whose sum is 52 and the sum of whose products in pairs is 624.
- (v) The rational number which equals the number $2.\overline{357}$ with recurring decimal is
 (A) $\frac{2357}{999}$ (B) $\frac{2379}{997}$
 (C) $\frac{785}{333}$ (D) $\frac{2355}{1001}$

6. HARMONIC PROGRESSION (H.P.):

A sequence is said to be in H.P. if the reciprocal of its terms are in AP.

If the sequence $a_1, a_2, a_3, \dots, a_n$ is an HP then $\frac{1}{a_1}, \frac{1}{a_2}, \dots, \frac{1}{a_n}$ is an AP. Here we do not have the formula for the sum of the n terms of an HP. The general form of a harmonic progression

$$\text{is } \frac{1}{a}, \frac{1}{a+d}, \frac{1}{a+2d}, \dots, \frac{1}{a+(n-1)d}$$

Note : No term of any H.P. can be zero.

- (i) If a, b, c are in HP, then

$$b = \frac{2ac}{a+c} \text{ or } \frac{a}{c} = \frac{a-b}{b-c}$$

Illustration 10 :

The sum of three numbers are in H.P. is 37 and the sum of their reciprocals is $1/4$. Find the numbers.

Solution :

Three numbers are in H.P. can be taken as

$$\frac{1}{a-d}, \frac{1}{a}, \frac{1}{a+d} \text{ then}$$

$$\frac{1}{a-d} + \frac{1}{a} + \frac{1}{a+d} = 37 \quad \dots\dots\dots (i)$$

$$\text{and } a-d + a + a+d = \frac{1}{4}$$

$$\Rightarrow a = \frac{1}{12}$$

from (i),

$$\frac{12}{1-12d} + 12 + \frac{12}{1+12d} = 37$$

$$\Rightarrow \frac{12}{1-12d} + \frac{12}{1+12d} = 25$$

$$\Rightarrow \frac{24}{1-144d^2} = 25$$

$$\Rightarrow 1 - 144d^2 = \frac{24}{25}$$

$$\Rightarrow d^2 = \frac{1}{25 \times 144}$$

$$\therefore d = \pm \frac{1}{60}$$

$$\therefore a - d, a, a + d \text{ are } \frac{1}{15}, \frac{1}{12}, \frac{1}{10} \text{ or } \frac{1}{10},$$

$$\frac{1}{12}, \frac{1}{15}$$

Hence, three numbers in H.P. are 15, 12, 10 or 10, 12, 15
Ans.

Illustration 11 :

Suppose a is a fixed real number such that $\frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{rz}$

If p, q, r are in A.P., then prove that x, y, z are in H.P.

Solution :

\therefore p, q, r are in A.P.

$$\therefore q-p = r-q \quad \dots\dots (i)$$

$$\Rightarrow p-q = q-r = k \text{ (let)}$$

$$\text{given } \frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{rz}$$

$$\Rightarrow \frac{\frac{a}{x}-1}{p} = \frac{\frac{a}{y}-1}{q} = \frac{\frac{a}{z}-1}{r}$$

$$\Rightarrow \frac{\left(\frac{a}{x}-1\right) - \left(\frac{a}{y}-1\right)}{p-q} = \frac{\left(\frac{a}{y}-1\right) - \left(\frac{a}{z}-1\right)}{q-r}$$

(by law of proportion)

$$\frac{\frac{a}{x}-a}{k} = \frac{\frac{a}{y}-a}{k}$$

$$\Rightarrow \frac{x-y}{k} = \frac{y-z}{k} \quad \{ \text{from (i)} \}$$

$$\Rightarrow a \left(\frac{1}{x} - \frac{1}{y} \right) = a \left(\frac{1}{y} - \frac{1}{z} \right)$$

$$\Rightarrow \frac{1}{x} - \frac{1}{y} = \frac{1}{y} - \frac{1}{z}$$

$$\therefore \frac{2}{y} = \frac{1}{x} = \frac{1}{z}$$

$$\therefore \frac{1}{x}, \frac{1}{y}, \frac{1}{z} \text{ are in A.P.}$$

Hence x, y, z are in H.P.

Do yourself - 4 :

(i) If the 7th term of a H.P. is 8 and the 8th term is 7. Then find the 28th term.

(ii) In a H.P., if 5th term is 6 and 3rd term is 10. Find the 2nd term.

(iii) If the pth, qth and rth terms of a H.P. are a, b, c respectively, then prove that

$$\frac{q-r}{a} + \frac{r-p}{b} + \frac{p-q}{c} = 0$$

7. MEANS

(a) ARITHMETIC MEAN :

If three terms are in A.P. then the middle term is called the A.M. between the other two, so if a, b, c are in A.P., b is A.M. of a & c. So A.M. of a and c = $\frac{a+c}{2} = b$.

n-ARITHMETIC MEANS BETWEEN TWO NUMBERS :

If a, b be any two given numbers & a, A₁, A₂, ..., A_n, b are in AP, then A₁, A₂, ..., A_n are the 'n' A.M.'s between a & b then.

A₁ = a + d, A₂ = a + 2d, ..., A_n = a + nd or b - d, where

$$d = \frac{b-a}{n+1}$$

$$\Rightarrow A_1 = a + \frac{b-a}{n+1}, A_2 = a + \left(\frac{2b-a}{n+1} \right), \dots\dots$$

Note : Sum of n A.M.'s inserted between a & b is equal to n times the single A.M. between a & b

i.e. $\sum_{r=1}^n A_r = nA$ where A is the single A.M. between a & b.

(b) GEOMETRIC MEAN :

If a, b, c are in G.P., then b is the G.M. between a & c , $b^2 = ac$. So G.M. of a and $c = \sqrt{ac} = b$

n-GEOMETRIC MEANS BETWEEN TWO NUMBERS :

If a, b are two given positive numbers & $a, G_1, G_2, \dots, G_n, b$ are in G.P. Then $G_1, G_2, G_3, \dots, G_n$ are ' n ' G.Ms between a & b , where $b = ar^{n+1}$

$$\Rightarrow r = (b/a)^{1/n+1}$$

$$G_1 = a(b/a)^{1/n+1},$$

$$G_2 = a(b/a)^{2/n+1}, \dots,$$

$$G_n = a(b/a)^{1/n+1},$$

$$= ar,$$

$$= ar^2, \dots,$$

$$= ar^n = b/r$$

Note : The product of n G.Ms between a & b is equal to n^{th} power of the single

$$\text{G.M. between } a \text{ & } b \text{ i.e. } \prod_{r=1}^n G_r = (G)^n$$

where G is the single G.M. between a & b

(c) HARMONIC MEAN :

If a, b, c are in H.P., then b is H.M. between a & c . So H.M. of a and

$$c = \frac{2ac}{a+c} = b.$$

Insertion of ' n ' HM's between a and b :

a, H₁, H₂, H₃, ..., H_n, b → H.P

$$\frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2}, \frac{1}{H_3}, \dots, \frac{1}{H_n}, \frac{1}{b} \rightarrow$$

A.P.

$$\frac{1}{b} = \frac{1}{a} + (n+1)d$$

$$\Rightarrow D = \frac{b-a}{n+1}$$

$$\frac{1}{H_n} = \frac{1}{a} + n \left(\frac{\frac{1}{b} - \frac{1}{a}}{n+1} \right)$$

Do yourself - 5 :

- (i) If $\frac{a^n + b^n}{a^{n-1} + b^{n-1}}$ is the G.M. between a & b then find the value of ' n '.
 - (ii) If b is the harmonic mean between a and c , then prove that
- $$\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{a} + \frac{1}{c}.$$

8. ARITHMETICO - GEOMETRIC SERIES :

A series, each term of which is formed by multiplying the corresponding term of an A.P. & G.P. is called the Arithmetico-Geometric Series , e.g. $1 + 3x + 5x^2 + 7x^3 + \dots$

Here $1, 3, 5, \dots$ are in A.P. & $1, x, x^2, x^3, \dots$ are in G.P.

(a) SUM OF N TERMS OF AN ARITHMETICO-GEOMETRIC SERIES :

Let $S_n = a + (a+d)r + (a+2d)r^2 + \dots + [a+(n-1)d]r^{n-1}$

$$\text{then } S_n = \frac{a}{1-r} + \frac{dr(1-r^{n-1})}{(1-r)^2} - \frac{[a+(n-1)d]r^n}{1-r}, r \neq 1$$

(b) SUM TO INFINITY :

If $0 < |r| < 1$ & $n \rightarrow \infty$, then $\lim_{n \rightarrow \infty} r^n = 0$,

$$S_\infty = \frac{a}{1-r} + \frac{dr}{(1-r)^2}$$

Illustration 12 :

Find the sum of series $4 - 9x + 16x^2 - 25x^3 + 36x^4 - 49x^5 + \dots \infty$.

Solution :

Let $S = 4 - 9x + 16x^2 - 25x^3 + 36x^4 - 49x^5 + \dots \infty$
 $- Sx = -4x + 9x^2 - 16x^3 + 25x^4 - 36x^5 + \dots \infty$

On subtraction, we get

$$S(1+x) = 4 - 5x + 7x^2 - 9x^3 + 11x^4 - 13x^5 + \dots \infty$$

$$-S(1+x)x = -4x + 5x^2 - 7x^3 + 9x^4 - 11x^5 + \dots \infty$$

On subtraction, we get

$$S(1+x)^2 = 4 - x + 2x^2 - 2x^3 + 2x^4 - 2x^5 + \dots \infty$$

$$= 4 - x + 2x^2 (1 - x + x^2 - \dots \infty) = 4 - x + \frac{2x^2}{1+x} = \frac{4+3x+x^2}{1+x}$$

$$S = \frac{4+3x+x^2}{(1+x)^3} \quad \text{Ans.}$$

Illustration 13 :

Find the sum of series upto n terms

$$\left(\frac{2n+1}{2n-1}\right) + 3\left(\frac{2n+1}{2n-1}\right)^2 + 5\left(\frac{2n+1}{2n-1}\right)^3 + \dots$$

Solution :

For $x \neq 1$, let

$$S = x + 3x^2 + 5x^3 + \dots + (2n-3)x^{n-1} + (2n-1)x^n \quad \dots \text{(i)}$$

$$\Rightarrow xS = x^2 + 3x^3 + \dots + (2n-5)x^{n-1} + (2n-3)x^n + (2n-1)x^{n+1} \quad \dots \text{(ii)}$$

Subtracting (ii) from (i), we get

$$(1-x)S = x + 2x^2 + 2x^3 + \dots + 2x^{n-1} + 2x^n - (2n-1)x^{n+1} = x + \frac{2x^2(1-x^{n-1})}{1-x} - (2n-1)x^{n+1}$$

$$= \frac{x}{1-x} [1 - x + 2x - 2x^n - (2n-1)x^n + (2n-1)x^{n+1}]$$

$$\Rightarrow S = \frac{x}{(1-x)^2} [(2n-1)x^{n+1} - (2n+1)x^n + 1 + x]$$

$$\text{Thus } \left(\frac{2n+1}{2n-1}\right) + 3\left(\frac{2n+1}{2n-1}\right)^2 + \dots + (2n-1)\left(\frac{2n+1}{2n-1}\right)^n$$

$$= \left(\frac{2n+1}{2n-1}\right)\left(\frac{2n-1}{2}\right)^2$$

$$\left[(2n-1)\left(\frac{2n+1}{2n-1}\right)^{n+1} - (2n+1)\left(\frac{2n+1}{2n-1}\right)^n + 1 + \frac{2n+1}{2n-1} \right] \\ = \frac{4n^2-1}{4} \cdot \frac{4n}{2n-1} = n(2n+1) \quad \text{Ans.}$$

Do yourself - 6 :

(i) Find sum to n terms of the series

$$3 + 5 \times \frac{1}{4} + 7 \times \frac{1}{4^2} + \dots$$

(ii) If the sum to the infinity of the series

$$3 + 5r + 7r^2 + \dots \text{ is } \frac{44}{9}, \text{ then find the value of } r.$$

(iii) If the sum to infinity of the series

$$3 + (3+d) \cdot \frac{1}{4} + (3+2d) \cdot \frac{1}{4^2} + \dots \text{ is } \frac{44}{9} \text{ then find } d.$$

9. SIGMA NOTATIONS (Σ)

THEOREMS :

$$(a) \sum_{r=1}^n (a_r \pm b_r) = \sum_{r=1}^n a_r \pm \sum_{r=1}^n b_r$$

$$(b) \sum_{r=1}^n ka_r = k \sum_{r=1}^n a_r$$

$$(c) \sum_{r=1}^n k = nk ; \text{ where } k \text{ is a constant.}$$

10. RESULTS

$$(a) \sum_{r=1}^n r = \frac{n(n+1)}{2} \quad (\text{sum of the first } n \text{ natural numbers})$$

$$(b) \sum_{r=1}^n r^2 = \frac{n(n+1)(2n+1)}{6} \quad (\text{sum of the squares of the first } n \text{ natural numbers})$$

$$(c) \sum_{r=1}^n r^3 = \frac{n^2(n+1)^2}{4} = \left[\sum_{r=1}^n r \right]^2 \quad (\text{sum of the cubes of the first } n \text{ natural numbers})$$

$$(d) \sum_{r=1}^n r^4 = \frac{n}{30} (n+1)(2n+1)(3n^2+3n-1)$$

(e) $\sum_{r=1}^n (2r-1) = n^2$ (sum of first n odd natural numbers)

(f) $\sum_{r=1}^n 2r = n(n+1)$ (sum of first n even natural numbers)

Note :

If n^{th} term of a sequence is given by $T_n = an^3 + bn^2 + cn + d$ where a, b, c, d are constants, then sum of n terms $S_n = \sum T_n = a\sum n^3 + b\sum n^2 + c\sum n + \sum d$

This can be evaluated using the above results.

Illustration 14 :

Sum up to 16 terms of the series

$$\frac{1^3}{1} + \frac{1^3 + 2^3}{1+3} + \frac{1^3 + 2^3 + 3^3}{1+3+5} + \dots \text{is}$$

(A) 450

(B) 456

(C) 446

(D) none of these

Solution :

$$t_n = \frac{1^3 + 2^3 + 3^3 + \dots + n^3}{1+3+5+\dots(2n-1)}$$

$$= \frac{\left\{ \frac{n(n+1)}{2} \right\}^2}{\frac{n}{2}\{2+2(n-1)\}} = \frac{\frac{n^2(n+1)^2}{4}}{n^2}$$

$$= \frac{\frac{(n+1)^2}{4}}{\frac{n^2}{4}} = \frac{n^2}{4} + \frac{n}{2} + \frac{1}{4}$$

$$\therefore S_n = \sum t_n = \frac{1}{4} \sum v^2 + \frac{1}{2} \sum v + \frac{1}{4} \sum 1 = \frac{1}{4} .$$

$$\frac{n(n-1)(2n-1)}{6} + \frac{1}{2} \cdot \frac{n(n+1)}{2} + \frac{1}{4} \cdot n$$

$$\therefore S_{16} = \frac{16 \cdot 17 \cdot 33}{24} + \frac{16 \cdot 17}{4} + \frac{16}{4} = 446$$

Ans.(C)

11. METHOD OF DIFFERENCE :

Some times the n^{th} term of a sequence or a series can not be determined by the method, we have

discussed earlier. So we compute the difference between the successive terms of given sequence for obtained the n^{th} terms.

If $T_1, T_2, T_3, \dots, T_n$ are the terms of a sequence then some times the terms $T_2 - T_1, T_3 - T_2, \dots$

constitute an AP/GP. n^{th} term of the series is determined & the sum to n terms of the sequence can easily be obtained.

Case 1 :

(a) If difference series are in A.P., then

Let $T_n = an^2 + bn + c$, where a, b, c are constant

(b) If difference of difference series are in A.P.

Let $T_n = an^3 + bn^2 + cn + d$, where a, b, c, d are constant

Case 2 :

(a) If difference are in G.P., then

Let $T_n = ar^n + b$, where r is common ratio & a, b are constant

(b) If difference of difference are in G.P., then

Let $T_n = ar^n + bn + c$, where r is common ratio & a, b, c are constant

Determine constant by putting $n = 1, 2, 3, \dots, n$ and putting the value of T_1, T_2, T_3, \dots and sum of series ($S_n = \sum T_n$)

Do yourself - 7 :

(i) Find the sum of the series upto n terms

$$1 + \frac{1+2}{2} + \frac{1+2+3}{3} + \frac{1+2+3+4}{4} + \dots$$

(ii) Find the sum of 'n' terms of the series whose n^{th} term is $t_n = 3n^2 + 2n$.

Miscellaneous Illustration :

Illustration 15 :

If $\sum_{r=1}^n T_r = \frac{n}{8}(n+1)(n+2)(n+3)$, then

$$\text{find } \sum_{r=1}^n \frac{1}{T_r}.$$

Solution :

$$\begin{aligned}
 & \because T_n = S_n - S_{n-1} \\
 &= \sum_{r=1}^n T_r - \sum_{r=1}^{n-1} T_r = \frac{n(n+1)(n+2)(n+3)}{8} - \\
 &\quad \frac{(n-1)(n+1)(n+2)}{8} = \frac{n(n+1)(n+2)}{8} \\
 &\quad [(n+3) - (n-1)] \\
 T_n &= \frac{n(n+1)(n+2)}{8}(4) \\
 &= \frac{n(n+1)(n+2)}{2} \\
 \Rightarrow \frac{1}{T_n} &= \frac{2}{n(n+1)(n+2)} \\
 &= \frac{(n+2)-n}{n(n+1)(n+2)} \\
 &= \frac{1}{n(n+1)} - \frac{1}{(n+1)(n+2)} \quad \dots\dots\dots(i) \\
 \text{Let } V_n &= \frac{1}{n(n+1)} \\
 \therefore \frac{1}{T_n} &= V_n - V_{n+1} \\
 \text{Putting } n = 1, 2, 3, \dots, n \\
 \Rightarrow \frac{1}{T_1} + \frac{1}{T_2} + \frac{1}{T_3} + \dots + \frac{1}{T_n} &= (V_1 - V_{n+1}) \\
 \Rightarrow \sum_{r=1}^n \frac{1}{T_r} &= \frac{n^2 + 3n}{2(n+1)(n+2)}
 \end{aligned}$$

Illustration 16 :

Find the sum of n terms of the series 1 . 3 . 5 + 3 . 5 . 7 + 5 . 7 . 9 +

Solution :

The n^{th} term is $(2n-1)(2n+1)(2n+3)$

$$\begin{aligned}
 T_n &= (2n-1)(2n+1)(2n+3) \\
 T_n &= \frac{1}{8} (2n-1)(2n+1)(2n+3) \{(2n+ \\
 &5) - (2n-3)\} \\
 &= \frac{1}{8} (V_n - V_{n-1}) [\text{Let } V_n = (2n-1)(2n+ \\
 &1)(2n+3)(2n+5)]
 \end{aligned}$$

$$\begin{aligned}
 S_n &= \sum T_n = \frac{1}{8} [V_n - V_0] \\
 &\therefore S_n = \frac{(2n-1)(2n+1)(2n+3)(2n+5)}{8} + \frac{15}{8} \\
 &= n(2n^3 + 8n^2 + 7n - 2) \quad \text{Ans.}
 \end{aligned}$$

Illustration 17 :

Find the sum of n terms of the series 3 + 7 + 14 + 24 + 37 +

Solution :

Clearly here the differences between the successive terms are

7 - 3, 14 - 7, 24 - 14, i.e. 4, 7, 10, 13,....., which are in A.P.

Let $S = 3 + 7 + 14 + 24 + \dots + T_n$

$S = 3 + 7 + 14 + \dots + T_{n-1} + T_n$

Subtracting, we get

$0 = 3 + [4 + 7 + 10 + 13 + \dots \text{ (n-1) terms}] - T_n$

$\therefore T_n = 3 + S_{n-1}$ of an A.P. whose $a = 4$ and $d = 3$.

$$\begin{aligned}
 \therefore T_n &= 3 + \left(\frac{n-1}{2} \right) (2.4(n-2)3) = \\
 &\quad \frac{6+(n-1)(3n+2)}{4} \text{ or,}
 \end{aligned}$$

$$T_n = \frac{1}{2} (3n^2 - n + 4)$$

Now putting $n = 1, 2, 3, \dots, n$ and adding

$$\begin{aligned}
 \therefore S_n &= \frac{1}{2} [3\sum n^2 - \sum n + 4n] \\
 &= \frac{1}{2} \left[3 \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} + 4n \right] \\
 &= \frac{n}{2} (n^2 + n + 4) \quad \text{Ans.}
 \end{aligned}$$

Aliter Method :

$$\text{Let } T_n = an^2 + bn + c$$

$$\text{Now, } T_1 = 3 = a + b + c \quad \dots\dots\dots(i)$$

$$T_2 = 7 = 4a + 2b + c \quad \dots\dots\dots(ii)$$

$$T_3 = 14 = 8a + 3b + c \quad \dots\dots\dots(iii)$$

Solving (i), (ii) & (iii) we get

$$a = \frac{3}{2}, b = -\frac{1}{2} \text{ & } c = 2$$

$$\therefore T_n = \frac{1}{2} (3n^2 - n + 4)$$

$$\Rightarrow S_n = \sum T_n = \frac{1}{2} [3\sum n^2 - \sum n + 4n]$$

$$= \frac{1}{2} \left[3 \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} + 4n \right]$$

$$= \frac{n}{2} (n^2 + n + 4) \quad \text{Ans.}$$

Illustration 18 :

Find the sum of n-terms of the series
 $1 + 4 + 10 + 22 + \dots$

Solution :

$$\text{Let } S = 1 + 4 + 10 + 22 + \dots + T_n \quad \dots \text{(i)}$$

$$S = 1 + 4 + 10 + \dots + T_{n-1} + T_n \quad \dots \text{(ii)}$$

$$(i) - (ii)$$

$$\Rightarrow T_n = 1 + (3 + 6 + 12 + \dots + T_n - T_{n-1})$$

$$T_n = 1 + 3 \left(\frac{2^{n-1} - 1}{2-1} \right)$$

$$T_n = 3 \cdot 2^{n-1} - 2$$

So

$$S_n = \sum T_n = 3\sum 2^{n-1} - \sum 2$$

$$= 3 \left(\frac{2^n - 1}{2-1} \right) - 2n = 3 \cdot 2^n - 2n - 3 \quad \text{Ans.}$$

Aliter Method :

$$\text{Let } T_n = ar^n + b, \text{ where } r = 2$$

$$\text{Now } T_1 = 1 = ar + b \quad \dots \text{(i)}$$

$$T_2 = 4 = ar^2 + b \quad \dots \text{(ii)}$$

Solving (i) & (ii), we get

$$a = \frac{3}{2}, b = -2 \therefore T_n = 3 \cdot 2^{n-1} - 2$$

$$\Rightarrow S_n = \sum T_n = 3\sum 2^{n-1} - \sum 2$$

$$= 3 \left(\frac{2^n - 1}{2-1} \right) - 2n = 3 \cdot 2^n - 2n - 3$$

Ans.

Illustration 19 :

The series of natural numbers is divided into groups (1), (2, 3, 4), (5, 6, 7, 8, 9) and so on. Show that the sum of the numbers in n^{th} group is $n^3 + (n-1)^3$

Solution :

The groups are (1), (2, 3, 4), (5, 6, 7, 8, 9)

The number of terms in the groups are 1, 3, 5.....

∴ The number of terms in the n^{th} group = $(2n-1)$ the last term of the n^{th} group is n^2

If we count from last term common difference should be -1

So the sum of numbers in the n^{th} group

$$= \left(\frac{2n-1}{2} \right) \{ 2n^2 + (2n-2)(-1) \}$$

$$= (2n-1)(n^2 - n + 1)$$

$$= 2n^3 - 3n^2 + 3n - 1 = n^3 + (n-1)^3$$

Illustration 20 :

Find the natural number 'a' for which

$$\sum_{k=1}^n f(a+k) = 16(2^n - 1), \text{ where the}$$

function f satisfied $f(x+y) = f(x)f(y)$.
 $f(y)$ for all natural number x, y and further $f(1) = 2$.

Solution :

It is given that

$$f(x+y) = f(x)f(y) \text{ and } f(1) = 2$$

$$f(1+1) = f(1)f(1)$$

$$\Rightarrow f(2) = 2^2, f(1+2) = f(1)f(2)$$

$$\Rightarrow f(3) = 2^3, f(2+2) = f(2)f(2)$$

$$\Rightarrow f(4) = 2^4$$

Similarly $f(k) = 2^k$ and $f(a) = 2^a$

$$\text{Hence, } \sum_{k=1}^n f(a+k) = \sum_{k=1}^n f(a)f(k) = f(a)$$

$$\sum_{k=1}^n f(k) = 20^a \{ 2^1 + 2^2 + \dots + 2^n \}$$

$$= 2^a \left\{ \frac{2(2^n - 1)}{2-1} \right\} = 2^{a+1}(2^n - 1)$$

$$\text{But } \sum_{k=1}^n f(a+k) = 16(2^n - 1)$$

$$2^{a+1}(2^n - 1) = 16(2^n - 1)$$

$$\therefore 2^{a+1} = 2^4$$

$$\therefore a+1 = 4$$

$$\Rightarrow a = 3 \quad \text{Ans.}$$

19. IMPORTANT RESULTS :

(ix) $\sin\alpha + \sin(\alpha + \beta) + \sin(\alpha + 2\beta) + \dots + \sin(\alpha + (n-1)\beta)$

$$= \frac{\sin\left\{\alpha + \left(\frac{n-1}{2}\right)\beta\right\} \sin\left(\frac{n\beta}{2}\right)}{\sin\left(\frac{\beta}{2}\right)}$$

(x) $\cos\alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos(\alpha + (n-1)\beta)$

$$= \frac{\cos\left\{\alpha + \left(\frac{n-1}{2}\right)\beta\right\} \sin\left(\frac{n\beta}{2}\right)}{\sin\left(\frac{\beta}{2}\right)}$$

Do yourself - 8 :

Evaluate $\sin \frac{\pi}{n} + \sin \frac{3\pi}{n} + \sin \frac{5\pi}{n} + \dots + \sin \frac{(2n-1)\pi}{n}$ to n terms

Miscellaneous Illustration :
Illustration 21:

Prove that

$$\tan\alpha + 2 \tan 2\alpha + 2^2 \tan^2 \alpha + \dots + 2^{n-1} \tan 2^{n-1}\alpha + 2^n \cot 2^n \alpha = \cot \alpha$$

Solution :

We know $\tan \theta = \cot \theta - 2 \cot 2\theta$

.....(i)

Putting $\theta = \alpha, 2\alpha, 2^2\alpha, \dots$ in (i), we get

$$\tan\alpha = (\cot \alpha - 2 \cot 2\alpha)$$

$$2(\tan 2\alpha) = 2(\cot 2\alpha - 2 \cot 2^2\alpha)$$

$$2^2(\tan 2^2\alpha) = 2^2(\cot 2^2\alpha - 2 \cot 2^3\alpha)$$

$$\dots \dots \dots \\ 2^{n-1}(\tan 2^{n-1}\alpha) = 2^{n-1}(\cot 2^{n-1}\alpha - 2 \cot 2^n\alpha)$$

Adding,

$$\tan\alpha + 2 \tan 2\alpha + 2^2 \tan^2 \alpha + \dots + 2^{n-1}$$

$$\tan 2^{n-1}\alpha = \cot \alpha - 2^n \cot 2^n \alpha$$

$$\therefore \tan\alpha + 2 \tan 2\alpha + 2^2 \tan^2 \alpha + \dots + 2^{n-1} \\ \tan 2^{n-1}\alpha + 2^n \cot 2^n \alpha = \cot \alpha$$

ANSWERS FOR DO YOURSELF

1 : (i) (a) $\frac{2}{1}, \frac{4}{2}, \frac{8}{3}, \frac{16}{4}, \dots$,
(b) $\frac{2}{3}, \frac{4}{9}, \frac{2}{27}, \frac{4}{81}, \dots$

(iii) -(p + q)

2 : (i) 900 (ii) 14

3 : (i) 931 (ii) 4^5
(iv) 4, 12, 36 (v) C

4 : (i) 2 (ii) 15

5 : (i) $\frac{1}{2}$

6 : (i) $4 + \frac{8}{9} \left(1 - \frac{1}{4^{n-1}}\right) - \left(\frac{2n+1}{3 \times 4^{n-1}}\right)$
(ii) $\frac{1}{4}$ (iii) 2

7 : (i) $\frac{n(n+3)}{4}$ (ii)
 $\frac{n(n+1)(2n+3)}{2}$

8 : 0

Result JEE Main

Result Highlights 2019



100 Percentile
HIMANSHU GAURAV SINGH
2019 (*SDCCP)



99.99 Percentile
GAURAV KRISHAN GUPTA
2020 (DLP)



99.98 Percentile
SARTHAK ROUT
2020 (CCP)



99.98 Percentile
VIBHAV AGGARWAL
2019 (CCP)



99.97 Percentile
RITVIK GUPTA
2020 (DLP)



99.97 Percentile
BHAVYA JAIN
2020 (CCP)



99.96 Percentile
AYUSH PATTNAIK
2019 (CCP)



99.96 Percentile
SAYANTAN DHAR
2020 (DLP)

Result Highlights 2018



AIR
29

Sarthak Behera



AIR
36

Shashank Agrawal



AIR
46

Shreya Pathak



AIR
48

Raaghav Raaj



AIR
2
SC

Aayush Kadam



AIR
18
OBC

Gunjan

Result Highlights 2017



AIR
12

Abhay Goyal



AIR
14

Piyush Tibarwal



AIR
20

Abhinav Kumar



AIR
38

Tushar Gautam



AIR
49

Saurav Sahoo



AIR
60

Pratyush

Mentored by Team



AIR
4

Kushal Babel
2014



AIR
15

Gourika Bang
2015



AIR
21

Amit Malav
2013



AIR
24

Suraj Sanjay Jog
2012



AIR
38

Chitraang Murdia
2014



AIR
58

Pulkit Tandon
2012



AIR
68

Kapil Shobhnani
2016



AIR
70

Utkarsh Sinha
2015



AIR
80

S John S
2015