

## Lecture 2.1: Intuition Behind Sampling Distributions

2014/01/29

# Behavior of Statistics

Say we draw a random sample of size  $n = 100$  from any distribution that has true population mean  $\mu = 5$ .

# Behavior of Statistics

Say we draw a random sample of size  $n = 100$  from any distribution that has true population mean  $\mu = 5$ .

Two Important Conceptual Questions:

# Behavior of Statistics

Say we draw a random sample of size  $n = 100$  from any distribution that has true population mean  $\mu = 5$ .

Two Important Conceptual Questions:

1. If we compute the sample mean  $\bar{x}$  of these points, are we going to always get exactly 5?

# Behavior of Statistics

Say we draw a random sample of size  $n = 100$  from any distribution that has true population mean  $\mu = 5$ .

## Two Important Conceptual Questions:

1. If we compute the sample mean  $\bar{x}$  of these points, are we going to always get exactly 5?
2. Say we do this once and the sample mean is  $\bar{x} = 5.025$ . If we repeat this procedure (i.e. generate a **new** sample of 100 points and compute  $\bar{x}$ ) are we going to get  $\bar{x} = 5.025$  exactly?

# Behavior of Statistics

Do this for the 1st time

We get, say,  $\bar{x} = 4.831$

# Behavior of Statistics

Do this for the 1st time

We get, say,  $\bar{x} = 4.831$

Do this for the 2nd time

We get, say,  $\bar{x} = 5.104$

# Behavior of Statistics

Do this for the 1st time

Do this for the 2nd time

Do this for the 3rd time

We get, say,  $\bar{x} = 4.831$

We get, say,  $\bar{x} = 5.104$

We get, say,  $\bar{x} = 4.965$



# Behavior of Statistics

Do this for the 1st time

Do this for the 2nd time

Do this for the 3rd time

...

We get, say,  $\bar{x} = 4.831$

We get, say,  $\bar{x} = 5.104$

We get, say,  $\bar{x} = 4.965$

# Behavior of Statistics

Do this for the 1st time	We get, say, $\bar{x} = 4.831$
Do this for the 2nd time	We get, say, $\bar{x} = 5.104$
Do this for the 3rd time	We get, say, $\bar{x} = 4.965$
...	
Do this for the 1000th time	We get, say, $\bar{x} = 4.957$

# Behavior of Statistics

Do this for the 1st time	We get, say, $\bar{x} = 4.831$
Do this for the 2nd time	We get, say, $\bar{x} = 5.104$
Do this for the 3rd time	We get, say, $\bar{x} = 4.965$
...	
Do this for the 1000th time	We get, say, $\bar{x} = 4.957$


The **sampling distribution** describes the random behavior of these estimates.

## Example: Sampling Distribution of the Sample Mean

Consider the histogram of 1000 instances of  $\bar{x}$ , where each  $\bar{x}$  is computed from a sample of  $n = 100$   $\text{Normal}(\mu = 5, \sigma^2 = 4)$  RV's.

## Example: Sampling Distribution of the Sample Mean

Consider the histogram of 1000 instances of  $\bar{x}$ , where each  $\bar{x}$  is computed from a sample of  $n = 100$   $\text{Normal}(\mu = 5, \sigma^2 = 4)$  RV's.



## Standard Error of the Sample Mean $\bar{X}$

The standard deviation associated with a statistic is called the **standard error**.

# Standard Error of the Sample Mean $\bar{X}$

The standard deviation associated with a statistic is called the **standard error**.

If the  $X_i$ 's have standard deviation  $\sigma$ , the **standard error of the sample mean** is

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

# Standard Error of the Sample Mean $\bar{X}$

The standard deviation associated with a statistic is called the **standard error**.

If the  $X_i$ 's have standard deviation  $\sigma$ , the **standard error of the sample mean** is

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$


Notice the  $\sqrt{n}$  in the denominator:  $n$  increases, SE decreases!



## Standard Error of the Sample Mean $\bar{X}$

Recall  $n = 100$  with  $X_i \sim \text{Normal}(\mu = 5, \sigma^2 = 4)$ . The standard error is the SD of the sampling distribution:

$$SE = \frac{\sigma}{\sqrt{n}} = \frac{2}{\sqrt{100}} = \frac{2}{10} = 0.2$$



## Standard Error of the Sample Mean $\bar{x}$

Now compare the sampling distributions based on

## Standard Error of the Sample Mean $\bar{x}$

Now compare the sampling distributions based on

- ▶ 1000 instances of  $\bar{x}$  where  $n = 100$ .  $SE = \frac{2}{\sqrt{100}} = 0.2$

## Standard Error of the Sample Mean $\bar{x}$

Now compare the sampling distributions based on

- ▶ 1000 instances of  $\bar{x}$  where  $n = 100$ .  $SE = \frac{2}{\sqrt{100}} = 0.2$
- ▶ 1000 instances of  $\bar{x}$  where  $n = 1000$ .  $SE = \frac{2}{\sqrt{1000}} = 0.0632$

# Standard Error of the Sample Mean $\bar{x}$

Now compare the sampling distributions based on

- ▶ 1000 instances of  $\bar{x}$  where  $n = 100$ .  $SE = \frac{2}{\sqrt{100}} = 0.2$
- ▶ 1000 instances of  $\bar{x}$  where  $n = 1000$ .  $SE = \frac{2}{\sqrt{1000}} = 0.0632$

