

Article

Far-Infrared Star Formation Rates of Quasar Host Galaxies from Multiwavelength Spectral Energy Distribution Decomposition

Xiaotong Feng ^{1,2*} , Xue-Bing Wu ^{1,2} , Yuming Fu ^{3,4} , Yuxuan Pang ⁵ , Rui Zhu ^{1,2}  and Huimei Wang ^{1,2} ¹ Department of Astronomy, School of Physics, Peking University, Beijing 100871, China² Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China³ Leiden Observatory, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands⁴ Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands⁵ School of Astronomy and Space Science, University of Chinese Academy of Sciences (UCAS), Beijing 100049, China

* Correspondence: xtfeng@pku.edu.cn

Abstract

Reliable star formation rates (SFRs) are essential for studying the connection between black hole growth and quasar host galaxies. We study the far-infrared (FIR) SFRs and the host galaxy properties of 202 SDSS and PG quasars at $0.02 < z \lesssim 0.8$, spanning $\log(\text{SFR}_{\text{FIR}}/M_{\odot} \text{ yr}^{-1}) \simeq -0.45\text{--}2.76$, using multiwavelength spectral energy distribution (SED) decomposition. The photometry covers wavelengths from the optical to the FIR and is supplemented by JCMT/SCUBA-2 observations at 450 and 850 μm . We model the SEDs with CIGALE and AGNfitter and adopt multiple cold dust templates to quantify systematic uncertainties. The median model-dependent scatter among the five FIR SFR estimates is 0.14 dex, and AGNfitter gives FIR SFRs lower than the mean CIGALE estimate by a median of 0.09 dex. For the 58 quasars with SCUBA-2 coverage, including SCUBA-2 data changes the adopted FIR SFR by only ~ 0.01 dex on average but can affect individual sources with limited Herschel coverage or radio-loud emission. Within our FIR-constrained sample, many quasar hosts lie on or above the star-forming main sequence, but the redshift-dependent FIR selection of the SDSS subsample limits conclusions about the full quasar-host population. We find no clear correlation between the main-sequence (MS) offset and the direct Eddington ratio, while the offset is positively related to the infrared-based $L_{\text{tor}}/L_{\text{Edd}}$ proxy. The minimum radiation field intensity in the dust model, U_{min} , increases with bolometric luminosity and dust temperature. WISE W2 (4.6 μm) and W3 (12 μm) combined with Herschel bands can also provide useful empirical indicators of f_{AGN} .

Keywords: quasars; active galactic nuclei; quasar host galaxies; star formation rates; spectral energy distribution decomposition; far-infrared emission; SCUBA-2 observations



Academic Editor: Stefano Vercellone

Received: 25 May 2026

Revised: 3 July 2026

Accepted: 7 July 2026

Published: 16 July 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC BY\) license](https://creativecommons.org/licenses/by/4.0/).

1. Introduction

The growth of supermassive black holes (SMBHs) and the evolution of their host galaxies are closely linked. Empirical correlations between the black hole mass and the properties of the host galaxy bulge (bulge luminosity, stellar mass, stellar velocity dispersion, etc.) suggest that the growth of SMBHs and galaxy formation are linked over a long time [1–3]. The luminous quasars provide an important phase for studying this connection, because rapid black hole accretion and the growth of their host galaxies can occur simultaneously. Therefore, the reliable measurement of the star formation rate (SFR) in a quasar host galaxy is essential to study whether black hole growth is related to star formation.

However, the measurement of SFRs of quasar host galaxies is not simple. Many SFR indicators can be contaminated by the strong emission of the central active galactic nucleus (AGN). The accretion disk dominates optical and ultraviolet continuum, the narrow-line region (NLR) contributes to nebular emission lines, the dusty torus emission is strong at the near (NIR)- to-mid-infrared (MIR) band. Mid-infrared features, such as PAH emission and the [Ne II] line, can also trace star formation [4,5]. However, in luminous quasars, strong AGN mid-infrared emission can make PAH features difficult to measure, while [Ne II] may include emission from the AGN narrow-line region [6,7]. X-ray emission in luminous quasars is also generally dominated by accretion-powered AGN emission and therefore does not directly trace star formation [8,9]. Radio emission and jets can also affect the radio continuum. So, most of the widely used SFR indicators of normal star-forming galaxies are unreliable (UV continuum, optical recombination lines, NIR emission, and radio continuum), unless the AGN contribution is carefully excluded [10,11].

Far-infrared (FIR) emission is one of the useful indicators of SFRs of AGN host galaxies. In star-forming galaxies, ultraviolet and optical photons from young stars are absorbed by dust and reradiated in the infrared (IR) band. So the FIR emission is generally less affected by the AGN torus than NIR and MIR emission and can be a relatively clean tracer of dust-obscured star formation. Many studies have shown that the FIR emission of normal AGNs is dominated by the host galaxy and can be used to probe the connection between star formation and AGN activity [12,13]. However, for luminous quasars, AGN torus may also contribute to FIR emission, and the radio emission from central AGN may affect the submillimeter-to-radio SED. Therefore, directly converting the FIR luminosity into an SFR may overestimate the SFR of quasar host galaxies if the AGN contribution is not properly decomposed.

Therefore, a more robust method to measure the SFR is to do the multiwavelength spectral energy distribution (SED) fitting and decompose the stellar, dust, AGN, and radio components. The SED-fitting decomposition can separate the cold dust emission which is associated with star formation from the AGN hot dust torus. Then, the FIR luminosity of the cold dust component can be used to estimate the FIR SFR. However, the reliability of this method is related to the quality and coverage of the FIR photometric data. Recent simulations have shown that the recovery of star-forming and AGN infrared luminosities depends on the wavelength coverage of the infrared SED. FIR data are especially important for constraining the cold dust component [14]. If there are additional submillimeter constraints, they can be important for reducing the uncertainties of the cold dust component, especially for the sources with no or limited Herschel detections. In addition, different SED-fitting codes can give different SFRs even when they use the same photometric data, because they adopt different model assumptions [15]. The AGN contribution at FIR wavelengths is one source of this uncertainty. Different torus models use different dust geometries and grain properties and can therefore predict different FIR SED shapes. For example, the models of Fritz et al. [16] and Siebenmorgen et al. [17] make different assumptions about the dust distribution and composition. As a result, the separation between AGN-heated dust and star-formation-heated dust can depend on the adopted model, especially when the FIR data are limited. These issues motivate the use of multiple SED models and extended FIR/submillimeter coverage to obtain robust SFR estimates for quasar host galaxies.

The relation between star formation and AGN activity remains debated. Some studies found the correlation between the host galaxy SFR and AGN luminosity, black hole accretion rate, and Eddington ratio, while other studies also found weak or no correlations once redshift and stellar mass are controlled [18–23]. Part of this diversity may be caused by different timescales. The FIR emission traces average star formation over tens to hundreds of Myr, but the variability in AGN accretion has much shorter timescale. It is therefore

important to derive a robust SFR of a quasar host galaxy. Then, we can compare them with the AGN properties while accounting for redshift and sample selection effects.

The main goal of this work is to derive robust FIR-based SFRs for low-redshift type-I quasars and to quantify the main uncertainties in these measurements. We study a sample of 202 quasars at $z \sim 0.02\text{--}0.8$, including 115 SDSS and 87 PG quasars. We compile multiwavelength photometry from the optical to the FIR and supplement the archival data with JCMT/SCUBA-2 observations at 450 and 850 μm . We use two independent SED-fitting approaches, CIGALE and AGNfitter, together with multiple cold dust templates. We first examine how the adopted models and the inclusion of SCUBA-2 data affect the derived SFRs and dust properties. We then use the resulting host galaxy properties to study the locations of quasar hosts relative to the star-forming main sequence and their relations with black hole accretion and dust heating. As an additional application, we examine whether simple infrared colors can provide empirical estimates of the infrared AGN fraction.

This paper is organized as follows. Section 2 describes the sample selection and observations. Section 3 presents the SED-fitting methods and adopted model parameters. Section 4 gives the FIR SFRs and host galaxy properties, discusses the effect of the SCUBA-2 data, and examines the connection between star formation and AGN activity. We summarize our main results in Section 5.

2. Sample and Observations

2.1. Sample Selection

Our sample consists of 202 quasars. We started with the catalog of 105,783 quasars in the Sloan Digital Sky Survey Data Release 7 (SDSS DR7) [24–26] compiled by Shen et al. [27]. This catalog provides a comprehensive set of physical measurements for SDSS DR7 quasars, including continuum measurements around the $H\alpha$, $H\beta$, $Mg\ II$, and $C\ IV$ regions; measurements of broad and narrow emission lines; and black hole masses estimated from multiple broad emission lines. The parent SDSS DR7 catalog includes luminous quasars with $M_i < -22.0$ and, in general, at least one broad emission line with $FWHM > 1000\text{ km s}^{-1}$. A small number of spectroscopically confirmed quasars with complex absorption spectra were also retained. This exception does not refer to the inclusion of narrow-line AGNs. Because this sample will also be used for a companion study of $[O\ II]\ \lambda 3727$ -based SFRs, and the SDSS DR7 spectral wavelength coverage is 3800–9200 $\text{\AA}$, we restrict the redshift range to $z \sim 0.02\text{--}0.8$, yielding 21,075 quasars. To estimate far-infrared SFRs, we need far-infrared data. We therefore cross-match the 21,075 quasars with Herschel [28] catalogs provided by the NASA/IPAC Infrared Science Archive (IRSA)¹. The adopted far-infrared photometry includes the 70, 100, and 160 μm bands from the Photoconductor Array Camera and Spectrometer (PACS) [29] and the 250, 350, and 500 μm bands from the Spectral and Photometric Imaging Receiver (SPIRE) [30]. We require each source to be detected in at least two of these six Herschel bands to provide meaningful constraints on the far-infrared SED. After applying this criterion, we obtain a sample of 120 SDSS quasars.

Figure 1 shows the distribution of the number of sources detected in the six Herschel bands (70, 100, 160, 250, 350, and 500 μm) and the distribution of the number of detected bands for each source. For our SDSS sample, the 250 μm band has the highest detection number (93 of 120), while the number of detections decreases at 350 μm and 500 μm . Most of the sample (100 of 120) are detected in no more than three Herschel bands. As a result, the long-wavelength side of the Rayleigh–Jeans tail is poorly sampled for most sources, which may lead to relatively weak constraints on the far-infrared SED and affect the estimates of far-infrared luminosity, SFR, cold dust temperature, and other derived quantities.




Figure 1. Herschel detection statistics for the SDSS and PG quasars. (a,b) show the SDSS subsample: (a) distribution of the number of sources detected in six Herschel bands and (b) distribution of the number of detected Herschel bands for each source. (c,d) show the same distributions for the PG quasars.

To improve the constraints on the long-wavelength part of the far-infrared SED, we obtained additional submillimeter observations with the Submillimetre Common-User Bolometer Array 2 on the James Clerk Maxwell Telescope (JCMT/SCUBA-2) [31]. The observations are described in Section 2.2.

We also include 87 PG quasars with $z < 0.5$ from the sample analyzed by Shangguan et al. [32] to enlarge the sample, because these objects have relatively extensive Herschel photometric coverage [32]. These objects originate from the Palomar–Green Bright Quasar Survey [33], with optical spectroscopic properties compiled by Boroson and Green [34]. The PG quasars do not follow the same absolute- i -band magnitude selection as the SDSS subsample. While the SDSS parent catalog adopts $M_i < -22.0$, the PG survey was selected using bright optical magnitude and ultraviolet-excess criteria. Thus, the PG quasars should not be regarded as a homogeneous low-redshift extension of the SDSS sample, and possible differences in luminosity and color selection should be considered when interpreting the combined sample. We adopt the Herschel photometry compiled by Shangguan et al. [32], including the available PACS 70, 100, and 160 μm and SPIRE 250, 350, and 500 μm measurements, together with the reported uncertainties and upper limits. These data were not

re-extracted in this work. As shown in Figure 1, the PG sample has generally more complete PACS coverage than the SDSS subsample, whereas the SPIRE detection rate decreases toward longer wavelengths in both samples. Compared with the SDSS quasars, a larger fraction of PG quasars have three or more Herschel detections. A summary of the selection criteria, redshift distributions, and luminosity ranges of the two subsamples is given in Table 1. A total of 5 of the 87 PG quasars are also included in the initially selected SDSS sample of 120 objects. To avoid double counting, these overlapping sources are assigned to the PG subsample, yielding a final sample of 202 unique quasars, comprising 115 SDSS quasars and 87 PG quasars.

Table 1. Comparison of the SDSS and PG subsamples. The listed numbers refer to unique quasars in the final sample. The initial SDSS selection contains 120 objects, five of which overlap with the PG sample. These duplicates are counted only once, resulting in 115 SDSS and 87 PG quasars. For the numerical quantities, values are reported as the median with the 16th and 84th percentiles.

Property	SDSS Subsample	PG Subsample
Number of sources	115	87
Redshift range	$0.02 \leq z \leq 0.82$	$z < 0.5$
Median z	$0.42^{+0.31}_{-0.26}$	$0.14^{+0.18}_{-0.08}$
Median $\log(L_{\text{bol}}/\text{erg s}^{-1})$	$45.53^{+0.46}_{-0.32}$	$45.79^{+0.78}_{-0.57}$
Parent-sample selection	$M_i < -22$; broad-line SDSS DR7 quasars	Bright optical magnitude and UV excess PG/BQS selection
Herschel data source	PACS/SPIRE photometry cross-matched from the IRSA Herschel point-source catalogs (HPPSC v1c and SPIRE PSC v2c) in this work	PACS/SPIRE fluxes, uncertainties, and upper limits adopted from Shangguan et al. [32]

2.2. SCUBA-2 Observations

SCUBA-2 is the submillimeter continuum camera of the 15 m JCMT on Maunakea, Hawaii. It can observe simultaneously at 450 μm and 850 μm . The 450 μm band provides a higher angular resolution and is therefore useful for resolving fine spatial structures, whereas the 850 μm band has higher sensitivity and is more suitable for detecting faint sources.

For our 120 selected SDSS quasars, we carried out several rounds of JCMT/SCUBA-2 observations under the programs M18BP050, M19AP020, M19BP041, and M20AP044. Observations were obtained for 44 sources in total. The targets were not selected randomly from the parent sample. We first used the available Herschel photometry to perform preliminary cold dust SED fits and predict the expected SCUBA-2 flux densities. These predictions were used to estimate the integration time required to reach the planned sensitivity. In assigning observational priorities, we considered both the estimated observing time and the existing Herschel coverage, giving higher priority to sources with fewer Herschel detections in order to improve the constraints on the long-wavelength FIR SED. Owing to the limited telescope time, some sources with very low predicted submillimeter flux densities and correspondingly long estimated integration times were not observed. Thus, the SCUBA-2-observed subset was selected to balance observational feasibility with the need for improved long-wavelength constraints. It should not be regarded as a random or statistically complete subset of the 120 SDSS FIR-selected quasars. The observations were performed in the “CV Daisy” (CV stands for constant velocity) observing mode, which is designed for small and compact sources of order 3 arcmin or less. The main beam size is 7.9'' at 450 μm and 13.0'' at 850 μm . The observations were obtained predominantly under

moderate-to-good atmospheric conditions. The zenith atmospheric opacity at 225 GHz had a median value of $\tau_{225} = 0.067$, with the 16th–84th percentile range of $\tau_{225} = 0.056$ – 0.085 . The full range of the start and end values recorded for the individual observations was $\tau_{225} = 0.001$ – 0.111 . The effective on-source exposure time for each target ranges from ~ 0.2 to ~ 3.5 h.

The SCUBA-2 data were reduced using the ORAC-DR pipeline [35] within the STARLINK 2018A software [36]. Since most of our targets are faint, we adopted the configuration file `dimmconfig_blank_field.lis`, which is ideal for faint cosmological sources and blank fields. The flux calibration was performed using the standard SCUBA-2 flux conversion factors recommended by the JCMT/Starlink pipeline. As the optical coordinates of all targets are well determined, source detection was performed around the known optical positions. We searched for submillimeter counterparts within one beam FWHM of the optical coordinates, corresponding to $7.9''$ at $450\ \mu\text{m}$ and $13.0''$ at $850\ \mu\text{m}$.

A source is considered detected if there is a cluster of pixels with a signal-to-noise ratio $\geq 3\sigma$ near the optical position and the emission morphology is consistent with the point spread function. Since the targets are expected to be unresolved point sources, the peak flux density was adopted as the flux density of the detected source. If no $\geq 3\sigma$ signal is found near the optical position, we adopted the corresponding 3σ value as the upper limit on the flux density. The observational results are listed in Table 2. At $450\ \mu\text{m}$ and $850\ \mu\text{m}$, two and seven sources are detected at $\geq 3\sigma$, respectively, while the remaining sources are assigned 3σ flux-density upper limits.

Table 2. SCUBA-2 observational data for the 44 SDSS quasars.

Name (1)	R.A. (deg) (2)	Dec. (deg) (3)	z (4)	F_{450} (mJy) (5)	F_{850} (mJy) (6)	Obs. (7)
J011430.24+000420.94	18.6260	0.0725	0.455	<136.90	<6.92	20A
J014003.99-002541.77	25.0166	−0.4283	0.770	<818.89	<7.60	20A
J015243.15+002039.65	28.1798	0.3443	0.578	<216.07	68.23 ± 2.64	20A
J015950.25+002340.82	29.9594	0.3947	0.163	<237.88	<7.23	19B
J021859.87+002855.80	34.7495	0.4822	0.351	<317.81	<8.24	20A
J090026.51+204158.70	135.1105	20.6996	0.706	<86.87	<4.51	19A
J090158.88+002313.87	135.4953	0.3872	0.196	<285.36	<6.20	19B
J092159.40+450912.38	140.4975	45.1534	0.235	126.15 ± 31.37	<7.80	19A
J092635.12+072446.43	141.6463	7.4129	0.189	<199.37	<12.00	19A
J095017.06+215022.41	147.5711	21.8396	0.455	<265.07	<7.77	20A
J105959.93+574848.17	164.9997	57.8134	0.453	<143.40	6.98 ± 1.79	20A
J110036.64+564134.89	165.1527	56.6930	0.834	148.23 ± 45.78	<6.96	20A
J111830.29+402554.02	169.6262	40.4317	0.154	<251.74	<10.60	19A
J115253.68+000131.90	178.2237	0.0255	0.823	<119.77	<5.18	20A
J120226.76-012915.28	180.6115	−1.4876	0.150	<155.18	6.52 ± 2.04	19A
J120312.14+015321.30	180.8006	1.8892	0.296	<83.32	<4.77	20A
J121037.81+053805.88	182.6575	5.6350	0.436	<87.25	<3.74	19A
J121946.54+145259.37	184.9439	14.8832	0.401	<278.92	<7.74	19A
J122011.88+020342.21	185.0495	2.0617	0.240	<236.71	131.57 ± 4.02	19A
J122102.95-000733.74	185.2623	−0.1260	0.366	<229.49	<6.12	19A
J122317.80+092306.94	185.8242	9.3853	0.682	<110.44	<4.63	19A
J122526.21+141332.24	186.3592	14.2256	0.760	<695.52	<10.10	19A
J122822.10+114606.83	187.0921	11.7686	0.365	<355.78	<8.13	19A
J122839.20+035749.29	187.1633	3.9637	0.608	<445.35	<8.94	19A
J123436.54+123918.66	188.6522	12.6552	0.777	<103.89	4.29 ± 1.32	19A
J124511.26+335610.12	191.2969	33.9361	0.711	<92.09	<4.90	20A
J125317.57+310550.64	193.3232	31.0974	0.782	<99.35	<6.32	20A

Table 2. Cont.

Name	R.A.	Dec.	<i>z</i>	<i>F</i> ₄₅₀	<i>F</i> ₈₅₀	Obs.
(1)	(deg) (2)	(deg) (3)	(4)	(mJy) (5)	(mJy) (6)	(7)
J125703.80+250457.56	194.2658	25.0827	0.821	<133.65	<4.49	19A
J125711.97+274216.45	194.2999	27.7046	0.793	<85.01	<4.19	20A
J130622.96+225752.95	196.5957	22.9647	0.758	<80.81	<6.64	20A
J131247.97+250756.71	198.1999	25.1324	0.425	<89.84	<4.87	20A
J132919.84+250626.45	202.3327	25.1073	0.781	<133.21	<6.69	20A
J140621.89+222346.54	211.5912	22.3963	0.098	<116.46	<4.95	19A
J140655.66+015712.88	211.7319	1.9536	0.427	<108.39	<4.87	20A
J140700.40+282714.65	211.7517	28.4541	0.077	<124.33	<6.70	20A
J141700.83+445606.39	214.2535	44.9351	0.113	<167.56	<5.70	19A
J145001.69+022006.75	222.5070	2.3352	0.520	<113.76	<4.31	19A
J145108.76+270926.92	222.7865	27.1575	0.064	<105.18	<4.91	19A
J145538.73+002238.06	223.9114	0.3772	0.434	<81.94	<4.67	20A
J152114.26+222743.87	230.3094	22.4622	0.136	<102.20	<4.76	19A
J161413.20+260416.21	243.5550	26.0712	0.131	<149.57	<5.52	19A
J163352.34+402115.66	248.4681	40.3544	0.782	<94.68	4.07 ± 1.30	19A
J163915.81+412833.70	249.8159	41.4760	0.690	<358.96	77.81 ± 6.01	19A
J171352.43+584201.25	258.4685	58.7003	0.521	<212.11	<5.67	19A

Note: Column (1): SDSS name. Column (2): right ascension (J2000). Column (3): declination (J2000). Column (4): redshift. Columns (5) and (6): flux densities at 450 μ m and 850 μ m, respectively; upper limits are given at the 3σ level. Column (7): observation semester.

For the PG quasars, we cross-matched our sample with the public JCMT/SCUBA-2 archival data available from the Canadian Astronomy Data Centre (CADC)². Then, we reduced the data using STARLINK. This procedure yielded SCUBA-2 data for 14 PG quasars. For the 450 μ m band, some measurements are unavailable because of the missed raw observations or unreliable reduction results. So, only the 850 μ m measurements are available for part of our PG sample. The final usable SCUBA-2 data for these nine PG quasars include three 3σ flux-density upper limits at 450 μ m and two $\geq 3\sigma$ detections plus seven 3σ flux-density upper limits at 850 μ m.

3. SED Fitting

Since quasars emission is very strong across a broad range of the electromagnetic spectrum, SED fitting can give more reliable FIR SFRs and host galaxy properties for our quasar sample. By decomposing the observed SED into different individual components, we can account for the AGN-heated hot dust emission in the MIR band and possible contamination in the radio band. Then, we can isolate the FIR cold dust emission associated with star formation.

3.1. Multiwavelength Data

We compiled multiwavelength data from optical to FIR for each quasar. For the radio-loud quasars, we also collected available radio measurements. The complete photometric data used as input for both CIGALE and AGNfitter are provided in Supplementary Table S1 in machine-readable format. The catalog products, versions, and corresponding references are summarized in Table 3. The NASA/IPAC Infrared Science Archive (IRSA) and the NASA/IPAC Extragalactic Database (NED)³ were used for data access and cross-identification.

Table 3. Photometric data products adopted in this work. Source-by-source references for radio and millimeter measurements are provided in Supplementary Table S1 in machine-readable format.

Facility	Bands	Catalog or Data Product	References
Optical to mid-infrared			
SDSS	<i>ugriz</i>	SDSS DR7 imaging photometry	[24,27]
2MASS	<i>J, H, K_s</i>	2MASS All-Sky Point Source Catalog, compiled in SDSS DR7 quasar-property catalog	[27,37,38]
WISE	W1, W2, W3, W4	WISE Preliminary Data Release, 14 April 2011	[27,39]
Spitzer	IRAC 3.6–8.0 μm ; MIPS 24 μm	SEIP Source List, Cryogenic Release 3.0	[40,41]
Far-infrared and submillimeter			
Herschel/PACS	70, 100, 160 μm	Herschel/PACS Point Source Catalogue, HPPSC v1c	[29]
Herschel/SPIRE	250, 350, 500 μm	Standard SPIRE Point Source Catalog v2c	[30]
PG infrared data	NIR to FIR	Published photometric compilation	[32]
SCUBA-2	450, 850 μm	JCMT observations and archival data	This work
Radio and millimeter			
FIRST	1.4 GHz	VLA FIRST Survey Catalog, 14 December 2017 version	[42–44]
Other radio/mm data	150 MHz–150 GHz; 1.2 mm	Individual published measurements	Supplementary Table S1

Note. Source-by-source radio and millimeter measurements are provided in Supplementary Table S1, together with their original literature sources [45–80].

3.2. SED Fitting Using CIGALE

CIGALE⁴ (Code Investigating GALaxy Emission) [81–83] is a widely used SED-fitting code. It models multiwavelength photometric data from FUV to radio and estimates physical properties, such as SFR, stellar mass, dust attenuation, dust emission, and AGN contribution. CIGALE relies on a predefined model grid, which is computed once and then applied to all sources. It is very efficient for large samples. Reliable results require sufficiently dense sampling of the relevant parameter space. However, the memory requirement increases as the number of models, photometric bands, and different physical parameters increases. Therefore, to keep the memory requirements manageable, we adopt a coarser sampling at first. Then, we further refine the grid around the preliminary best-fit values for a few key parameters. In CIGALE, the SED is constructed through a sequence of independent modules, each describing a specific physical component. These modules typically include star formation history (SFH), simple stellar population (SSP), nebular emission, dust attenuation, dust emission, AGN, X-ray, radio, rest-frame parameters, and intergalactic medium (IGM). We use the following models and parameters for each module: (1) SFH: the delayed SFH with optional exponential burst. (2) SSP: BC03 [84], with the Chabrier [85] initial mass function (IMF) and solar metallicity, $Z = 0.02$. Since the BC03 module in CIGALE does not include the Kroupa [86] IMF which is adopted elsewhere in this work, the relevant CIGALE-derived quantities are converted to that IMF following Madau and Dickinson [87]. (3) Dust attenuation: modified Calzetti et al. [88] attenuation law. (4) Dust emission: we adopt four models, dale2014 [89], dl2014 [90], casey2012 [91], and THEMIS [92]. (5) AGN emission: skirtor2016 [93,94]. (6) Radio: synchrotron emission based on Helou et al. [95]. The parameters of these models are listed in Table 4. Since the FIR emission is the primary focus of this work, we adopt four dust emission models for the dust module: dale2014, dl2014, casey2012, and THEMIS. As an example, Figure 2 presents the CIGALE SED-fitting results of J130622.96+225752.95 obtained

with the four adopted cold dust emission models. The AGNfitter decomposition of the same quasar is shown in panel (e), enabling a direct comparison between the two fitting approaches.

Table 4. CIGALE model parameters.

Parameter	Symbol	Input Values
SFH: delayed star formation history with an additional burst		
E-folding time of the main stellar population model (Myr)	τ_{main}	100, 500, 1000, 5000, 10,000
E-folding time of the late starburst population model (Myr)	τ_{burst}	20,000
Mass fraction of the late burst population	f_{burst}	0.0, 0.01, 0.1, 0.3
Age of the main stellar population in the galaxy (Myr)	t_{main}	100, 500, 1000, 5000, 10,000
Age of the late burst (Myr)	t_{burst}	20
SSP: Bruzual & Charlot [84]		
Initial mass function	–	Chabrier [85]
Metallicity	Z	0.02
Age of the separation between the young and the old stellar population (Myr)	t_{sep}	10
Dust attenuation: modified Calzetti et al. [88] attenuation law		
Color excess of the nebular line emission	$E(B - V)_{\text{lines}}$	0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, 0.9
Reduction factor applied to $E(B - V)_{\text{lines}}$ to compute the stellar continuum attenuation $E(B - V)_{\star}$	$f_{E(B-V)}$	0.44
Dust emission: dale2014 [89]		
AGN fraction	f_{AGN}	0
Alpha slope	α	1.0 to 3.0
Dust emission: dl2014 [90]		
Mass fraction of PAH	q_{PAH}	0.47 to 5.95
Minimum radiation field	U_{min}	1.0 to 30.0
Power-law slope $dU/dM \propto U^{\alpha}$	α	2.0
Fraction illuminated from U_{min} to U_{max}	γ	0.02 to 0.20
Dust emission: casey2012 [91]		
Temperature of the dust	T_{dust}	20 to 90 K
Emissivity index of the dust	β	1.0 to 3.0
Mid-infrared power-law slope	α	2.0
Dust emission: THEMIS [92]		
Mass fraction of hydrocarbon solids, i.e., a-C(:H) smaller than 1.5 nm, also known as HAC	q_{HAC}	0.02 to 0.32
Minimum radiation field	U_{min}	1.0 to 30.0
Power-law slope $dU/dM \propto U^{\alpha}$	α	2.0
Fraction illuminated from U_{min} to U_{max}	γ	0.02 to 0.2
AGN emission: skitor2016 [93,94]		
Average edge-on optical depth at 9.7 μm	τ	3 to 11
Power-law exponent that sets radial gradient of dust density	p	1.0
Index that sets dust density gradient with polar angle	q	1.0
Angle measured between the equatorial plane and edge of the torus	θ	40
Ratio of outer to inner radius, $R_{\text{out}}/R_{\text{in}}$	R	20
Fraction of total dust mass inside clumps	M_{cl}	0.97
Inclination, i.e., viewing angle	i	0 to 50
AGN fraction	f_{AGN}	0.01 to 0.99
Polar-dust color excess	$E(B - V)_{\text{PD}}$	0, 0.2, 0.4
Radio		
The value of the FIR/radio correlation coefficient for star formation	$q_{\text{IR,SF}}$	2.58
The slope of the power-law synchrotron emission related to star formation	α_{SF}	0.8
The radio-loudness parameter for AGN	R_{AGN}	0.01 to 5000
The slope of the power-law AGN radio emission, assumed isotropic	α_{AGN}	0.7 or 0.6, 0.7, 0.8, 0.9, 1.0




Figure 2. Comparison of the SED decompositions of J130622.96+225752.95 obtained with CIGALE and AGNfitter. (a–d) show the CIGALE results obtained with the four adopted cold dust emission models: *dale2014*, *d12014*, *casey2012*, and *THEMIS*. (e) shows the AGNfitter result for the same quasar. All panels use the same observed photometry and upper limits. The lower panels show the relative residuals. The component colors follow the native output conventions of the two fitting codes; their physical meanings are indicated in the individual panels.

3.3. SED Fitting Using AGNfitter

AGNfitter⁵ [96,97] is an SED-fitting code specifically designed for AGNs. It can model multiwavelength photometric data from the radio, submillimeter, IR, optical, and UV bands to the X-ray. It uses a Bayesian MCMC approach to decompose the observed SED into the physical components associated with the AGN and its host galaxy. This Bayesian framework estimates model parameters and their uncertainties from posterior probability distribution functions (PDFs). This provides more information than a single best-fit solution. Owing to its computational efficiency, AGNfitter is very suitable for statistical studies of large samples of AGNs and their host galaxies.

The SED components of AGNfitter include stellar population, cold dust emission from star-forming regions, AGN dusty torus, accretion disk, and radio emission. Since no X-ray

data are used in this work, we do not include the X-ray component. AGNfitter allows for theoretical, empirical, and semi-empirical templates for different components. The models and parameter ranges we used for each component are summarized in Table 5. These components include (1) stellar emission: BC03 stellar population synthesis models [84]; (2) cold dust emission: S17 model [98]; (3) AGN dusty torus: SKIRTOR model [94]; (4) accretion disk: THB21 model [99]; and (5) radio emission: a simple power-law (SPL) model. As an example, AGNfitter fitting result for the quasar J130622.96+225752.95 shows in Figure 2e.

Table 5. Models and parameters adopted in AGNfitter.

Component	Model	Parameter	Description	Range
Stellar emission	BC03	τ	Time scale of the exponential SFH [log year]	[0.05, age(z)]
		age	Galaxy age [log year]	[7, age(z)]
		$E(B - V)_{\text{gal}}$	Galaxy reddening	[0, 0.6]
		GA	Galaxy normalization	[−10, 10]
Cold dust emission	S17	T_{dust}	Dust temperature [K]	[14.24, 42]
		fracPAH	PAH fraction	[0, 0.05]
		SB	Starburst normalization	[−10, 10]
Torus emission	SKIRTOR	incl	Torus inclination angle [°]	[0, 90]
		oa	Opening angle [°]	[10, 80]
		τ_{ν}	Optical depth at 9.7 μm	[3, 11]
		TO	Torus normalization	[−10, 10]
Accretion disk	THB21	$E(B - V)_{\text{bbb}}$	BBB reddening	[0, 1]
		BB	BBB normalization	[−10, 10]
Radio	SPL	α	Slope of synchrotron emission [log ν]	[−2, 1]
		RAD	Radio normalization	[−10, 10]

4. Results and Discussion

4.1. FIR SFR Estimates and Model Comparison

We estimate the FIR SFRs of the quasar host galaxies using the infrared luminosity–SFR calibration of Kennicutt [10], with the coefficient converted to the Kroupa [86] IMF following Madau and Dickinson [87]:

$$\text{SFR} (M_{\odot} \text{ yr}^{-1}) = 3.02 \times 10^{-44} L_{\text{FIR}} (\text{erg s}^{-1}). \quad (1)$$

Here, L_{FIR} is measured by integrating the cold dust component obtained from the SED fitting over the infrared wavelength range of 8–1000 μm . For each quasar, we obtain five FIR SFR estimates: four from CIGALE using different cold dust templates, and one from AGNfitter.

Figure 3 compares the FIR SFRs of five different models of each source. The mean SFR of the five estimations is shown on the abscissa, and each FIR SFR of the five different models are shown on the ordinate.

To quantify the model-dependent uncertainty of the FIR SFRs, we calculated, for each quasar, the standard deviation of the logarithmic SFR estimates obtained from the four CIGALE cold dust templates and AGNfitter. Among the four CIGALE templates, the median source-by-source scatter is 0.10 dex, corresponding to a multiplicative factor of 1.26 or an approximately 26% difference in SFR. The 16th–84th percentile range is 0.02–0.23 dex. Relative to the mean CIGALE estimate, AGNfitter gives a median offset of −0.09 dex, corresponding to an SFR approximately 19% lower. When all five SFR estimates are considered together, the median model-dependent scatter is 0.14 dex, corresponding to a factor of 1.38 or approximately 38%. The 16th–84th percentile range is 0.06–0.24 dex.

Therefore, for quasars with photometric coverage and SED assumptions comparable to those adopted here, a systematic uncertainty of approximately 0.14 dex, or about 38%, provides a practical estimate of the uncertainty associated with the choice of SED-fitting code and cold dust template. This uncertainty reflects model dependence and should be considered separately from the formal fitting uncertainty within an individual model.




Figure 3. Comparison of the five FIR SFR estimates. The abscissa shows the mean SFR, while the ordinate shows the FIR SFR derived from each model. The blue, green, red, and yellow points represent the CIGALE results obtained with the casey2012, dale2014, dl2014, and THEMIS cold dust templates, respectively. The orange points represent the AGNfitter results. The lower panel shows the residuals relative to the mean SFR.

The selected SED-fitting results are summarized in Table 6. Only a portion of the table is shown here; the full table for all the 202 quasars is provided in Appendix A. When the same parameter is produced by multiple templates, we adopt the average value as the final estimate.

4.2. Effect of SCUBA-2 Data on the SED-Fitting Results

To examine the impact of the SCUBA-2 observations on the SED-fitting results, we performed two fits for each source with each template, one including and the other excluding the SCUBA-2 data. In total, five models were considered: the four CIGALE cold dust templates and AGNfitter. Since the influence of the SCUBA-2 data may depend on the available far-infrared constraints and radio properties, we divided the full sample into subsamples according to the number of detected Herschel bands and the radio classification.

We first compare the FIR SFRs derived with and without the SCUBA-2 data. For each source, the FIR SFR is taken as the average value from the five models. The result is shown in Figure 4. The abscissa shows the FIR SFR obtained from the SED fitting including the SCUBA-2 data, while the ordinate shows the FIR SFR obtained without the SCUBA-2 data, together with the corresponding residuals. Red symbols indicate sources detected in no more than two of the six Herschel bands, while blue symbols indicate sources detected

in three or more Herschel bands. Triangles denote radio-loud sources, and circles denote radio-quiet sources. For the full sample and for each subsample divided by the number of Herschel detections and radio classification, we calculate the mean difference between the results obtained without and with the SCUBA-2 data. These mean offsets are labeled in the upper left and lower right corners of the figure. The same color and symbol scheme is used in the following comparisons of the other five model parameters.

Table 6. Selected SED-fitting results for the quasar sample. Only a portion of the table is shown here; the full table for all 202 quasars is provided in Appendix A.

Name	z	$\log M_{\star}$ ($M_{\odot}$)	$\log \text{SFR}_{\text{FIR}}$ ($M_{\odot} \text{ yr}^{-1}$)	T_{dust} (K)	U_{min}	f_{AGN}	$\log M_{\text{BH}}$ ($M_{\odot}$)	$\log \lambda_{\text{Edd}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
J001115.69+011459.23	0.580	10.86 ± 0.53	2.34 ± 0.15	43.52 ± 2.99	14.53 ± 8.68	0.03 ± 0.01	8.37	−0.95
J002421.84+002508.49	0.599	10.22 ± 0.12	2.12 ± 0.15	30.89 ± 3.56	2.73 ± 2.37	0.29 ± 0.05	7.36	−0.16
J005905.51+000651.67	0.719	11.56 ± 0.31	2.00 ± 0.38	45.86 ± 9.04	11.14 ± 8.83	0.70 ± 0.07	8.94	−0.64
J011430.24+000420.94	0.455	11.70 ± 0.08	1.43 ± 0.14	21.02 ± 3.12	1.18 ± 1.02	0.20 ± 0.07	8.42	−1.32
J011536.92-000011.01	0.532	11.05 ± 0.18	1.72 ± 0.14	21.92 ± 2.72	1.06 ± 0.40	0.34 ± 0.09	8.24	−1.15

Note: Column (1): source name. Column (2): redshift. Column (3): stellar mass. Column (4): FIR SFR derived from the cold dust component integrated over 8–1000 μm . Column (5): dust temperature. Column (6): minimum radiation field intensity. Column (7): fractional AGN contribution to the total infrared luminosity, measured over 8–1000 μm ; when a parameter is returned by multiple SED-fitting templates, we report the average value. Columns (8) and (9) give the black hole mass and Eddington ratio, respectively. For the SDSS quasars, M_{BH} and L_{bol} are adopted from Shen et al. [27]; for the PG quasars, they are adopted from Shangguan et al. [32]. The Eddington ratios are calculated as $\lambda_{\text{Edd}} = L_{\text{bol}}/L_{\text{Edd}}$.

Unless otherwise specified, the comparison with SCUBA-2 data includes all 58 quasars with available SCUBA-2 observations, including both direct detections and sources with SCUBA-2 upper limits. We define a SCUBA-2 detection as an $S/N \geq 3$ in either the 450 or 850 μm band. The remaining sources with SCUBA-2 coverage are treated as upper-limit-only sources.

The FIR SFRs derived without the SCUBA-2 data are, on average, lower by only 0.01 dex than those obtained when the SCUBA-2 data are included. Sources with fewer Herschel detections show a larger scatter than those with better Herschel coverage. Similarly, radio-loud sources exhibit a larger scatter than radio-quiet sources. In particular, for some radio-loud sources, the difference between the two FIR SFR estimates can reach ~ 0.12 dex. Overall, including the SCUBA-2 data has only a minor effect on the mean FIR SFR of the full sample. However, its impact can be appreciable for individual sources with limited Herschel detections or for radio-loud quasars.

To assess whether directly measured submillimeter fluxes provide stronger constraints than upper limits alone, we repeated the comparison separately for the SCUBA-2-detected and upper-limit-only subsets. The results are summarized in Table 7. The mean change in the adopted FIR SFR is close to zero for both subsets. The SCUBA-2-detected sources show a substantially larger source-to-source scatter, 0.059 dex, than the upper-limit-only sources, 0.011 dex. Their individual changes range from -0.110 to $+0.133$ dex, whereas the upper-limit-only sources show a much narrower range of -0.011 to $+0.058$ dex. Thus, direct SCUBA-2 detections do not introduce a significant systematic shift in the mean FIR SFR but can lead to larger adjustments in the fitted FIR SFRs for individual sources.

In addition to the FIR SFR, we compare the dust-related parameters derived with and without the SCUBA-2 data for the five SED-fitting models, including the four CIGALE cold dust templates and AGNfitter. The detailed comparisons are presented in Appendix B.




Figure 4. Comparison of the FIR SFRs derived from SED fitting with and without the SCUBA-2 data. The abscissa shows the FIR SFR obtained when the SCUBA-2 data are included, while the ordinate shows the FIR SFR obtained when the SCUBA-2 data are excluded. Red symbols indicate sources detected in no more than two of the six Herschel bands, and blue symbols indicate sources detected in three or more Herschel bands. Triangles and circles represent radio-loud and radio-quiet sources, respectively. The mean offsets for the full sample and different subsamples are labeled in the upper left and lower right corners.

Table 7. Effect of SCUBA-2 constraints on the adopted FIR SFR. We define $\Delta \log \text{SFR}_{\text{FIR}} = \log \text{SFR}_{\text{with SCUBA}} - \log \text{SFR}_{\text{no SCUBA}}$. A SCUBA-2 detection is defined as an $S/N \geq 3$ in either the 450 or 850 μm band. The quoted uncertainties are the standard deviations among sources.

Subsample	<i>N</i>	Mean $\Delta \log \text{SFR}_{\text{FIR}}$ (dex)	Median (dex)	Range (dex)
All observed	58	0.006 ± 0.026	0.003	−0.110 to +0.133
Detected	11	0.008 ± 0.059	0.001	−0.110 to +0.133
Upper-limit only	47	0.006 ± 0.011	0.003	−0.011 to +0.058

We can see that excluding the SCUBA-2 data has only a minor effect on the mean dust parameter estimates for the full sample. However, the impact becomes more noticeable for sources with limited Herschel detections and for radio-loud quasars. For example, when the SCUBA-2 data are excluded, the casey2012 dust temperature can differ by up to ~ 18 K for individual sources, and the difference in $\log U_{\text{min}}$ for the dl2014 template can reach ~ 0.7 dex for some radio-loud quasars. Similar behavior is found for the THEMIS and AGNfitter results. These results indicate that the SCUBA-2 data do not significantly change the average properties of the full sample, but they provide useful additional constraints for individual sources, especially for those with sparse far-infrared coverage or radio-loud sources.

Although the comparison above quantifies the effect of including the SCUBA-2 measurements in the SED fitting, it does not address a separate observational uncertainty associated with the relatively large beams at FIR and submillimeter wavelengths. The Herschel/SPIRE and SCUBA-2 fluxes may include contributions from nearby dusty galaxies

that are physically associated with the quasar or are unrelated projected sources. Such source confusion is expected to be more relevant at the higher-redshift end of the sample, where the physical area covered by a beam is larger. High-resolution ALMA observations of FIR-bright SDSS quasars at $1 < z < 4$ have shown that multiple submillimeter counterparts can occur within a single SPIRE beam [100].

We visually inspected the SCUBA-2 maps and did not identify obvious cases with multiple resolved submillimeter peaks near the target positions. Nevertheless, the angular resolutions of SPIRE and SCUBA-2 are insufficient to exclude unresolved companions or projected dusty galaxies. In particular, the SCUBA-2 beam FWHM values are $7.9''$ at $450\ \mu\text{m}$ and $13.0''$ at $850\ \mu\text{m}$, so optically faint or obscured FIR contributors may remain unresolved. We therefore do not apply a uniform correction to the SPIRE or SCUBA-2 fluxes. The quantitative results of Hatziminaoglou et al. [100] cannot be directly transferred to our sample because their quasars are at substantially higher redshifts and were selected to be FIR bright, whereas our sample spans $0.02 < z \lesssim 0.8$. Source confusion should nevertheless be regarded as an additional source-level systematic uncertainty, particularly for the highest-redshift and most FIR-luminous quasars. It may contribute to the scatter in the inferred FIR luminosities, SFRs, and dust parameters, but spatially resolved submillimeter observations are required to quantify its effect for individual sources.

4.3. Main-Sequence Offset and Black Hole Accretion

4.3.1. Main-Sequence Location of Quasar Host Galaxies

Using SED fitting, we obtain far-infrared star formation rates and stellar masses for the 202 quasars in our sample. These measurements allow us to investigate the connection between black hole accretion and galaxy evolution by examining the locations of quasar host galaxies relative to the star-forming main sequence (MS). The possible model-related scatter in stellar mass and its impact on the following analysis are further discussed in Section 4.5.

The main sequence describes a tight correlation between star formation rate and stellar mass for star-forming galaxies. It has been observed over a wide range of redshifts [101–106]. Previous studies have found different results regarding the locations of quasar host galaxies relative to the MS. Some studies found quasar hosts to lie below the MS [107,108] and some found them to be consistent with the MS [109–111], while others found them to lie above the MS [112–115]. These discrepancies may be caused by many things: differences in sample selection, uncertainties and methodological differences in SFR and stellar mass measurements, the adopted shape and normalization of the MS relation.

The locations of our quasar host galaxies on the MS are shown in Figure 5. The black dashed line and gray shaded region represent the MS relation at $z = 0$ and its width (± 0.4 dex) from Popesso et al. [101], while the orange dashed line and yellow shaded region show the corresponding relation and width at $z = 0.5$. Blue points denote quasars at $z < 0.5$, and red points denote quasars at $z > 0.5$.

For the $z < 0.5$ subsample, most quasar hosts lie on or above the MS. The $z > 0.5$ subsample is located at systematically higher SFRs, with most sources lying above the MS. However, the apparent high-redshift excess should be interpreted with caution because the SDSS subsample is selected to have detections in at least two Herschel bands. We quantify the resulting redshift-dependent FIR selection effect in Section 4.3.3.

Since this redshift division is relatively coarse, and Popesso et al. [101] provide an MS relation that explicitly evolves with redshift, we further quantify the offset of each quasar from the MS at its own redshift. This reduces the effect of redshift evolution. We define the MS offset as

$$\Delta_{\text{MS}} = \log \text{SFR} - \log \text{SFR}_{\text{MS}}, \quad (2)$$

where $\log \text{SFR}_{\text{MS}}$ is the SFR expected from the MS relation at the redshift and stellar mass of each quasar.




Figure 5. Relation between star formation rate and stellar mass for the host galaxies of the 202 quasars. The black and orange dashed lines show the star-forming main-sequence relations at $z = 0$ and $z = 0.5$ from Popesso et al. [101], respectively. The gray and yellow shaded regions indicate the corresponding main-sequence widths. The quasars are divided into two redshift bins: $z < 0.5$ sources are shown as blue points, and $z > 0.5$ sources are shown as red points.

4.3.2. Star Formation and Black Hole Accretion

Several studies of AGNs with different luminosities have reported correlations between host galaxy star formation and AGN luminosity, black hole accretion rate, or Eddington ratio [18,21–23]. Therefore, we compare Δ_{MS} with the black hole accretion rate and Eddington ratio, as shown in Figure 6. The sample is divided into four redshift bins, $z < 0.1$, $0.1 < z < 0.3$, $0.3 < z < 0.5$, and $z > 0.5$, shown in blue, green, orange, and red, respectively. In each redshift bin, the sources are further divided into three bins along the abscissa, and the median trends are shown. The left panel of Figure 6 shows that a weak positive correlation is seen only in the lowest-redshift bin, $z < 0.1$, whereas the higher-redshift bins show little or no correlation. Therefore, the full-sample trend should be interpreted with caution. The right panel shows that Δ_{MS} is largely uncorrelated with the Eddington ratio, both for the full sample and within individual redshift bins.

The FIR SFR traces star formation averaged over relatively long timescales, whereas L_{bol} , BHAR, and the Eddington ratio trace more instantaneous black hole activity. The weak or nearly flat relation between AGN luminosity and the SFR of host galaxies does not necessarily mean there is no physical connection between black hole growth and star formation. Instead, it is more likely due to the strong variability in the AGN accretion rate on short timescales, whereas the SFR reflects the average star formation activity over longer timescales. (e.g., [19,20]). In addition, although we account for the redshift evolution of the main sequence using the relation of Popesso et al. [101], the higher-redshift quasars in our sample still tend to have larger Δ_{MS} . This may be affected by the residual redshift dependence or sample-selection effects.

We also compare the sSFR with the black hole accretion rate and Eddington ratio in Figure 7, using the same redshift bins as in Figure 6. The sSFR–BHAR relation is affected by redshift, as higher-redshift quasars tend to have both a higher sSFR and higher BHAR. However, the trends within individual redshift bins are weak or do not show. The sSFR–Eddington ratio relation also does not show a clear trend for the full sample or within individual redshift bins.




Figure 6. Relation between main-sequence offset, Δ_{MS} , and (a) black hole accretion rate and (b) Eddington ratio. The sample is divided into four redshift bins: $z < 0.1$ (blue), $0.1 < z < 0.3$ (green), $0.3 < z < 0.5$ (orange), and $z > 0.5$ (red). In each redshift bin, the sources are further divided into three bins along the abscissa, and the median trends are shown.




Figure 7. Relation between specific star formation rate and (a) black hole accretion rate and (b) Eddington ratio. The sample is divided into four redshift bins: $z < 0.1$ (blue), $0.1 < z < 0.3$ (green), $0.3 < z < 0.5$ (orange), and $z > 0.5$ (red). In each redshift bin, the sources are further divided into three bins along the abscissa, and the median trends are shown.

Therefore, for the nearby quasar subsample at $z < 0.1$, there is a weak positive trend between the star formation of host galaxies and the black hole accretion. But for quasars at $z > 0.1$, our data do not show clear evidence for a direct, instantaneous connection between global star formation and current black hole accretion activity. However, these results do not rule out a long-term coevolution between quasars and their host galaxies. Instead, connection between star formation and black hole growth is likely affected by

redshift evolution, sample selection, and the different timescales between FIR SFR and AGN accretion indicators. For samples spanning a broad redshift range, these effects should be carefully controlled when we study the relation between star formation and central black hole activity.

As an additional test, we examined the relation between Δ_{MS} and two infrared-based AGN indicators derived from AGNfitter: the torus luminosity integrated over 1–30 μm , L_{tor} , and the ratio $L_{\text{tor}}/L_{\text{Edd}}$. These quantities are expected to be less affected by short-timescale optical variability than optical-continuum-based AGN luminosities. The results are shown in Figure 8, and the Kendall- τ test results are listed in Table A2.




Figure 8. Relation between the main-sequence offset, Δ_{MS} , and infrared-based AGN indicators derived from AGNfitter: (a) the torus luminosity integrated over 1–30 μm , L_{tor} , and (b) the torus-to-Eddington luminosity ratio, $L_{\text{tor}}/L_{\text{Edd}}$. The color scheme and redshift bins are the same as those in Figure 6. In each redshift bin, the sources are further divided into three bins along the abscissa, and the median trends are shown. The Kendall- τ correlation coefficients and the corresponding two-sided p -values are listed in Table A2.

For L_{tor} , Δ_{MS} shows positive correlations in the $z < 0.1$, $0.1 < z < 0.3$, and $z > 0.5$ bins, while no significant trend is found at $0.3 < z < 0.5$. The result is clearer when the torus luminosity is normalized by the Eddington luminosity. A positive correlation between Δ_{MS} and $L_{\text{tor}}/L_{\text{Edd}}$ is found in all four redshift bins. The relation is strongest at $0.3 < z < 0.5$, while it is weaker at $z > 0.5$.

We also tested whether these results could be caused by residual redshift trends within each redshift bin. The quantity $\log(L_{\text{tor}}/L_{\text{Edd}})$ does not show a significant correlation with redshift in any individual bin. Therefore, the positive $\Delta_{\text{MS}}-L_{\text{tor}}/L_{\text{Edd}}$ relation cannot be explained simply by redshift mixing within the bins. In addition, we recalculated Δ_{MS} using only the CIGALE-derived FIR SFRs and stellar masses. The correlations with the AGNfitter torus quantities remain qualitatively similar, suggesting that they are not mainly caused by covariance within one SED-fitting method.

We note that $L_{\text{tor}}/L_{\text{Edd}}$ is not a direct measurement of the Eddington ratio. It may also depend on torus geometry, covering factor, and uncertainties in the SED decomposition. In addition, the redshift-dependent FIR selection discussed in Section 4.3.3 should be considered when interpreting these relations.

We note that the Popesso et al. MS relation is based on a Chabrier IMF, whereas our FIR SFR calibration is converted to a Kroupa IMF. The difference between these two IMFs is small, at the level of $\lesssim 0.05$ dex [87]. Moreover, the IMF conversion would shift both the SFR and stellar mass in nearly the same direction, so the relative position with respect

to the MS and the derived Δ_{MS} values are only weakly affected. This difference is much smaller than the typical uncertainties in our SED-based SFRs and stellar masses. Therefore, the IMF difference is expected to have only a minor effect on the relative location of our quasars with respect to the MS and on the derived Δ_{MS} values.

More importantly, the MS comparison should be interpreted in the context of our FIR-constrained sample. For the SDSS subsample, we require detections in at least two Herschel bands, which naturally favors quasars with stronger FIR emission and higher FIR-based SFRs. Quasars without sufficient Herschel detections, which may have lower cold dust luminosities and lower SFRs, are not included in this comparison. Therefore, our result should not be interpreted as a general statement that all quasars lie on or above the MS. Instead, it shows that many FIR-constrained quasars in our sample are located on or above the MS. Although the PG quasars generally have better FIR coverage, the effects of FIR non-detections and upper limits should still be considered when drawing conclusions about the overall quasar population.

4.3.3. Redshift-Dependent FIR Selection Effects

Because the SDSS subsample requires detections in at least two Herschel bands, its effective FIR sensitivity is expected to vary with the redshift. To quantify this effect, we estimated an approximate two-band Herschel sensitivity for each source using its fitted SED and the available Herschel photometric uncertainties. We converted the corresponding FIR luminosity limit into an SFR limit and evaluated the associated MS offset at the median redshift and stellar mass in each redshift bin. The results are summarized in Table 8.

Table 8. Approximate effective two-band Herschel sensitivity of the SDSS subsample. For each source, we estimate the FIR luminosity required to reach the adopted detection threshold in at least two Herschel bands using its fitted SED and photometric uncertainties. The corresponding limiting SFR is evaluated at the median redshift and stellar mass in each redshift bin. We define $\Delta_{\text{MS,lim}} = \log \text{SFR}_{\text{lim}} - \log \text{SFR}_{\text{MS}}(M_{\star,\text{med}}, z_{\text{med}})$. These values are approximate effective selection thresholds rather than formal completeness limits.

Redshift Bin	<i>N</i>	<i>z</i> _{med}	log <i>M</i> _{★,med} (<i>M</i> _☉)	log SFR _{lim} (<i>M</i> _☉ , yr ^{−1})	Δ _{MS,lim} (dex)
<i>z</i> < 0.1	5	0.09	10.24	−0.20	−0.42
0.1 < <i>z</i> < 0.3	30	0.18	10.60	0.70	0.19
0.3 < <i>z</i> < 0.5	33	0.37	10.30	1.43	0.86
<i>z</i> > 0.5	47	0.70	10.69	1.93	0.87

The estimated selection threshold is well below the MS at *z* < 0.1, with $\Delta_{\text{MS,lim}} \simeq -0.42$ dex. It increases to $\Delta_{\text{MS,lim}} \simeq 0.19$ dex at $0.1 < z < 0.3$ and reaches approximately 0.86–0.87 dex at *z* > 0.3. These values are approximate effective thresholds rather than formal completeness limits, because the Herschel band coverage and depths vary among sources. Nevertheless, they indicate that the SDSS selection becomes progressively less sensitive to MS and below-MS hosts with increasing redshift.

Therefore, the elevated Δ_{MS} distribution at high redshift in Figure 5 should not be interpreted as evidence that the full quasar-host population has intrinsically enhanced SFRs. This effect should also be considered when interpreting the $\Delta_{\text{MS}}\text{--BHAR}$ and $\Delta_{\text{MS}}\text{--}\lambda_{\text{Edd}}$ relations in Figure 6, as well as the sSFR trends in Figure 7. In particular, the higher-redshift bins mainly represent the FIR-selected, high-SFR portion of the quasar-host population, and may not be representative of quasars without Herschel detections and with lower SFRs. This selection effect is also relevant to FIR-selected quasar samples at higher redshift, which preferentially include FIR-luminous hosts with high SFRs [116].

4.4. Radiation Field Intensity and Dust Heating

Based on the radiation field intensity, $U_{\min}$, derived from the SED fitting, we compare $U_{\min}$ with the quasar bolometric luminosity, L_{bol} , and the cold dust temperature, T_{dust} , in Figure 9. The Kendall- τ coefficient and the corresponding p value are shown in the lower right corner of each panel. To better illustrate the overall trends, we divide the full sample into three bins and show the median values as blue points.




Figure 9. Comparison of $U_{\min}$ with (a) bolometric luminosity, L_{bol} , and (b) cold dust temperature, T_{dust} . The Kendall- τ coefficient and the corresponding p value are shown in the lower-right corner of each panel. Blue points show the median values in three bins of the full sample.

The left panel shows that $U_{\min}$ tends to increase with increasing L_{bol} , while the right panel shows a positive correlation between $U_{\min}$ and T_{dust} . Since $U_{\min}$ traces the intensity of the radiation field heating the diffuse dust, and T_{dust} is also sensitive to dust heating, the positive correlation between these two quantities is expected. More importantly, the correlation between $U_{\min}$ and L_{bol} suggests that more luminous quasars may expose the cold dust in their host galaxies to a stronger radiation field.

This trend is consistent with the scenario in which AGN radiation contributes to heating dust on host galaxy scales. Shangguan et al. [32] found that quasar host galaxies tend to have higher $U_{\min}$ values than normal galaxies. It suggests that the enhanced radiation field may be caused by AGN heating of the interstellar medium. They also found that the inferred cold dust temperature may be biased high if the FIR SED contains a contribution from warmer dust. Several recent studies further support the possibility that AGN radiation can heat dust on scales larger than the classical torus. Schneider et al. [117] used radiative transfer modeling of the $z \sim 6$ quasar SDSS J1148+5251 and found that AGN radiation may power a substantial fraction of the observed FIR emission. McKinney et al. [118] showed that in dust-enshrouded AGN, host galaxy-scale diffuse dust heated by the AGN can increase the FIR flux at $\lambda \gtrsim 100 \mu\text{m}$. Di Mascia et al. [119] further showed that AGN radiation can heat ISM dust in simulated $z \sim 6$ quasar hosts and lead to overestimated TIR-based SFRs if this contribution is ignored.

Our results are broadly consistent with these results, but the correlation should be interpreted carefully. Our SED fitting does not provide spatial information, so we cannot determine where the dust is heated. If a large fraction of the cold dust is located far from the nucleus in star-forming regions, it may be heated mainly by local star formation and only weakly affected by AGN radiation. The enhanced $U_{\min}$ may instead be related to dust closer to the central region, more compact star formation, or both. In addition, the studies mentioned above mostly focus on very luminous, high-redshift, or dust-enshrouded

systems, while our sample consists mainly of lower-redshift type-I quasars. In addition, other factors may also contribute to this trend, such as redshift evolution, sample selection, enhanced star formation, and degeneracies in the SED fitting. Therefore, this relation provides suggestive evidence for a possible AGN effect, but it does not prove that AGN heating is the only origin of the enhanced radiation field.

4.5. Possible Effects of Stellar Mass

Stellar mass may also affect the relations discussed above, because it is connected with black hole mass and may therefore indirectly affect the correlations involving AGN luminosity or black hole accretion rate. We make additional tests about stellar mass. Before examining stellar mass-dependent trends, we first estimate the model related scatter in the stellar masses. We compared the stellar masses obtained from different cold dust templates. The median scatter among these stellar masses is about 0.24 dex. This shows that the stellar masses of type-I quasars are also model dependent. It is likely because the bright AGN emission can contaminate the optical and NIR continuum, making it difficult to accurately separate the host galaxy stellar component in the SED fitting. This scatter may introduce additional uncertainty in the sSFR and Δ_{MS} , so the related trends should be interpreted statistically. We divide the sample into three stellar mass bins ($\log(M_*/M_\odot) < 10.5$, $10.5 < \log(M_*/M_\odot) < 11.0$, $\log(M_*/M_\odot) > 11.0$) and repeat related analysis in these bins.

Figure 10a shows the relation between Δ_{MS} and the BHAR in three stellar mass bins. The median trends are not same among the different stellar mass bins, and the scatter is large. The low and intermediate mass subsamples show an increase in Δ_{MS} from low-to-moderate BHAR, while the high mass subsample shows a weaker trend and a generally lower Δ_{MS} at a given BHAR.




Figure 10. (a) Relation between the main-sequence offset, Δ_{MS} , and black hole accretion rate in three stellar mass bins. (b) Relation between the minimum radiation field intensity, U_{min} , and AGN bolometric luminosity in the same stellar mass bins. Blue, green, and red symbols represent sources with $\log(M_*/M_\odot) < 10.5$, $10.5 < \log(M_*/M_\odot) < 11.0$, and $\log(M_*/M_\odot) > 11.0$, respectively. Large squares and connecting lines show the median values in bins along the abscissa, and the error bars indicate the corresponding scatter.

Figure 10b shows the relation between U_{min} and L_{bol} in the same stellar mass bins. The increasing trend of U_{min} with L_{bol} is still visible within different stellar mass bins, suggesting that this relation is unlikely to be solely driven by stellar mass. We also directly checked the relations between stellar mass and dust-heating parameters. Stellar mass

does not show a positive correlation with either $U_{\min}$ or T_{dust} . Therefore, although stellar radiation is certainly an important heating source for cold dust, the observed $U_{\min}$ – L_{bol} relation may also include a contribution from AGN radiation.

Additional stellar mass binned tests are shown in Appendix D. The relation between Δ_{MS} and the Eddington ratio also does not show the same trend in different stellar mass bins. In addition, the positive relation between T_{dust} and $U_{\min}$ remains visible within different stellar mass bins, indicating that this dust parameter relation is not mainly driven by stellar mass.

4.6. Effects of FIR Data Quality

The SDSS subsample was originally selected to have detections in at least two Herschel bands. However, the long-wavelength FIR coverage still varies substantially among the sources. To test whether the main results are strongly affected by sources with limited SPIRE constraints, we define two higher-quality SPIRE subsamples within the 115 unique SDSS quasars. The first requires an $S/N_{250} \geq 5$ at 250 μm , while the second requires detections with an $S/N \geq 3$ in at least two SPIRE bands. Each subsample contains 51 quasars, and they overlap in 35 objects.

Table 9 compares the full SDSS FIR-constrained sample with the two SPIRE-quality subsamples. Both SPIRE-selected subsamples have higher median redshifts, FIR SFRs, and main-sequence offsets than the full SDSS sample. In particular, their median $\log \text{SFR}_{\text{FIR}}$ values are about 0.4 dex higher than that of the full SDSS sample, and all selected sources lie above the adopted main sequence. Therefore, these subsets preferentially select FIR-brighter quasars and should not be regarded as unbiased representations of the full SDSS quasar population.

Table 9. Summary of the full SDSS FIR-constrained sample and the two SPIRE-quality subsamples. Values are reported as the median with the 16th and 84th percentiles. The two SPIRE-quality subsamples overlap in 35 sources. Kendall– τ statistics are calculated from individual sources.

Quantity	Full SDSS Sample	$S/N_{250} \geq 5$	≥ 2 SPIRE Detections
Number of sources	115	51	51
Median z	$0.42^{+0.31}_{-0.26}$	$0.50^{+0.26}_{-0.22}$	$0.60^{+0.17}_{-0.27}$
Median $\log(M_*/M_\odot)$	$10.65^{+0.50}_{-0.48}$	$10.71^{+0.49}_{-0.49}$	$10.85^{+0.51}_{-0.58}$
Median $\log(\text{SFR}_{\text{FIR}}/M_\odot \text{ yr}^{-1})$	$1.71^{+0.60}_{-0.54}$	$2.08^{+0.31}_{-0.63}$	$2.11^{+0.30}_{-0.63}$
Median Δ_{MS}	$1.00^{+0.34}_{-0.48}$	$1.23^{+0.22}_{-0.51}$	$1.19^{+0.19}_{-0.49}$
Fraction with $\Delta_{\text{MS}} \geq 0$	98%	100%	100%
Kendall– τ for $\log U_{\min}$ – $\log L_{\text{bol}}$	0.24	0.16	0.28
p value	$<10^{-3}$	0.103	$<10^{-2}$

The corresponding comparisons are shown in Appendix E (Figures A5–A8). The SPIRE-quality quasars remain predominantly above the star-forming main sequence, indicating that the elevated SFRs of the FIR-constrained SDSS sample are not driven solely by sources with weak long-wavelength constraints. However, because both tests are applied within the original Herschel-selected sample and preferentially retain FIR-brighter objects, they do not remove the underlying FIR selection bias. The main-sequence result should therefore still be interpreted only for quasars with adequate FIR constraints.

For the relations between Δ_{MS} and the BHAR or Eddington ratio, the two SPIRE-quality subsamples occupy broadly similar regions of parameter space to the full SDSS sample. Because of the reduced sample sizes and the remaining redshift dependence, we do not repeat the redshift-binned median-trend analysis for these subsets. The redshift-resolved results in Figures 6 and 7 therefore remain the basis for interpreting the relations between star formation and black hole accretion.

The $U_{\min}-L_{\text{bol}}$ relation remains positive in both SPIRE-quality subsamples, but its statistical significance depends on the adopted FIR-quality criterion. The $S/N_{250} \geq 5$ subsample gives $\tau = 0.16$ with $p = 0.103$, whereas the subsample with at least two SPIRE detections gives $\tau = 0.28$ with $p < 10^{-2}$. These results are qualitatively consistent with the positive trend found for the full sample but show that its statistical strength is sensitive to the FIR data selection. Thus, the observed $U_{\min}-L_{\text{bol}}$ relation should be interpreted cautiously and may also be affected by sample selection, host galaxy properties, and SED-fitting degeneracies.

4.7. Infrared Colors as Tracers of the AGN Contribution

The SED-fitting results provide the fractional AGN contribution to the total infrared luminosity, f_{AGN} , integrated over 8–1000 μm . Figure 11 shows the distribution of $f_{\text{AGN}}(8\text{--}1000\text{ }\mu\text{m})$ for the full sample and for the two redshift subsamples. The AGN contribution to the total infrared luminosity is non-negligible, especially for the $z < 0.5$ quasars. Therefore, if the total infrared luminosity is used to infer the SFR without decomposing the AGN component, the SFR may be overestimated.




Figure 11. Distribution of $f_{\text{AGN}}(8\text{--}1000\text{ }\mu\text{m})$. The gray histogram shows the full sample, while the blue and red histograms show the $z < 0.5$ and $z > 0.5$ subsamples, respectively.

Since MIR/FIR flux-density ratios trace the relative importance of AGN-heated warm dust and host galaxy cold dust, we also examine whether infrared colors can be used as empirical indicators of $f_{\text{AGN}}(8\text{--}1000\text{ }\mu\text{m})$. Among the 24 WISE-to-Herschel colors examined in this work, W2/70 μm , W3/70 μm , and W3/100 μm show the strongest correlations with f_{AGN} . These three representative relations are shown in Figure 12, while the full set of WISE-to-Herschel color comparisons is provided in Appendix F.

The observed correlations suggest that these colors can be used as practical empirical estimators of f_{AGN} for quasars with both WISE and Herschel detections. After estimating f_{AGN} , the AGN-heated dust contribution to the total infrared luminosity can be separated from the cold dust emission associated with star formation, leading to a more reliable estimate of the host galaxy FIR SFR.

However, the reliability of these color- f_{AGN} relations depends on the quality of the FIR constraints. Sources with three or more Herschel detections generally follow the fitted relations more closely, whereas sources with two or fewer Herschel detections show larger deviations and tend to have a higher inferred f_{AGN} at a given color. This offset may have

two related origins. First, these objects may be intrinsically fainter in cold dust emission. A lower cold dust luminosity would naturally increase the relative AGN contribution to the total infrared luminosity, leading to a higher inferred f_{AGN} . Second, with only a few Herschel detections, the FIR peak and long-wavelength side of the cold dust SED are poorly constrained, increasing the degeneracy between the AGN and cold dust components in the SED fitting.

Therefore, the color-based f_{AGN} indicators should be regarded as most reliable for sources with good FIR coverage. They should be used with caution for objects with two or fewer Herschel detections. In addition, the observed colors can be affected by redshift, K-correction, photometric coverage, host galaxy contamination in the shorter WISE bands, and selection effects. Since the fitted relations are calibrated using radio-quiet sources with detections in at least three Herschel bands, they may not be directly applicable to radio-loud quasars or sources with poor FIR constraints.




Figure 12. $f_{\text{AGN}}(8-1000 \mu\text{m})$ as a function of the three strongest WISE-to-Herschel color diagnostics: (a) $W2/70 \mu\text{m}$, (b) $W3/70 \mu\text{m}$, and (c) $W3/100 \mu\text{m}$. Red and black symbols denote sources detected in at least three and in no more than two Herschel bands, respectively. Filled and open symbols represent radio-quiet and radio-loud quasars, respectively. The black dashed line and blue shaded region show the Bayesian linear regression fit and its uncertainty for the radio-quiet sources with detections in at least three Herschel bands. The fitted relation, intrinsic scatter ϵ , and correlation coefficient ρ are shown in each panel.

5. Summary and Conclusions

We study the FIR SFRs and host galaxy properties of 202 SDSS and PG quasars at $0.02 < z \lesssim 0.8$ using multiwavelength SED decomposition. The photometric data cover the optical-to-FIR bands and include JCMT/SCUBA-2 observations at 450 and 850 μm . We fit the SEDs with CIGALE and AGNfitter and use different cold dust templates to estimate systematic uncertainties. Our main results are summarized as follows.

1. For each quasar, we derive five FIR SFR estimates using four CIGALE cold dust templates and AGNfitter. The median scatter among the four CIGALE template results is 0.10 dex, corresponding to approximately 26%. AGNfitter gives SFRs lower than the mean CIGALE estimate by a median of 0.09 dex, or approximately 19%. Considering all five estimates together, the median model-dependent scatter is 0.14 dex, corresponding to approximately 38%. Thus, a systematic uncertainty of about 0.14 dex should be considered for sources with photometric coverage and modeling assumptions similar to those adopted here.
2. For the 58 quasars with SCUBA-2 coverage, removing the SCUBA-2 data changes the adopted FIR SFR by only ~ 0.01 dex on average. The mean change is close to zero for both the SCUBA-2-detected and upper-limit-only subsets. However, the change for individual sources ranges from -0.11 to $+0.13$ dex and can be larger for quasars with limited Herschel coverage or radio-loud emission.

3. Within our FIR-constrained sample, many quasar hosts lie on or above the star-forming MS. However, for the SDSS subsample, the effective FIR selection threshold increases strongly with redshift, from $\Delta_{\text{MS,lim}} \simeq -0.42$ dex at $z < 0.1$ to $\simeq 0.86\text{--}0.87$ dex at $z > 0.3$. The higher-redshift bins therefore mainly represent the FIR-bright, high-SFR part of the quasar-host population. The full-sample $\Delta_{\text{MS}}\text{--BHAR}$ trend is not seen in all redshift bins, and no clear correlation is found between Δ_{MS} or sSFR and the direct Eddington ratio.
4. As an additional test, Δ_{MS} is positively correlated with the infrared-based $L_{\text{tor}}/L_{\text{Edd}}$ ratio in all four redshift bins, while the relation with L_{tor} alone is weaker. The $L_{\text{tor}}/L_{\text{Edd}}$ ratio is not a direct Eddington-ratio measurement and can also depend on torus properties and SED decomposition. The redshift-dependent FIR selection should also be considered when interpreting this relation.
5. The radiation field intensity, U_{min} , tends to increase with quasar bolometric luminosity and cold dust temperature. This suggests that more luminous quasars may be related to stronger radiation fields. However, redshift effects, enhanced star formation, and SED-fitting degeneracies may also contribute to this trend.
6. We test 24 WISE-to-Herschel colors as indicators of $f_{\text{AGN}}(8\text{--}1000\text{ }\mu\text{m})$. W2/70 μm , W3/70 μm , and W3/100 μm show the strongest correlations. These colors may provide simple estimates of the AGN infrared contribution. But they should be used carefully for radio-loud sources or objects with weak FIR constraints.

Overall, multi-component SED decomposition is important for obtaining FIR-based SFRs in quasar host galaxies. SCUBA-2 data do not strongly change the average SFRs of the SCUBA-2-observed subset but can improve the constraints for individual sources. The relations between host galaxy star formation and AGN properties should be interpreted with care because of redshift evolution, FIR selection effects, model dependence, and the different timescales traced by FIR SFRs and AGN accretion indicators.

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/universe12070213/s1>: Table S1: Machine-readable photometric data used for the SED fitting of the 202 quasars. The accompanying README file describes the table columns, units, upper-limit convention, and the meaning of the Reference column.

Author Contributions: Conceptualization, X.F. and X.-B.W.; methodology, X.F., X.-B.W., Y.F. and Y.P.; software, X.F.; validation, X.F., Y.F. and Y.P.; formal analysis, X.F.; investigation, X.F., X.-B.W., Y.F., Y.P., R.Z. and H.W.; resources, X.F. and X.-B.W.; data curation, X.F. and Y.F.; writing—original draft preparation, X.F.; writing—review and editing, X.F., X.-B.W., Y.F., Y.P., R.Z. and H.W.; visualization, X.F.; supervision, X.-B.W.; project administration, X.-B.W.; funding acquisition, X.-B.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Key R&D Program of China (2025YFA1614100, 2025YFA1614101) and the National Science Foundation of China (grant No. 12133001).

Data Availability Statement: The complete photometric data used as input for the SED fitting are provided in Supplementary Table S1 in machine-readable format. The public photometric, spectroscopic, and archival data used in this study are available from SDSS, IRSA, NED, CADC, and the relevant mission archives. Additional derived data supporting the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgments: This work was supported by the High-performance Computing Platform of Peking University. We thank the staff of the James Clerk Maxwell Telescope for their support of the SCUBA-2 observations. The James Clerk Maxwell Telescope is operated by the East Asian Observatory on behalf of the National Astronomical Observatory of Japan, Academia Sinica Institute of Astronomy and Astrophysics, the Korea Astronomy and Space Science Institute, the National Astronomical Research Institute of Thailand, and organizations in the United Kingdom and Canada.

This work is based in part on observations obtained with SCUBA-2 on the JCMT under programs M18BP050, M19AP020, M19BP041, and M20AP044. This work also makes use of archival data from the Canadian Astronomy Data Centre (CADC), operated by the National Research Council of Canada with the support of the Canadian Space Agency. This research has made use of the NASA/IPAC Infrared Science Archive (IRSA) and the NASA/IPAC Extragalactic Database (NED), which are funded by the National Aeronautics and Space Administration and operated by the California Institute of Technology. This work is based in part on observations made with the Herschel Space Observatory, an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. This publication also makes use of data products from the Wide-field Infrared Survey Explorer (WISE), a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by NASA. This work also makes use of data products from the Sloan Digital Sky Survey (SDSS), 2MASS, and the *Spitzer* Space Telescope.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

AGN	Active galactic nucleus
BHAR	Black hole accretion rate
CADC	Canadian Astronomy Data Centre
CIGALE	Code Investigating GALaxy Emission
FIR	Far-infrared
FWHM	Full width at half maximum
IMF	Initial mass function
IRAC	Infrared Array Camera
IRSA	NASA/IPAC Infrared Science Archive
JCMT	James Clerk Maxwell Telescope
MCMC	Markov chain Monte Carlo
MIPS	Multiband Imaging Photometer for Spitzer
MIR	Mid-infrared
MS	Main sequence
NED	NASA/IPAC Extragalactic Database
NIR	Near-infrared
PACS	Photoconductor Array Camera and Spectrometer
PDF	Probability distribution function
PG	Palomar–Green
SCUBA-2	Submillimetre Common-User Bolometer Array 2
SDSS	Sloan Digital Sky Survey
SED	Spectral energy distribution
SFH	Star formation history
SFR	Star formation rate
sSFR	Specific star formation rate
SMBH	Supermassive black hole
SPIRE	Spectral and Photometric Imaging Receiver
SSP	Simple stellar population
WISE	Wide-field Infrared Survey Explorer

Appendix A. SED-Fitting Results

Table A1. Selected SED-fitting results and adopted black hole properties for the 202 quasars

Name	z	$\log M_{\star}$ ($M_{\odot}$)	$\log \text{SFR}_{\text{FIR}}$ ($M_{\odot} \text{ yr}^{-1}$)	T_{dust} (K)	U_{min}	f_{AGN}	$\log M_{\text{BH}}$ ($M_{\odot}$)	$\log \lambda_{\text{Edd}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
J001115.69+011459.23	0.580	10.86 ± 0.53	2.34 ± 0.15	43.52 ± 2.99	14.53 ± 8.68	0.03 ± 0.01	8.37	−0.95
J002421.84+002508.49	0.599	10.22 ± 0.12	2.12 ± 0.15	30.89 ± 3.56	2.73 ± 2.37	0.29 ± 0.05	7.36	−0.16
J005905.51+000651.67	0.719	11.56 ± 0.31	2.00 ± 0.38	45.86 ± 9.04	11.14 ± 8.83	0.70 ± 0.07	8.94	−0.64
J011430.24+000420.94	0.455	11.70 ± 0.08	1.43 ± 0.14	21.02 ± 3.12	1.18 ± 1.02	0.20 ± 0.07	8.42	−1.32
J011536.92-000011.01	0.532	11.05 ± 0.18	1.72 ± 0.14	21.92 ± 2.72	1.06 ± 0.40	0.34 ± 0.09	8.24	−1.15
J013354.30+011034.00	0.704	11.03 ± 0.51	1.92 ± 0.24	29.08 ± 5.97	2.56 ± 2.62	0.25 ± 0.07	8.48	−1.09
J014003.99-002541.77	0.770	10.59 ± 0.17	2.40 ± 0.19	41.33 ± 4.08	2.79 ± 3.24	0.29 ± 0.07	8.76	−1.11
J015243.15+002039.65	0.578	11.74 ± 0.16	1.48 ± 0.26	17.30 ± 0.61	1.59 ± 2.43	0.11 ± 0.03	8.22	−0.96
J015950.25+002340.82	0.163	10.46 ± 0.08	2.40 ± 0.12	55.98 ± 0.22	30.00 ± 0.03	0.19 ± 0.00	8.06	−0.42
J021859.87+002855.80	0.351	10.19 ± 0.06	1.96 ± 0.18	32.74 ± 3.17	13.79 ± 7.91	0.51 ± 0.07	7.85	−0.37
J022430.61-000038.91	0.431	9.83 ± 0.26	1.41 ± 0.13	30.57 ± 4.08	2.73 ± 1.35	0.31 ± 0.08	7.40	−0.25
J090026.51+204158.70	0.706	11.20 ± 0.04	2.35 ± 0.06	42.50 ± 1.93	28.81 ± 3.69	0.63 ± 0.02	8.79	−0.60
J090158.88+002313.87	0.196	9.88 ± 0.24	1.52 ± 0.09	45.36 ± 3.68	18.82 ± 4.24	0.25 ± 0.04	8.38	−1.36
J090933.49+425346.51	0.670	11.96 ± 0.24	2.46 ± 0.14	17.19 ± 0.44	1.00 ± 0.02	0.56 ± 0.01	9.73	−2.02
J092159.40+450912.38	0.235	9.93 ± 0.07	1.88 ± 0.07	45.86 ± 0.71	30.00 ± 0.01	0.01 ± 0.00	8.64	−1.53
J092554.72+195405.13	0.192	10.98 ± 0.44	0.85 ± 0.14	57.07 ± 6.51	19.01 ± 9.30	0.93 ± 0.01	9.10	−1.19
J092635.12+072446.43	0.189	11.22 ± 0.08	1.25 ± 0.11	30.81 ± 3.91	2.85 ± 0.78	0.17 ± 0.01	9.26	−2.39
J094652.58+131953.83	0.133	10.29 ± 0.37	1.20 ± 0.19	61.04 ± 3.20	29.31 ± 2.86	0.78 ± 0.02	8.39	−0.80
J094745.14+072520.58	0.086	10.85 ± 0.01	0.04 ± 0.12	46.33 ± 3.60	3.49 ± 4.16	0.91 ± 0.01	8.23	−1.51
J095017.06+215022.41	0.455	10.28 ± 0.28	1.67 ± 0.07	33.01 ± 6.36	7.10 ± 6.23	0.50 ± 0.05	8.31	−0.54
J095240.17+515249.91	0.554	10.83 ± 0.59	1.63 ± 0.19	43.06 ± 8.28	6.64 ± 6.44	0.58 ± 0.07	8.50	−0.80
J095819.87+022903.51	0.345	10.06 ± 0.26	1.18 ± 0.04	30.39 ± 2.63	4.92 ± 0.75	0.37 ± 0.03	8.63	−1.65
J100043.14+020637.25	0.360	10.72 ± 0.11	1.14 ± 0.11	40.40 ± 5.48	9.38 ± 6.56	0.21 ± 0.04	7.89	−0.75
J100232.13+023537.33	0.658	10.98 ± 0.34	1.75 ± 0.11	40.45 ± 7.09	16.09 ± 8.13	0.32 ± 0.03	8.14	−0.42
J100420.13+051300.48	0.160	10.31 ± 0.44	0.93 ± 0.22	61.11 ± 2.96	22.30 ± 8.48	0.86 ± 0.02	7.74	−0.30
J103651.94+575950.96	0.500	10.46 ± 0.25	1.65 ± 0.13	34.05 ± 3.58	4.53 ± 3.08	0.30 ± 0.03	8.45	−1.03
J104009.33+560343.28	0.393	11.29 ± 0.05	0.98 ± 0.68	44.22 ± 11.79	11.11 ± 8.88	0.70 ± 0.04	7.89	−0.25
J104505.39+561118.34	0.428	11.04 ± 0.19	1.88 ± 0.10	47.59 ± 2.90	21.46 ± 6.41	0.01 ± 0.00	9.80	−2.48
J104739.49+563507.19	0.303	10.24 ± 0.40	1.71 ± 0.15	50.84 ± 1.91	6.32 ± 6.98	0.08 ± 0.01	8.07	−1.03
J105106.12+591625.24	0.767	10.82 ± 0.69	2.03 ± 0.16	29.89 ± 5.10	2.96 ± 2.63	0.40 ± 0.05	9.49	−1.42
J105143.89+335926.71	0.167	10.51 ± 0.20	0.86 ± 0.11	57.61 ± 6.17	10.12 ± 8.14	0.73 ± 0.03	8.27	−0.91
J105151.44-005117.66	0.359	10.83 ± 0.22	2.34 ± 0.12	65.93 ± 0.36	30.00 ± 0.22	0.60 ± 0.01	9.17	−0.72
J105705.41+580437.46	0.140	11.15 ± 0.06	1.17 ± 0.12	34.37 ± 4.61	4.72 ± 0.69	0.22 ± 0.02	7.86	−0.90
J105959.93+574848.17	0.453	10.01 ± 0.21	1.58 ± 0.18	27.65 ± 6.53	4.37 ± 4.75	0.60 ± 0.08	8.27	−0.68
J110036.64+564134.89	0.834	10.61 ± 0.18	2.42 ± 0.19	37.48 ± 6.26	9.56 ± 7.76	0.22 ± 0.05	8.38	−0.25
J111706.40+441333.30	0.144	10.79 ± 0.16	0.81 ± 0.13	65.25 ± 1.01	28.10 ± 4.82	0.93 ± 0.01	8.77	−1.13
J111830.29+402554.02	0.154	10.07 ± 0.26	1.41 ± 0.05	35.09 ± 1.54	6.31 ± 3.83	0.44 ± 0.01	8.45	−0.94
J112019.62+130320.10	0.314	10.29 ± 0.35	1.46 ± 0.12	59.43 ± 8.44	13.62 ± 8.88	0.46 ± 0.04	7.80	−0.57
J112048.99+133821.91	0.513	10.00 ± 0.19	1.60 ± 0.05	50.19 ± 10.36	4.99 ± 5.23	0.34 ± 0.03	8.63	−1.23
J112759.26+360207.00	0.667	10.61 ± 0.24	2.19 ± 0.26	29.24 ± 6.55	3.04 ± 4.31	0.38 ± 0.05	8.80	−0.98
J114121.76+014803.58	0.382	10.27 ± 0.05	1.82 ± 0.14	27.85 ± 3.06	2.75 ± 1.48	0.44 ± 0.05	8.59	−0.88
J114933.88+222227.07	0.554	10.29 ± 0.34	1.66 ± 0.07	36.79 ± 5.03	5.43 ± 6.54	0.40 ± 0.03	8.62	−0.99
J115253.68+000131.90	0.823	11.14 ± 0.28	2.23 ± 0.25	36.33 ± 7.13	7.52 ± 7.54	0.30 ± 0.10	8.50	−0.81
J120226.76-012915.28	0.150	10.21 ± 0.05	2.24 ± 0.10	37.58 ± 1.69	27.70 ± 4.84	0.14 ± 0.00	7.28	−0.26
J120312.14+015321.30	0.296	10.68 ± 0.37	1.45 ± 0.21	30.29 ± 3.74	1.99 ± 1.23	0.28 ± 0.05	8.86	−1.82
J120442.11+275411.80	0.165	10.06 ± 0.27	1.39 ± 0.14	65.72 ± 0.50	29.92 ± 1.01	0.68 ± 0.01	8.36	−1.34
J120734.63+150643.69	0.750	10.58 ± 0.32	2.07 ± 0.24	41.94 ± 12.34	11.19 ± 8.91	0.69 ± 0.06	8.63	−0.44
J121037.81+053805.88	0.436	10.07 ± 0.73	1.63 ± 0.09	46.50 ± 6.27	11.95 ± 8.56	0.33 ± 0.04	8.24	−0.95
J121728.37+065110.00	0.589	10.66 ± 0.13	1.94 ± 0.42	60.29 ± 7.12	9.53 ± 8.35	0.54 ± 0.05	8.57	−0.77
J121836.71+155908.43	0.766	10.33 ± 0.14	2.15 ± 0.11	39.50 ± 6.27	8.48 ± 6.85	0.42 ± 0.06	8.25	−0.39
J121906.57+160243.06	0.761	10.52 ± 0.17	2.28 ± 0.21	52.99 ± 10.16	22.71 ± 8.31	0.53 ± 0.06	7.93	−0.80
J121945.03+082117.95	0.228	10.41 ± 0.18	1.78 ± 0.13	57.41 ± 5.81	4.96 ± 2.33	0.22 ± 0.02	8.33	−1.18
J121946.54+145259.37	0.401	9.77 ± 0.11	1.17 ± 0.31	31.51 ± 6.09	4.81 ± 5.42	0.41 ± 0.03	8.79	−1.35
J122011.88+020342.21	0.240	11.05 ± 0.22	1.43 ± 0.32	27.19 ± 6.67	2.74 ± 3.78	1.00 ± 0.00	8.87	−0.87

Table A1. Cont.

Name	z	$\log M_{\star}$ ($M_{\odot}$)	$\log \text{SFR}_{\text{FIR}}$ ($M_{\odot} \text{ yr}^{-1}$)	T_{dust} (K)	U_{min}	f_{AGN}	$\log M_{\text{BH}}$ ($M_{\odot}$)	$\log \lambda_{\text{Edd}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
J122026.72+062748.20	0.349	10.78 ± 0.34	1.64 ± 0.09	33.74 ± 3.47	4.83 ± 2.10	0.19 ± 0.02	8.63	−1.52
J122102.50+155447.04	0.229	10.10 ± 0.63	1.18 ± 0.21	35.46 ± 7.19	4.67 ± 5.59	0.55 ± 0.06	8.00	−0.83
J122102.95−000733.74	0.366	10.71 ± 0.17	2.21 ± 0.10	50.05 ± 3.60	21.78 ± 8.18	0.19 ± 0.02	7.75	−0.33
J122106.50+114625.46	0.340	11.04 ± 0.07	1.52 ± 0.07	39.79 ± 6.39	3.72 ± 4.29	0.17 ± 0.03	8.37	−1.19
J122307.37−002124.73	0.804	11.42 ± 0.22	2.23 ± 0.23	36.72 ± 5.31	4.05 ± 5.06	0.19 ± 0.06	—	—
J122312.17+095017.72	0.277	10.21 ± 0.31	1.55 ± 0.28	46.70 ± 4.90	4.42 ± 5.58	0.47 ± 0.10	8.05	−0.63
J122317.80+092306.94	0.682	10.96 ± 0.06	2.76 ± 0.10	48.72 ± 2.06	29.99 ± 0.26	0.17 ± 0.01	8.92	−0.62
J122404.62+045637.94	0.358	10.22 ± 0.25	1.76 ± 0.24	57.12 ± 9.15	5.11 ± 5.23	0.47 ± 0.04	8.25	−0.69
J122520.13+084450.76	0.535	10.52 ± 0.22	2.06 ± 0.04	34.01 ± 1.02	5.13 ± 1.55	0.42 ± 0.02	8.66	−0.60
J122526.21+141332.24	0.760	10.61 ± 0.22	2.35 ± 0.12	42.56 ± 4.87	21.34 ± 8.11	0.18 ± 0.02	8.03	−0.48
J122641.50+055906.81	0.290	11.36 ± 0.15	1.31 ± 0.18	31.93 ± 5.44	2.69 ± 2.09	0.24 ± 0.08	8.85	−1.56
J122822.10+114606.83	0.365	11.09 ± 0.26	1.86 ± 0.07	28.26 ± 2.65	4.83 ± 0.56	0.14 ± 0.03	8.84	−1.52
J122839.20+035749.29	0.608	10.57 ± 0.14	2.20 ± 0.10	39.38 ± 4.61	5.79 ± 5.14	0.37 ± 0.03	9.03	−1.32
J123436.54+123918.66	0.777	10.59 ± 0.13	2.44 ± 0.16	52.04 ± 3.17	26.37 ± 6.30	0.23 ± 0.05	8.73	−0.99
J123800.92+621336.09	0.440	10.33 ± 0.41	1.84 ± 0.16	39.80 ± 1.74	4.94 ± 0.77	0.17 ± 0.03	7.94	−0.59
J124511.26+335610.12	0.711	10.69 ± 0.24	2.19 ± 0.23	35.96 ± 8.06	5.32 ± 5.67	0.72 ± 0.08	9.01	−0.46
J125257.36+331555.81	0.367	11.36 ± 0.07	1.32 ± 0.16	21.81 ± 4.75	1.66 ± 1.07	0.27 ± 0.05	8.16	−1.03
J125317.57+310550.64	0.782	11.61 ± 0.06	2.39 ± 0.21	54.60 ± 5.54	23.04 ± 8.04	0.74 ± 0.07	9.02	−0.41
J125553.05+272405.23	0.316	11.48 ± 0.14	1.03 ± 0.14	28.67 ± 5.56	6.98 ± 6.59	0.40 ± 0.06	8.86	−1.45
J125703.80+250457.56	0.821	10.89 ± 0.21	2.63 ± 0.15	42.65 ± 4.85	6.94 ± 6.80	0.25 ± 0.05	8.19	−0.31
J125711.97+274216.45	0.793	10.55 ± 0.31	2.09 ± 0.14	40.90 ± 6.18	9.33 ± 8.17	0.44 ± 0.03	8.74	−0.88
J125757.23+322929.29	0.806	12.20 ± 0.16	2.60 ± 0.26	17.75 ± 2.03	1.00 ± 0.04	0.83 ± 0.03	8.74	−0.85
J130622.96+225752.95	0.758	10.65 ± 0.43	2.06 ± 0.31	31.47 ± 8.35	5.02 ± 5.48	0.36 ± 0.10	8.44	−1.15
J130947.00+081948.23	0.154	10.72 ± 0.31	1.00 ± 0.22	58.64 ± 8.67	29.72 ± 1.87	0.83 ± 0.01	8.59	−0.93
J131247.97+250756.71	0.425	10.99 ± 0.36	2.21 ± 0.21	40.35 ± 3.07	19.48 ± 9.57	0.31 ± 0.07	8.63	−1.09
J131312.13+284730.01	0.259	11.21 ± 0.18	0.93 ± 0.17	20.49 ± 3.52	1.54 ± 1.28	0.47 ± 0.09	8.21	−1.21
J131531.66+265414.73	0.620	10.24 ± 0.23	1.92 ± 0.21	37.98 ± 5.94	4.13 ± 5.32	0.33 ± 0.08	7.88	−0.65
J132919.84+250626.45	0.781	10.44 ± 0.19	2.22 ± 0.20	40.34 ± 7.64	4.14 ± 4.26	0.34 ± 0.08	8.81	−0.85
J133005.71+254243.77	0.598	10.85 ± 0.23	2.48 ± 0.34	54.85 ± 3.20	16.61 ± 9.25	0.73 ± 0.07	9.11	−1.34
J133442.29+320939.60	0.649	10.71 ± 0.10	2.39 ± 0.30	48.41 ± 6.56	27.62 ± 5.11	0.88 ± 0.03	8.52	−0.55
J134356.74+253847.69	0.086	10.24 ± 0.17	0.45 ± 0.08	59.64 ± 3.01	4.33 ± 4.46	0.69 ± 0.02	7.92	−1.09
J135632.80+210352.35	0.301	9.70 ± 0.32	1.28 ± 0.21	64.77 ± 2.82	22.78 ± 8.30	0.81 ± 0.02	8.82	−1.08
J140621.89+222346.54	0.098	9.96 ± 0.23	0.49 ± 0.10	30.74 ± 3.30	2.67 ± 1.10	0.70 ± 0.01	7.36	−0.32
J140655.66+015712.88	0.427	10.32 ± 0.12	2.10 ± 0.19	46.44 ± 5.30	26.06 ± 6.68	0.31 ± 0.07	7.96	−0.43
J140700.40+282714.65	0.077	9.53 ± 0.33	1.21 ± 0.16	56.24 ± 4.23	4.13 ± 3.16	0.63 ± 0.01	8.79	−1.63
J141637.45+003352.28	0.434	10.47 ± 0.15	2.32 ± 0.19	65.79 ± 0.52	25.26 ± 6.59	0.27 ± 0.02	8.67	−1.31
J141644.62+190541.92	0.365	9.85 ± 0.13	1.67 ± 0.11	48.70 ± 5.31	20.35 ± 7.44	0.23 ± 0.03	7.51	−0.16
J141700.83+445606.39	0.113	11.34 ± 0.09	0.73 ± 0.27	28.77 ± 5.10	2.00 ± 1.62	0.61 ± 0.01	7.93	−0.89
J142052.44+525622.42	0.677	10.69 ± 0.19	2.17 ± 0.29	49.66 ± 4.64	10.98 ± 8.75	0.69 ± 0.08	8.57	−0.63
J142648.78+005323.24	0.220	11.00 ± 0.11	1.40 ± 0.11	33.16 ± 3.41	4.11 ± 1.23	0.24 ± 0.07	8.40	−1.22
J142710.94+013023.22	0.704	10.80 ± 0.31	2.45 ± 0.27	44.87 ± 4.24	11.43 ± 8.79	0.40 ± 0.09	8.24	−0.37
J142753.79+345248.35	0.514	10.01 ± 0.47	1.58 ± 0.11	29.40 ± 4.75	3.24 ± 3.84	0.39 ± 0.08	8.00	−0.48
J142918.15+592106.65	0.739	11.08 ± 0.43	2.70 ± 0.14	17.78 ± 1.02	1.00 ± 0.00	0.11 ± 0.01	8.94	−1.50
J142943.07+474726.23	0.221	10.50 ± 0.58	1.27 ± 0.30	65.31 ± 3.38	23.42 ± 8.27	0.59 ± 0.01	7.95	−0.47
J143624.81−002905.36	0.325	9.98 ± 0.24	1.66 ± 0.31	43.24 ± 6.56	6.18 ± 6.58	0.74 ± 0.08	7.73	−0.30
J144231.82+014353.43	0.280	11.29 ± 0.20	0.99 ± 0.14	20.51 ± 3.16	1.33 ± 0.88	0.44 ± 0.08	8.62	−1.66
J145001.69+022006.75	0.520	10.45 ± 0.39	2.28 ± 0.14	36.81 ± 3.25	6.71 ± 5.15	0.15 ± 0.02	8.23	−0.63
J145108.76+270926.92	0.064	10.67 ± 0.18	0.63 ± 0.21	62.36 ± 1.33	13.99 ± 6.45	0.67 ± 0.01	7.29	−0.12
J145538.73+002238.06	0.434	11.43 ± 0.23	1.71 ± 0.21	26.64 ± 2.10	1.07 ± 0.36	0.10 ± 0.02	8.18	−0.91
J152114.26+222743.87	0.136	10.48 ± 0.32	1.02 ± 0.16	57.56 ± 2.47	6.54 ± 4.15	0.78 ± 0.01	7.90	−0.63
J154530.24+484608.98	0.399	10.84 ± 0.03	2.51 ± 0.08	64.33 ± 2.35	30.00 ± 0.16	0.55 ± 0.01	8.51	−0.26
J155444.58+082221.48	0.119	11.04 ± 0.10	0.22 ± 0.11	37.40 ± 4.42	16.60 ± 8.64	0.87 ± 0.01	7.73	−0.43
J161413.20+260416.21	0.131	10.05 ± 0.47	0.91 ± 0.26	34.42 ± 1.55	1.38 ± 0.86	0.54 ± 0.01	7.99	−0.62
J163352.34+402115.66	0.782	11.14 ± 0.54	2.14 ± 0.08	27.92 ± 1.51	3.49 ± 0.87	0.24 ± 0.03	8.99	−1.30
J163915.81+412833.70	0.690	11.12 ± 0.34	1.84 ± 0.05	17.17 ± 0.07	1.00 ± 0.03	0.30 ± 0.02	8.06	−0.80
J171033.22+584456.86	0.281	10.93 ± 0.39	1.53 ± 0.15	46.81 ± 2.70	4.89 ± 5.84	0.10 ± 0.02	7.79	−0.76
J171352.43+584201.25	0.521	10.94 ± 0.30	2.11 ± 0.20	53.23 ± 12.05	2.88 ± 2.00	0.46 ± 0.02	8.84	−0.92
J220759.39+001722.62	0.368	10.25 ± 0.18	1.40 ± 0.21	26.63 ± 4.29	4.26 ± 4.16	0.51 ± 0.05	7.75	−0.53

Table A1. Cont.

Name	z	$\log M_{\star}$ ($M_{\odot}$)	$\log \text{SFR}_{\text{FIR}}$ ($M_{\odot} \text{ yr}^{-1}$)	T_{dust} (K)	U_{min}	f_{AGN}	$\log M_{\text{BH}}$ ($M_{\odot}$)	$\log \lambda_{\text{Edd}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
J223607.68+134355.32	0.326	10.15 ± 0.36	1.64 ± 0.13	64.76 ± 1.19	28.53 ± 4.15	0.81 ± 0.01	8.57	−0.52
J233741.32+001743.77	0.762	10.21 ± 0.07	2.08 ± 0.05	32.85 ± 1.88	4.83 ± 0.72	0.26 ± 0.02	8.53	−1.21
J235156.13−010913.34	0.174	11.06 ± 0.39	1.46 ± 0.15	50.62 ± 5.96	7.90 ± 5.74	0.68 ± 0.01	8.81	−1.18
PG0003+158	0.450	11.25 ± 0.25	1.45 ± 0.23	51.92 ± 10.72	11.18 ± 8.94	1.00 ± 0.00	9.45	−0.56
PG0003+199	0.025	10.22 ± 0.20	$−0.09 \pm 0.24$	43.34 ± 5.65	28.06 ± 4.54	0.89 ± 0.00	7.52	−0.45
PG0007+106	0.089	11.19 ± 0.20	0.94 ± 0.17	24.38 ± 1.49	1.00 ± 0.07	0.70 ± 0.00	8.87	−1.18
PG0026+129	0.142	11.45 ± 0.16	0.49 ± 0.15	51.10 ± 5.96	16.84 ± 9.23	0.96 ± 0.01	8.12	−0.15
PG0043+039	0.384	10.68 ± 0.38	1.24 ± 0.04	50.24 ± 9.40	17.28 ± 9.36	1.00 ± 0.01	9.28	−0.87
PG0049+171	0.064	9.67 ± 0.19	$−0.37 \pm 0.29$	33.89 ± 6.23	5.60 ± 5.77	0.90 ± 0.01	8.45	−1.58
PG0050+124	0.061	10.31 ± 0.49	1.39 ± 0.28	34.23 ± 2.80	4.71 ± 0.95	0.53 ± 0.00	7.57	0.09
PG0052+251	0.155	11.61 ± 0.36	1.13 ± 0.17	27.81 ± 2.06	2.67 ± 0.98	0.78 ± 0.00	8.99	−1.09
PG0157+001	0.164	10.64 ± 0.12	2.53 ± 0.10	61.62 ± 2.43	29.82 ± 1.47	0.16 ± 0.00	8.31	−0.46
PG0804+761	0.100	10.65 ± 0.69	0.73 ± 0.27	56.08 ± 4.18	28.77 ± 3.81	0.97 ± 0.00	8.55	−0.62
PG0838+770	0.131	10.85 ± 0.30	1.06 ± 0.30	29.78 ± 2.96	3.22 ± 1.29	0.53 ± 0.01	8.29	−0.69
PG0844+349	0.064	10.34 ± 0.22	0.30 ± 0.23	24.91 ± 1.10	1.00 ± 0.04	0.84 ± 0.00	8.03	−0.67
PG0921+525	0.035	10.23 ± 0.09	$−0.32 \pm 0.15$	31.69 ± 2.40	4.73 ± 0.79	0.87 ± 0.00	7.45	−0.95
PG0923+201	0.190	11.05 ± 0.50	0.84 ± 0.16	55.80 ± 4.56	21.31 ± 8.80	0.95 ± 0.01	9.33	−1.42
PG0923+129	0.029	10.80 ± 0.08	0.56 ± 0.16	36.45 ± 3.15	4.94 ± 0.34	0.38 ± 0.00	7.52	−0.79
PG0934+013	0.050	10.15 ± 0.19	0.47 ± 0.16	39.13 ± 2.85	3.83 ± 1.04	0.30 ± 0.00	7.15	−0.40
PG0947+396	0.206	11.28 ± 0.29	1.23 ± 0.24	58.49 ± 6.77	19.71 ± 8.60	0.86 ± 0.01	8.81	−1.13
PG0953+414	0.239	10.35 ± 0.64	1.03 ± 0.13	40.22 ± 7.75	7.00 ± 6.54	0.94 ± 0.01	8.74	−0.49
PG1001+054	0.161	10.34 ± 0.49	0.84 ± 0.26	44.19 ± 10.84	14.43 ± 8.80	0.88 ± 0.01	7.87	−0.26
PG1004+130	0.240	10.60 ± 0.21	1.68 ± 0.18	64.44 ± 2.16	30.00 ± 0.02	0.74 ± 0.01	9.43	−1.02
PG1011−040	0.058	10.92 ± 0.23	0.50 ± 0.33	27.32 ± 2.74	3.59 ± 1.26	0.55 ± 0.00	7.43	−0.30
PG1012+008	0.185	11.15 ± 0.29	1.36 ± 0.17	41.43 ± 2.83	5.41 ± 2.37	0.73 ± 0.01	8.39	−0.51
PG1022+519	0.045	10.54 ± 0.19	0.33 ± 0.04	35.04 ± 1.31	5.16 ± 1.41	0.43 ± 0.00	7.25	−0.68
PG1048+342	0.167	11.29 ± 0.20	0.90 ± 0.17	30.95 ± 6.85	2.08 ± 1.72	0.75 ± 0.01	8.50	−0.92
PG1048−090	0.344	10.47 ± 0.70	1.04 ± 0.06	41.45 ± 10.56	14.47 ± 9.24	0.96 ± 0.05	9.37	−0.90
PG1049−005	0.357	10.83 ± 0.21	2.31 ± 0.15	61.63 ± 6.18	29.47 ± 2.61	0.64 ± 0.00	9.34	−0.84
PG1100+772	0.313	10.72 ± 0.44	1.65 ± 0.20	60.30 ± 5.17	24.21 ± 7.27	0.80 ± 0.01	9.44	−0.99
PG1103−006	0.425	10.24 ± 0.54	1.36 ± 0.25	54.52 ± 7.73	19.85 ± 9.13	0.93 ± 0.01	9.49	−0.95
PG1114+445	0.144	10.97 ± 0.11	0.79 ± 0.14	64.00 ± 2.48	26.85 ± 5.89	0.95 ± 0.00	8.72	−1.12
PG1115+407	0.154	10.33 ± 0.31	1.42 ± 0.04	34.67 ± 1.25	10.25 ± 6.17	0.45 ± 0.00	7.80	−0.31
PG1116+215	0.177	10.35 ± 0.70	1.16 ± 0.02	46.35 ± 5.05	24.49 ± 6.76	0.99 ± 0.00	8.69	−0.42
PG1119+120	0.049	10.62 ± 0.21	0.70 ± 0.31	49.80 ± 10.14	22.83 ± 7.04	0.50 ± 0.00	7.58	−0.58
PG1121+422	0.234	10.77 ± 0.28	0.72 ± 0.26	40.75 ± 11.68	12.26 ± 9.11	1.00 ± 0.00	8.17	−0.42
PG1126−041	0.060	10.90 ± 0.41	0.96 ± 0.22	41.61 ± 6.08	12.49 ± 6.37	0.60 ± 0.00	7.87	−0.61
PG1149−110	0.049	11.07 ± 0.18	0.43 ± 0.30	34.59 ± 3.90	3.06 ± 1.05	0.44 ± 0.01	8.04	−1.06
PG1151+117	0.176	10.89 ± 0.33	0.47 ± 0.03	47.17 ± 11.78	11.98 ± 8.99	0.96 ± 0.03	8.68	−1.05
PG1202+281	0.165	10.36 ± 0.26	1.33 ± 0.16	58.97 ± 8.51	26.02 ± 6.11	0.66 ± 0.01	8.74	−1.27
PG1211+143	0.085	9.94 ± 0.55	0.60 ± 0.14	58.37 ± 3.36	28.98 ± 3.48	0.95 ± 0.00	8.10	−0.16
PG1216+069	0.334	10.70 ± 0.40	0.95 ± 0.28	40.12 ± 12.53	9.79 ± 8.44	1.00 ± 0.00	9.36	−0.77
PG1226+023	0.158	11.83 ± 0.03	1.74 ± 0.02	57.58 ± 5.07	29.98 ± 0.53	1.00 ± 0.00	9.18	−0.29
PG1229+204	0.064	11.10 ± 0.16	0.67 ± 0.25	37.28 ± 3.41	4.19 ± 0.99	0.69 ± 0.00	8.26	−1.01
PG1244+026	0.048	9.53 ± 0.34	0.27 ± 0.29	51.83 ± 11.18	17.26 ± 8.64	0.52 ± 0.00	6.62	0.05
PG1259+593	0.472	11.45 ± 0.54	1.41 ± 0.16	39.74 ± 13.67	10.22 ± 8.54	1.00 ± 0.00	9.09	−0.31
PG1302−102	0.286	11.78 ± 0.39	1.74 ± 0.34	30.88 ± 0.14	1.00 ± 0.00	0.85 ± 0.01	9.05	−0.35
PG1307+085	0.155	10.55 ± 0.44	0.88 ± 0.29	40.43 ± 4.78	20.78 ± 8.52	0.88 ± 0.01	9.00	−1.12
PG1309+355	0.184	11.04 ± 0.14	1.22 ± 0.24	31.01 ± 3.23	2.18 ± 1.02	0.85 ± 0.01	8.48	−0.60
PG1310−108	0.035	9.72 ± 0.47	$−0.14 \pm 0.21$	31.80 ± 9.32	2.08 ± 1.12	0.84 ± 0.00	7.99	−1.39
PG1322+659	0.168	10.65 ± 0.34	1.15 ± 0.10	54.11 ± 4.78	23.16 ± 7.38	0.72 ± 0.01	8.42	−0.57
PG1341+258	0.087	10.80 ± 0.14	0.48 ± 0.16	53.87 ± 5.92	4.93 ± 5.30	0.75 ± 0.01	8.15	−0.94
PG1351+236	0.055	10.91 ± 0.05	0.73 ± 0.05	35.07 ± 2.41	5.10 ± 1.09	0.19 ± 0.01	8.67	−1.75
PG1351+640	0.087	9.94 ± 0.21	1.51 ± 0.25	63.74 ± 5.61	29.83 ± 1.49	0.52 ± 0.00	8.97	−1.26
PG1352+183	0.158	10.68 ± 0.36	0.37 ± 0.15	52.06 ± 8.70	14.90 ± 9.50	0.96 ± 0.03	8.56	−0.87
PG1354+213	0.300	9.96 ± 0.51	1.20 ± 0.25	52.48 ± 10.55	20.80 ± 8.81	0.85 ± 0.01	8.77	−0.92
PG1402+261	0.164	10.32 ± 0.07	1.55 ± 0.09	49.04 ± 3.97	29.76 ± 1.71	0.71 ± 0.00	8.08	−0.23
PG1404+226	0.098	10.39 ± 0.25	0.51 ± 0.14	27.96 ± 2.63	1.93 ± 1.09	0.74 ± 0.01	7.01	0.24

Table A1. Cont.

Name	z	$\log M_{\star}$ ($M_{\odot}$)	$\log \text{SFR}_{\text{FIR}}$ ($M_{\odot} \text{ yr}^{-1}$)	T_{dust} (K)	U_{min}	f_{AGN}	$\log M_{\text{BH}}$ ($M_{\odot}$)	$\log \lambda_{\text{Edd}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
PG1411+442	0.089	10.25 ± 0.52	0.57 ± 0.14	31.05 ± 1.80	4.12 ± 1.05	0.90 ± 0.00	8.20	−0.70
PG1415+451	0.114	11.48 ± 0.14	0.92 ± 0.33	33.72 ± 4.99	4.48 ± 1.96	0.66 ± 0.00	8.14	−0.71
PG1416-129	0.129	10.30 ± 0.48	0.70 ± 0.13	52.50 ± 6.96	21.43 ± 8.32	0.66 ± 0.01	9.19	−1.18
PG1425+267	0.366	10.87 ± 0.30	1.81 ± 0.09	64.89 ± 2.08	29.90 ± 1.13	0.72 ± 0.01	9.90	−1.27
PG1426+015	0.086	11.65 ± 0.20	1.17 ± 0.17	40.66 ± 1.69	5.03 ± 0.67	0.72 ± 0.00	9.15	−1.40
PG1427+480	0.221	10.39 ± 0.48	1.13 ± 0.34	54.60 ± 12.06	22.31 ± 8.62	0.61 ± 0.01	8.22	−0.59
PG1435-067	0.129	10.55 ± 0.46	0.31 ± 0.13	47.68 ± 6.88	2.13 ± 3.02	0.93 ± 0.01	8.50	−0.71
PG1440+356	0.077	11.31 ± 0.13	1.33 ± 0.08	43.37 ± 5.82	22.60 ± 6.20	0.44 ± 0.00	7.60	−0.18
PG1444+407	0.267	10.54 ± 0.28	1.96 ± 0.18	43.85 ± 12.16	12.45 ± 9.16	1.00 ± 0.00	8.44	−0.37
PG1448+273	0.065	10.81 ± 0.09	0.51 ± 0.28	47.17 ± 10.35	5.40 ± 4.66	0.64 ± 0.00	7.09	0.26
PG1501+106	0.036	10.50 ± 0.13	0.52 ± 0.23	64.71 ± 3.05	29.56 ± 2.32	0.52 ± 0.00	8.64	−1.48
PG1512+370	0.371	11.47 ± 0.23	1.47 ± 0.27	37.91 ± 8.96	3.21 ± 3.95	1.00 ± 0.06	9.53	−1.06
PG1519+226	0.137	10.68 ± 0.36	1.01 ± 0.22	50.59 ± 7.84	14.98 ± 7.36	0.83 ± 0.00	8.07	−0.49
PG1534+580	0.030	10.00 ± 0.15	0.09 ± 0.17	60.45 ± 5.01	17.60 ± 7.02	0.72 ± 0.00	8.30	−1.74
PG1535+547	0.038	10.62 ± 0.12	−0.31 ± 0.30	25.77 ± 1.96	1.22 ± 0.65	0.91 ± 0.00	7.30	−0.47
PG1543+489	0.400	10.84 ± 0.25	2.35 ± 0.18	57.39 ± 10.43	25.27 ± 7.03	0.54 ± 0.00	8.16	0.16
PG1545+210	0.266	10.45 ± 0.49	0.98 ± 0.03	50.82 ± 9.85	14.16 ± 9.40	0.95 ± 0.03	9.47	−1.17
PG1552+085	0.119	10.87 ± 0.22	0.24 ± 0.18	31.98 ± 7.65	5.81 ± 6.03	0.90 ± 0.01	7.67	−0.10
PG1612+261	0.131	10.42 ± 0.44	0.95 ± 0.31	40.16 ± 5.14	3.58 ± 2.30	0.50 ± 0.01	8.19	−0.60
PG1613+658	0.129	11.67 ± 0.37	1.76 ± 0.09	39.36 ± 3.63	15.08 ± 5.05	0.46 ± 0.00	9.32	−1.61
PG1617+175	0.114	9.70 ± 0.68	0.38 ± 0.15	51.27 ± 5.44	16.54 ± 9.24	0.95 ± 0.01	8.91	−1.20
PG1626+554	0.133	11.12 ± 0.16	0.16 ± 0.13	39.45 ± 11.53	12.01 ± 9.06	1.00 ± 0.00	8.63	−1.18
PG1700+518	0.282	11.37 ± 0.40	2.11 ± 0.26	41.60 ± 11.37	15.17 ± 9.65	0.66 ± 0.00	8.61	−0.02
PG1704+608	0.371	11.56 ± 0.17	2.17 ± 0.10	61.52 ± 2.25	30.00 ± 0.09	0.81 ± 0.00	9.55	−0.98
PG2112+059	0.466	10.88 ± 0.49	2.13 ± 0.28	47.45 ± 8.34	29.72 ± 1.87	0.90 ± 0.00	9.18	−0.12
PG2130+099	0.061	10.05 ± 0.43	1.01 ± 0.11	46.25 ± 2.22	5.53 ± 2.51	0.75 ± 0.00	8.04	−0.60
PG2209+184	0.070	11.35 ± 0.07	0.33 ± 0.08	24.54 ± 0.75	1.00 ± 0.01	0.67 ± 0.01	8.89	−1.55
PG2214+139	0.067	11.41 ± 0.09	0.20 ± 0.11	33.39 ± 4.33	3.20 ± 1.17	0.90 ± 0.00	8.68	−1.15
PG2233+134	0.325	10.01 ± 0.41	1.55 ± 0.21	60.58 ± 4.25	20.03 ± 8.38	0.85 ± 0.01	8.19	0.01
PG2251+113	0.323	11.87 ± 0.29	1.28 ± 0.10	45.53 ± 6.90	14.72 ± 8.90	0.90 ± 0.01	9.15	−0.59
PG2304+042	0.042	11.06 ± 0.07	−0.45 ± 0.18	57.51 ± 3.85	8.67 ± 7.92	0.81 ± 0.02	8.68	−1.74
PG2308+098	0.432	11.67 ± 0.35	1.29 ± 0.06	32.84 ± 9.93	6.52 ± 6.60	1.00 ± 0.00	9.76	−1.11

Note: Column (1): source name. Column (2): redshift. Column (3): stellar mass. Column (4): FIR SFR derived from the cold dust component integrated over 8–1000 μm . Column (5): dust temperature. Column (6): minimum radiation field intensity. Column (7): fractional AGN contribution to the total infrared luminosity, measured over 8–1000 μm ; when a parameter is returned by multiple SED-fitting templates, we report the average value. Columns (8) and (9) give the black hole mass and Eddington ratio, respectively. For the SDSS quasars, M_{BH} and L_{bol} are adopted from Shen et al. [27]; for the PG quasars, they are adopted from Shanguan et al. [32]. The Eddington ratios are calculated as $\lambda_{\text{Edd}} = L_{\text{bol}}/L_{\text{Edd}}$.

Appendix B. Additional Comparisons of Dust-Related SED-Fitting Parameters

In this appendix, we present additional comparisons of the dust-related parameters derived with and without the SCUBA-2 data. The comparisons include the parameters from the CIGALE cold dust templates and AGNfitter. Figures A1–A3 show the detailed comparisons. In all panels, the symbols and colors are the same as in Figure 4. Red and blue symbols denote sources detected in no more than two and in at least three of the six Herschel bands, respectively. Triangles and circles represent radio-loud and radio-quiet sources, respectively.

For the casey2012 template, excluding the SCUBA-2 data leads to slightly higher dust temperatures and emissivity indices, with mean offsets of 0.43 K and 0.02 for the full sample, respectively. The offsets are larger for sources with limited Herschel detections and for radio-loud quasars, with maximum differences reaching ~ 18 K in T_{dust} and ~ 0.7 in β . For the dale2014 template, the mean offset in α is negligible for the full sample, but some sources with sparse Herschel coverage show larger deviations. For the dl2014 and THEMIS templates, the mean offsets in γ are close to zero for the full sample, although individual radio-loud sources

can show noticeable differences. The effect on $\log U_{\min}$ is also small on average, but sources with fewer Herschel detections show larger scatter. The PAH- or HAC-related parameters are only weakly affected for the full sample, while radio-loud quasars exhibit larger object-to-object variations. For AGNfitter, excluding the SCUBA-2 data produces a small mean offset in T_{dust} for the full sample, but the offset becomes larger for sources with limited Herschel detections. The mean effect on fracPAH is small, although some radio-loud quasars show larger differences.




Figure A1. Comparison of the dust-related parameters derived with and without the SCUBA-2 data: (a) dust temperature, T_{dust} , from the casey2012 template; (b) dust emissivity index, β , from the casey2012 template; and (c) dust-heating intensity parameter, α , from the dale2014 template. The lower subpanel in each panel shows the difference between the two estimates. The symbols and colors are the same as in Figure 4.




Figure A2. Comparison of the dust-related parameters derived with and without the SCUBA-2 data: (a) γ from the dl2014 template; (b) $\log U_{\min}$ from the dl2014 template; (c) q_{PAH} from the dl2014 template; (d) γ from the THEMIS template; (e) $\log U_{\min}$ from the THEMIS template; and (f) q_{HAC} from the THEMIS template. The lower subpanel in each panel shows the difference between the two estimates. The symbols and colors are the same as in Figure 4.




Figure A3. Comparison of the AGNfitter dust-related parameters derived with and without the SCUBA-2 data: (a) dust temperature, T_{dust} ; and (b) the PAH fraction, fracPAH . The lower subpanel in each panel shows the difference between the two estimates. The symbols and colors are the same as in Figure 4.

Overall, these comparisons confirm that the SCUBA-2 data have little effect on the average dust parameter estimates for the full sample, but they can provide important constraints for individual sources with poor far-infrared coverage or radio-loud emission.

Appendix C. Kendall- τ Tests for Infrared-Based AGN Indicators

Table A2 lists the Kendall- τ test results for the relations between Δ_{MS} and the AGNfitter torus indicators. For L_{tor} , significant positive correlations are found in three redshift bins, while the $0.3 < z < 0.5$ bin shows no significant trend. In contrast, Δ_{MS} is positively correlated with $L_{\text{tor}}/L_{\text{Edd}}$ in all four redshift bins.

We also tested the residual redshift dependence of $\log(L_{\text{tor}}/L_{\text{Edd}})$ within each redshift bin. No significant correlation with redshift is found, with p -values of 0.25, 0.31, 0.24, and 0.18 from the lowest- to highest-redshift bins. Therefore, the $\Delta_{\text{MS}}-L_{\text{tor}}/L_{\text{Edd}}$ relation is not readily explained by a simple residual redshift trend within the bins.

Table A2. Kendall- τ test results for the relations between the main-sequence offset and AGNfitter torus indicators. The tests are based on individual sources within each redshift bin.

Redshift Bin	N	$\Delta_{\text{MS}}-L_{\text{tor}}$		$\Delta_{\text{MS}}-L_{\text{tor}}/L_{\text{Edd}}$	
		τ	p	τ	p
$z < 0.1$	38	0.45	8.3×10^{-5}	0.30	0.009
$0.1 < z < 0.3$	68	0.24	0.004	0.23	0.005
$0.3 < z < 0.5$	49	0.16	0.110	0.39	1.1×10^{-4}
$z > 0.5$	47	0.30	0.003	0.21	0.044

Appendix D. Additional Stellar Mass Tests

In Section 4.5, we discussed the possible effects of stellar mass on the relations involving star formation and dust heating. Here, we present two additional stellar mass binned tests. Figure A4a shows the relation between Δ_{MS} and Eddington ratio, and Figure A4b shows the relation between T_{dust} and U_{min} .




Figure A4. Additional stellar mass binned tests. (a) Relation between the main-sequence offset, Δ_{MS} , and Eddington ratio. (b) Relation between dust temperature, T_{dust} , and the minimum radiation field intensity, U_{min} . Blue, green, and red symbols represent sources with $\log(M_*/M_\odot) < 10.5$, $10.5 < \log(M_*/M_\odot) < 11.0$, and $\log(M_*/M_\odot) > 11.0$, respectively. Large squares and connecting lines show the median values in bins along the abscissa, and the error bars indicate the corresponding scatter.

Appendix E. Robustness Tests with SPIRE-Quality Subsamples

In this appendix, we compare the full SDSS FIR-constrained sample with two SPIRE-quality subsamples defined in Section 4.6. The left panels show the subsample with $S/N_{250} \geq 5$, while the right panels show the subsample with detections at $S/N \geq 3$ in at least two SPIRE bands. Gray open circles represent the full SDSS sample. Blue-filled circles and orange-filled squares represent the two SPIRE-quality subsamples, respectively.




Figure A5. Comparison of the FIR SFR–stellar mass relation between the full SDSS FIR-constrained sample and the two SPIRE-quality subsamples. The dashed curves and shaded regions show the adopted star-forming main sequence and its scatter. Gray open circles show the full SDSS sample. Blue-filled circles in the left panel represent sources with $S/N_{250} \geq 5$, while orange-filled squares in the right panel represent sources detected at $S/N \geq 3$ in at least two SPIRE bands.




Figure A6. Same as Figure A5 but for the relation between the main-sequence offset, Δ_{MS} , and black hole accretion rate.




Figure A7. Same as Figure A5 but for the relation between Δ_{MS} and Eddington ratio.




Figure A8. Same as Figure A5 but for the relation between the minimum radiation field intensity, U_{min} , and bolometric luminosity. The Kendall- τ coefficient and corresponding p value are calculated for the highlighted SPIRE-quality subsample in each panel.

Appendix F. WISE-to-Herschel Color Diagnostics

Figures A9–A12 show the full set of relations between $f_{\text{AGN}}(8\text{--}1000\ \mu\text{m})$ and the flux-density ratios between the four WISE bands and the six Herschel bands.




Figure A9. Same as Figure 12 but for WISE W1 and the six Herschel bands: (a) W1/70 μm , (b) W1/100 μm , (c) W1/160 μm , (d) W1/250 μm , (e) W1/350 μm , and (f) W1/500 μm . Red and black symbols denote sources detected in at least three and in no more than two Herschel bands, respectively. Filled and open symbols represent radio-quiet and radio-loud quasars, respectively. The black dashed line and blue shaded region show the Bayesian linear regression fit and its uncertainty for the radio-quiet sources detected in at least three Herschel bands. The fitted relation, intrinsic scatter ϵ , and correlation coefficient ρ are shown in each panel.




Figure A10. Cont.




Figure A10. Same as Figure 12 but for WISE W2 and the six Herschel bands: (a) W2/70 μm , (b) W2/100 μm , (c) W2/160 μm , (d) W2/250 μm , (e) W2/350 μm , and (f) W2/500 μm . Red and black symbols denote sources detected in at least three and in no more than two Herschel bands, respectively. Filled and open symbols represent radio-quiet and radio-loud quasars, respectively. The black dashed line and blue shaded region show the Bayesian linear regression fit and its uncertainty for the radio-quiet sources detected in at least three Herschel bands. The fitted relation, intrinsic scatter ϵ , and correlation coefficient ρ are shown in each panel.




Figure A11. Same as Figure 12 but for WISE W3 and the six Herschel bands: (a) W3/70 μm , (b) W3/100 μm , (c) W3/160 μm , (d) W3/250 μm , (e) W3/350 μm , and (f) W3/500 μm . Red and black symbols denote sources detected in at least three and in no more than two Herschel bands, respectively. Filled and open symbols represent radio-quiet and radio-loud quasars, respectively. The black dashed line and blue shaded region show the Bayesian linear regression fit and its uncertainty for the radio-quiet sources detected in at least three Herschel bands. The fitted relation, intrinsic scatter ϵ , and correlation coefficient ρ are shown in each panel.




Figure A12. Same as Figure 12 but for WISE W4 and the six Herschel bands: (a) W4/70 μm, (b) W4/100 μm, (c) W4/160 μm, (d) W4/250 μm, (e) W4/350 μm, and (f) W4/500 μm. Red and black symbols denote sources detected in at least three and in no more than two Herschel bands, respectively. Filled and open symbols represent radio-quiet and radio-loud quasars, respectively. The black dashed line and blue shaded region show the Bayesian linear regression fit and its uncertainty for the radio-quiet sources detected in at least three Herschel bands. The fitted relation, intrinsic scatter ϵ , and correlation coefficient ρ are shown in each panel.

Notes

- <https://irsa.ipac.caltech.edu/>, accessed on 5 May 2021
- <https://www.cadc-ccda.hia-ihp.nrc-cnrc.gc.ca/en/search/?collection=JCMT&Observation.instrument.name=SCUBA-2&Plane.calibrationLevel=2&noexec=true>, accessed on 20 June 2020
- <https://ned.ipac.caltech.edu/>, accessed on 16 June 2023
- <https://cigale.lam.fr/>, accessed on 16 April 2022
- <https://github.com/GabrielaCR/AGNfitter/>, accessed on 19 February 2024

References

- Gebhardt, K.; Bender, R.; Bower, G.; Dressler, A.; Faber, S.M.; Filippenko, A.V.; Green, R.; Grillmair, C.; Ho, L.C.; Kormendy, J.; et al. A Relationship between Nuclear Black Hole Mass and Galaxy Velocity Dispersion. *Astrophys. J. Lett.* **2000**, *539*, L13–L16. <https://doi.org/10.1086/312840>.
- Ferrarese, L.; Merritt, D. A Fundamental Relation between Supermassive Black Holes and Their Host Galaxies. *Astrophys. J. Lett.* **2000**, *539*, L9–L12. <https://doi.org/10.1086/312838>.
- Kormendy, J.; Ho, L.C. Coevolution (Or Not) of Supermassive Black Holes and Host Galaxies. *Annu. Rev. Astron. Astrophys.* **2013**, *51*, 511–653. <https://doi.org/10.1146/annurev-astro-082708-101811>.
- Shiple, H.V.; Papovich, C.; Rieke, G.H.; Brown, M.J.I.; Moustakas, J. A New Star Formation Rate Calibration from Polycyclic Aromatic Hydrocarbon Emission Features and Application to High-redshift Galaxies. *Astrophys. J.* **2016**, *818*, 60. <https://doi.org/10.3847/0004-637X/818/1/60>.
- Ho, L.C.; Keto, E. The Mid-Infrared Fine-Structure Lines of Neon as an Indicator of Star Formation Rate in Galaxies. *Astrophys. J.* **2007**, *658*, 314–318. <https://doi.org/10.1086/511260>.
- Shiple, H.V.; Papovich, C.; Rieke, G.H.; Dey, A.; Jannuzi, B.T.; Moustakas, J.; Weiner, B. Spitzer Spectroscopy of Infrared-luminous Galaxies: Diagnostics of Active Galactic Nuclei and Star Formation and Contribution to Total Infrared Luminosity. *Astrophys. J.* **2013**, *769*, 75. <https://doi.org/10.1088/0004-637X/769/1/75>.

7. Zhuang, M.Y.; Ho, L.C.; Shangguan, J. A New Method to Measure Star Formation Rates in Active Galaxies Using Mid-infrared Neon Emission Lines. *Astrophys. J.* **2019**, *873*, 103. <https://doi.org/10.3847/1538-4357/ab0650>.
8. Mineo, S.; Gilfanov, M.; Sunyaev, R. X-ray emission from star-forming galaxies—I. High-mass X-ray binaries. *Mon. Not. R. Astron. Soc.* **2012**, *419*, 2095–2115. <https://doi.org/10.1111/j.1365-2966.2011.19862.x>.
9. Brandt, W.N.; Alexander, D.M. Cosmic X-ray surveys of distant active galaxies. The demographics, physics, and ecology of growing supermassive black holes. *Astron. Astrophys. Rev.* **2015**, *23*, 1. <https://doi.org/10.1007/s00159-014-0081-z>.
10. Kennicutt, R.C., Jr. Star Formation in Galaxies Along the Hubble Sequence. *Annu. Rev. Astron. Astrophys.* **1998**, *36*, 189–232. <https://doi.org/10.1146/annurev.astro.36.1.189>.
11. Kennicutt, R.C.; Evans, N.J. Star Formation in the Milky Way and Nearby Galaxies. *Annu. Rev. Astron. Astrophys.* **2012**, *50*, 531–608. <https://doi.org/10.1146/annurev-astro-081811-125610>.
12. Mullaney, J.R.; Pannella, M.; Daddi, E.; Alexander, D.M.; Elbaz, D.; Hickox, R.C.; Bournaud, F.; Altieri, B.; Aussel, H.; Coia, D.; et al. GOODS-Herschel: The far-infrared view of star formation in active galactic nucleus host galaxies since $z \approx 3$. *Mon. Not. R. Astron. Soc.* **2012**, *419*, 95–115. <https://doi.org/10.1111/j.1365-2966.2011.19675.x>.
13. Rosario, D.J.; Santini, P.; Lutz, D.; Shao, L.; Maiolino, R.; Alexander, D.M.; Altieri, B.; Andreani, P.; Aussel, H.; Bauer, F.E.; et al. The mean star formation rate of X-ray selected active galaxies and its evolution from $z \sim 2.5$: Results from PEP-Herschel. *Astron. Astrophys.* **2012**, *545*, A45. <https://doi.org/10.1051/0004-6361/201219258>.
14. Farrah, D.; Ejercito, K.; Efstathiou, A.; Leisawitz, D.; Engholm, A.; Shivaee, I.; Bonato, M.; Clements, D.L.; Petty, S.; Pitchford, L.K.; et al. How Accurately Can Obscured Galaxy Luminosities Be Measured Using Spectral Energy Distribution Fitting of Near-through Far-infrared Observations? *Astrophys. J.* **2026**, *997*, 150. <https://doi.org/10.3847/1538-4357/ae21cd>.
15. Pacifici, C.; Iyer, K.G.; Mobasher, B.; da Cunha, E.; Acquaviva, V.; Burgarella, D.; Calistro Rivera, G.; Carnall, A.C.; Chang, Y.Y.; Chartab, N.; et al. The Art of Measuring Physical Parameters in Galaxies: A Critical Assessment of Spectral Energy Distribution Fitting Techniques. *Astrophys. J.* **2023**, *944*, 141. <https://doi.org/10.3847/1538-4357/acacff>.
16. Fritz, J.; Franceschini, A.; Hatziminaoglou, E. Revisiting the infrared spectra of active galactic nuclei with a new torus emission model. *Mon. Not. R. Astron. Soc.* **2006**, *366*, 767–786. <https://doi.org/10.1111/j.1365-2966.2006.09866.x>.
17. Siebenmorgen, R.; Heymann, F.; Efstathiou, A. Self-consistent two-phase AGN torus models*. SED library for observers. *Astron. Astrophys.* **2015**, *583*, A120. <https://doi.org/10.1051/0004-6361/201526034>.
18. Chen, C.T.J.; Hickox, R.C.; Alberts, S.; Brodwin, M.; Jones, C.; Murray, S.S.; Alexander, D.M.; Assef, R.J.; Brown, M.J.I.; Dey, A.; et al. A Correlation between Star Formation Rate and Average Black Hole Accretion in Star-forming Galaxies. *Astrophys. J.* **2013**, *773*, 3. <https://doi.org/10.1088/0004-637X/773/1/3>.
19. Hickox, R.C.; Mullaney, J.R.; Alexander, D.M.; Chen, C.T.J.; Civano, F.M.; Goulding, A.D.; Hainline, K.N. Black Hole Variability and the Star Formation-Active Galactic Nucleus Connection: Do All Star-forming Galaxies Host an Active Galactic Nucleus? *Astrophys. J.* **2014**, *782*, 9. <https://doi.org/10.1088/0004-637X/782/1/9>.
20. Stanley, F.; Harrison, C.M.; Alexander, D.M.; Swinbank, A.M.; Aird, J.A.; Del Moro, A.; Hickox, R.C.; Mullaney, J.R. A remarkably flat relationship between the average star formation rate and AGN luminosity for distant X-ray AGN. *Mon. Not. R. Astron. Soc.* **2015**, *453*, 591–604. <https://doi.org/10.1093/mnras/stv1678>.
21. Dai, Y.S.; Wilkes, B.J.; Bergeron, J.; Kuraszkiewicz, J.; Omont, A.; Atanas, A.; Teplitz, H.I. Is there a relationship between AGN and star formation in IR-bright AGNs? *Mon. Not. R. Astron. Soc.* **2018**, *478*, 4238–4254. <https://doi.org/10.1093/mnras/sty1341>.
22. Woo, J.H.; Son, D.; Rakshit, S. The Correlation of Outflow Kinematics with Star Formation Rate. VI. Gas Outflows in AGNs. *Astrophys. J.* **2020**, *901*, 66. <https://doi.org/10.3847/1538-4357/abad97>.
23. Zhuang, M.Y.; Ho, L.C. The Star-forming Main Sequence of the Host Galaxies of Low-redshift Quasars. *Astrophys. J.* **2022**, *934*, 130. <https://doi.org/10.3847/1538-4357/ac7aaf>.
24. Abazajian, K.N.; Adelman-McCarthy, J.K.; Agüeros, M.A.; Allam, S.S.; Allende Prieto, C.; An, D.; Anderson, K.S.J.; Anderson, S.F.; Annis, J.; Bahcall, N.A.; et al. The Seventh Data Release of the Sloan Digital Sky Survey. *Astrophys. J. Suppl. Ser.* **2009**, *182*, 543–558. <https://doi.org/10.1088/0067-0049/182/2/543>.
25. Schneider, D.P.; Richards, G.T.; Hall, P.B.; Strauss, M.A.; Anderson, S.F.; Boroson, T.A.; Ross, N.P.; Shen, Y.; Brandt, W.N.; Fan, X.; et al. The Sloan Digital Sky Survey Quasar Catalog. V. Seventh Data Release. *Astron. J.* **2010**, *139*, 2360. <https://doi.org/10.1088/0004-6256/139/6/2360>.
26. York, D.G.; Adelman, J.; Anderson, J.E., Jr.; Anderson, S.F.; Annis, J.; Bahcall, N.A.; Bakken, J.A.; Barkhouser, R.; Bastian, S.; Berman, E.; et al. The Sloan Digital Sky Survey: Technical Summary. *Astron. J.* **2000**, *120*, 1579–1587. <https://doi.org/10.1086/301513>.
27. Shen, Y.; Richards, G.T.; Strauss, M.A.; Hall, P.B.; Schneider, D.P.; Snedden, S.; Bizyaev, D.; Brewington, H.; Malanushenko, V.; Malanushenko, E.; et al. A Catalog of Quasar Properties from Sloan Digital Sky Survey Data Release 7. *Astrophys. J. Suppl. Ser.* **2011**, *194*, 45. <https://doi.org/10.1088/0067-0049/194/2/45>.

28. Pilbratt, G.L.; Riedinger, J.R.; Passvogel, T.; Crone, G.; Doyle, D.; Gageur, U.; Heras, A.M.; Jewell, C.; Metcalfe, L.; Ott, S.; et al. Herschel Space Observatory. An ESA facility for far-infrared and submillimetre astronomy. *Astron. Astrophys.* **2010**, *518*, L1. <https://doi.org/10.1051/0004-6361/201014759>.
29. Poglitsch, A.; Waelkens, C.; Geis, N.; Feuchtgruber, H.; Vandenbussche, B.; Rodriguez, L.; Krause, O.; Renotte, E.; van Hoof, C.; Saraceno, P.; et al. The Photodetector Array Camera and Spectrometer (PACS) on the Herschel Space Observatory. *Astron. Astrophys.* **2010**, *518*, L2. <https://doi.org/10.1051/0004-6361/201014535>.
30. Griffin, M.J.; Abergel, A.; Abreu, A.; Ade, P.A.R.; André, P.; Augueres, J.L.; Babbedge, T.; Bae, Y.; Baillie, T.; Baluteau, J.P.; et al. The Herschel-SPIRE instrument and its in-flight performance. *Astron. Astrophys.* **2010**, *518*, L3. <https://doi.org/10.1051/0004-6361/201014519>.
31. Holland, W.S.; Bintley, D.; Chapin, E.L.; Chrysostomou, A.; Davis, G.R.; Dempsey, J.T.; Duncan, W.D.; Fich, M.; Friberg, P.; Halpern, M.; et al. SCUBA-2: The 10 000 pixel bolometer camera on the James Clerk Maxwell Telescope. *Mon. Not. R. Astron. Soc.* **2013**, *430*, 2513–2533. <https://doi.org/10.1093/mnras/sts612>.
32. Shangguan, J.; Ho, L.C.; Xie, Y. On the Gas Content and Efficiency of AGN Feedback in Low-redshift Quasars. *Astrophys. J.* **2018**, *854*, 158. <https://doi.org/10.3847/1538-4357/aaa9be>.
33. Schmidt, M.; Green, R.F. Quasar evolution derived from the Palomar bright quasar survey and other complete quasar surveys. *Astrophys. J.* **1983**, *269*, 352–374. <https://doi.org/10.1086/161048>.
34. Boroson, T.A.; Green, R.F. The Emission-Line Properties of Low-Redshift Quasi-stellar Objects. *Astrophys. J. Suppl. Ser.* **1992**, *80*, 109. <https://doi.org/10.1086/191661>.
35. Jenness, T.; Economou, F. ORAC-DR: A generic data reduction pipeline infrastructure. *Astron. Comput.* **2015**, *9*, 40–48. <https://doi.org/10.1016/j.ascom.2014.10.005>.
36. Currie, M.J.; Berry, D.S.; Jenness, T.; Gibb, A.G.; Bell, G.S.; Draper, P.W. Starlink Software in 2013. In *Proceedings of the Astronomical Data Analysis Software and Systems XXIII*; Manset, N., Forshay, P., Eds.; Astronomical Society of the Pacific Conference Series; Astronomical Society of the Pacific: San Francisco, CA, USA, 2014; Volume 485, p. 391.
37. Skrutskie, M.F.; Cutri, R.M.; Stiening, R.; Weinberg, M.D.; Schneider, S.; Carpenter, J.M.; Beichman, C.; Capps, R.; Chester, T.; Elias, J.; et al. The Two Micron All Sky Survey (2MASS). *Astron. J.* **2006**, *131*, 1163–1183. <https://doi.org/10.1086/498708>.
38. Cohen, M.; Wheaton, W.A.; Megeath, S.T. Spectral Irradiance Calibration in the Infrared. XIV. The Absolute Calibration of 2MASS. *Astron. J.* **2003**, *126*, 1090–1096. <https://doi.org/10.1086/376474>.
39. Wright, E.L.; Eisenhardt, P.R.M.; Mainzer, A.K.; Ressler, M.E.; Cutri, R.M.; Jarrett, T.; Kirkpatrick, J.D.; Padgett, D.; McMillan, R.S.; Skrutskie, M.; et al. The Wide-field Infrared Survey Explorer (WISE): Mission Description and Initial On-orbit Performance. *Astron. J.* **2010**, *140*, 1868–1881. <https://doi.org/10.1088/0004-6256/140/6/1868>.
40. Fazio, G.G.; Hora, J.L.; Allen, L.E.; Ashby, M.L.N.; Barmby, P.; Deutsch, L.K.; Huang, J.S.; Kleiner, S.; Marengo, M.; Megeath, S.T.; et al. The Infrared Array Camera (IRAC) for the Spitzer Space Telescope. *Astrophys. J. Suppl. Ser.* **2004**, *154*, 10–17. <https://doi.org/10.1086/422843>.
41. Rieke, G.H.; Young, E.T.; Engelbracht, C.W.; Kelly, D.M.; Low, F.J.; Haller, E.E.; Beeman, J.W.; Gordon, K.D.; Stansberry, J.A.; Misselt, K.A.; et al. The Multiband Imaging Photometer for Spitzer (MIPS). *Astrophys. J. Suppl. Ser.* **2004**, *154*, 25–29. <https://doi.org/10.1086/422717>.
42. Becker, R.H.; White, R.L.; Helfand, D.J. The FIRST Survey: Faint Images of the Radio Sky at Twenty Centimeters. *Astrophys. J.* **1995**, *450*, 559. <https://doi.org/10.1086/176166>.
43. White, R.L.; Becker, R.H.; Helfand, D.J.; Gregg, M.D. A Catalog of 1.4 GHz Radio Sources from the FIRST Survey. *Astrophys. J.* **1997**, *475*, 479–493. <https://doi.org/10.1086/303564>.
44. Helfand, D.J.; White, R.L.; Becker, R.H. The Last of FIRST: The Final Catalog and Source Identifications. *Astrophys. J.* **2015**, *801*, 26. <https://doi.org/10.1088/0004-637X/801/1/26>.
45. Condon, J.J.; Cotton, W.D.; Greisen, E.W.; Yin, Q.F.; Perley, R.A.; Taylor, G.B.; Broderick, J.J. The NRAO VLA Sky Survey. *Astron. J.* **1998**, *115*, 1693–1716. <https://doi.org/10.1086/300337>.
46. Wall, J.V.; Shimmins, A.J.; Merkelijn, J.K. The Parkes 2700 MHz Survey. Catalogues for the ± 4 declination zone and for the selected regions. *Aust. J. Phys. Astrophys. Suppl.* **1971**, *19*, 1–68.
47. Griffith, M.R.; Wright, A.E.; Burke, B.F.; Ekers, R.D. The Parkes-MIT-NRAO (PMN) Surveys. VI. Source Catalog for the Equatorial Survey (-9.5 degrees $< \delta < +10.0$ degrees). *Astrophys. J. Suppl. Ser.* **1995**, *97*, 347. <https://doi.org/10.1086/192146>.
48. Owen, F.N.; Porcas, R.W.; Mufson, S.L.; Moffett, T.J. Observations of radio sources with flat spectra. *Astron. J.* **1978**, *83*, 685–696. <https://doi.org/10.1086/112253>.
49. Wright, A.; Otrupcek, R. Parkes Catalog, 1990, Australia telescope national facility. *PKS Cat.* **1990**.
50. Becker, R.H.; White, R.L.; Edwards, A.L. A New Catalog of 53,522 4.85 GHz Sources. *Astrophys. J. Suppl. Ser.* **1991**, *75*, 1. <https://doi.org/10.1086/191529>.
51. Hales, S.E.G.; Baldwin, J.E.; Warner, P.J. The 6C survey of radio sources—II. The zone $30^\circ < \delta < 51^\circ$, $08^h30^m < \alpha < 17^h30^m$. *Mon. Not. R. Astron. Soc.* **1988**, *234*, 919–936. <https://doi.org/10.1093/mnras/234.4.919>.

52. Kuehr, H.; Witzel, A.; Pauliny-Toth, I.I.K.; Nauber, U. A Catalogue of Extragalactic Radio Sources Having Flux Densities Greater than 1-JY at 5-GHZ. *Astron. Astrophys. Suppl. Ser.* **1981**, *45*, 367.
53. Gregory, P.C.; Condon, J.J. The 87GB Catalog of Radio Sources Covering 0 degrees < delta < +75 degrees at 4.85 GHz. *Astrophys. J. Suppl. Ser.* **1991**, *75*, 1011. <https://doi.org/10.1086/191559>.
54. Algaba, J.C.; Gabuzda, D.C.; Smith, P.S. Search for correlations between the optical and radio polarization of active galactic nuclei—I. VLBA polarization data at 15 + 22 + 43 GHz. *Mon. Not. R. Astron. Soc.* **2011**, *411*, 85–101. <https://doi.org/10.1111/j.1365-2966.2010.17654.x>.
55. Steppe, H.; Jeyakumar, S.; Saikia, D.J.; Salter, C.J. Millimeter-wavelength observations of compact steep-spectrum sources. *Astron. Astrophys. Suppl. Ser.* **1995**, *113*, 409.
56. Gear, W.K.; Stevens, J.A.; Hughes, D.H.; Litchfield, S.J.; Robson, E.I.; Terasanta, H.; Valtaoja, E.; Steppe, H.; Aller, M.F.; Aller, H.D. A comparison of the radio-submillimetre spectra of BL Lacertae objects and flat-spectrum radio quasars. *Mon. Not. R. Astron. Soc.* **1994**, *267*, 167–186. <https://doi.org/10.1093/mnras/267.1.167>.
57. Brotherton, M.S.; Singh, V.; Runnoe, J. Orientation and quasar black hole mass estimation. *Mon. Not. R. Astron. Soc.* **2015**, *454*, 3864–3871. <https://doi.org/10.1093/mnras/stv2186>.
58. Hardcastle, M.J.; Gürkan, G.; van Weeren, R.J.; Williams, W.L.; Best, P.N.; de Gasperin, F.; Rafferty, D.A.; Read, S.C.; Sabater, J.; Shimwell, T.W.; et al. LOFAR/H-ATLAS: A deep low-frequency survey of the Herschel-ATLAS North Galactic Pole field. *Mon. Not. R. Astron. Soc.* **2016**, *462*, 1910–1936. <https://doi.org/10.1093/mnras/stw1763>.
59. Richards, J.L.; Max-Moerbeck, W.; Pavlidou, V.; King, O.G.; Pearson, T.J.; Readhead, A.C.S.; Reeves, R.; Shepherd, M.C.; Stevenson, M.A.; Weintraub, L.C.; et al. Blazars in the Fermi Era: The OVRO 40 m Telescope Monitoring Program. *Astrophys. J. Suppl. Ser.* **2011**, *194*, 29. <https://doi.org/10.1088/0067-0049/194/2/29>.
60. Wright, E.L.; Chen, X.; Odegard, N.; Bennett, C.L.; Hill, R.S.; Hinshaw, G.; Jarosik, N.; Komatsu, E.; Nolte, M.R.; Page, L.; et al. Five-Year Wilkinson Microwave Anisotropy Probe Observations: Source Catalog. *Astrophys. J. Suppl. Ser.* **2009**, *180*, 283–295. <https://doi.org/10.1088/0067-0049/180/2/283>.
61. Vollmer, B.; Krichbaum, T.P.; Angelakis, E.; Kovalev, Y.Y. Quasi-simultaneous multi-frequency observations of inverted-spectrum GPS candidate sources. *Astron. Astrophys.* **2008**, *489*, 49–55. <https://doi.org/10.1051/0004-6361:20078857>.
62. Richards, J.L.; Hovatta, T.; Max-Moerbeck, W.; Pavlidou, V.; Pearson, T.J.; Readhead, A.C.S. Connecting radio variability to the characteristics of gamma-ray blazars. *Mon. Not. R. Astron. Soc.* **2014**, *438*, 3058–3069. <https://doi.org/10.1093/mnras/stt2412>.
63. Antón, S.; Browne, I.W.A.; Marchã, M.J.M.; Bondi, M.; Polatidis, A. The spectral energy distributions of the revised 200-mJy sample. *Mon. Not. R. Astron. Soc.* **2004**, *352*, 673–688. <https://doi.org/10.1111/j.1365-2966.2004.07961.x>.
64. White, R.L.; Becker, R.H. A New Catalog of 30,239 1.4 GHz Sources. *Astrophys. J. Suppl. Ser.* **1992**, *79*, 331. <https://doi.org/10.1086/191656>.
65. Pauliny-Toth, I.I.K.; Wade, C.M.; Heeschen, D.S. Positions and Flux Densities of Radio Sources. *Astrophys. J. Suppl. Ser.* **1966**, *13*, 65. <https://doi.org/10.1086/190137>.
66. Hales, S.E.G.; Mayer, C.J.; Warner, P.J.; Baldwin, J.E. The 6C survey of radio sources—IV. The zone 67 < delta < 82, 0 h < alpha < 24 h. *Mon. Not. R. Astron. Soc.* **1991**, *251*, 46. <https://doi.org/10.1093/mnras/251.1.46>.
67. Laing, R.A.; Peacock, J.A. The relation between radio luminosity and spectrum for extended extragalactic radio sources. *Mon. Not. R. Astron. Soc.* **1980**, *190*, 903–924. <https://doi.org/10.1093/mnras/190.4.903>.
68. Gilbert, G.M.; Riley, J.M.; Hardcastle, M.J.; Croston, J.H.; Pooley, G.G.; Alexander, P. High-resolution observations of a complete sample of 27 FR II radio galaxies and quasars with 0.3 < z < 0.6. *Mon. Not. R. Astron. Soc.* **2004**, *351*, 845–890. <https://doi.org/10.1111/j.1365-2966.2004.07824.x>.
69. Tingay, S.J.; Hancock, P.J.; Wayth, R.B.; Intema, H.; Jagannathan, P.; Mooley, K. A Multi-resolution, Multi-epoch Low Radio Frequency Survey of the Kepler K2 Mission Campaign 1 Field. *Astron. J.* **2016**, *152*, 82. <https://doi.org/10.3847/0004-6256/152/4/82>.
70. Sajina, A.; Partridge, B.; Evans, T.; Stefl, S.; Vechik, N.; Myers, S.; Dicker, S.; Korngut, P. High-frequency Radio Spectral Energy Distributions and Polarization Fractions of Sources in an Atacama Cosmology Telescope Survey Field. *Astrophys. J.* **2011**, *732*, 45. <https://doi.org/10.1088/0004-637X/732/1/45>.
71. Arshakian, T.G.; Torrealba, J.; Chavushyan, V.H.; Ros, E.; Lister, M.L.; Cruz-González, I.; Zensus, J.A. Radio-optical scrutiny of compact AGN: Correlations between properties of pc-scale jets and optical nuclear emission. *Astron. Astrophys.* **2010**, *520*, A62. <https://doi.org/10.1051/0004-6361/201014418>.
72. Bolton, J.G.; Savage, A.; Wright, A.E. The Parkes 2700 MHz survey (14th part): Catalogue and new optical identifications. *Aust. J. Phys. Astrophys. Suppl.* **1979**, *46*, 1–27.
73. Griffith, M.R.; Wright, A.E.; Burke, B.F.; Ekers, R.D. The Parkes-MIT-NRAO (PMN) Surveys. III. Source Catalog for the Tropical Survey (-29 degrees < delta < -9 degrees -3pt.5). *Astrophys. J. Suppl. Ser.* **1994**, *90*, 179. <https://doi.org/10.1086/191863>.
74. Haas, M.; Klaas, U.; Müller, S.A.H.; Bertoldi, F.; Camenzind, M.; Chini, R.; Krause, O.; Lemke, D.; Meisenheimer, K.; Richards, P.J.; et al. The ISO view of Palomar-Green quasars. *Astron. Astrophys.* **2003**, *402*, 87–111. <https://doi.org/10.1051/0004-6361:20030110>.

75. WalDRAM, E.M.; Yates, J.A.; Riley, J.M.; Warner, P.J. The 7C survey of radio sources at 151 MHz—A region covering RA 9^h to 16^h and Dec. 20° to 35°. *Mon. Not. R. Astron. Soc.* **1996**, *282*, 779–787. <https://doi.org/10.1093/mnras/282.3.779>.
76. Kuźmicz, A.; Jamroz, M.; Bronarska, K.; Janda-Boczar, K.; Saikia, D.J. An Updated Catalog of Giant Radio Sources. *Astrophys. J. Suppl. Ser.* **2018**, *238*, 9. <https://doi.org/10.3847/1538-4365/aad9ff>.
77. WalDRAM, E.M.; Bolton, R.C.; Pooley, G.G.; Riley, J.M. Some estimates of the source counts at Planck Surveyor frequencies, using the 9C survey at 15 GHz. *Mon. Not. R. Astron. Soc.* **2007**, *379*, 1442–1452. <https://doi.org/10.1111/j.1365-2966.2007.12025.x>.
78. van Bemmél, I.M.; Bertoldi, F. Millimeter observations of radio-loud active galaxies. *Astron. Astrophys.* **2001**, *368*, 414–419. <https://doi.org/10.1051/0004-6361:20010040>.
79. Kellermann, K.I.; Pauliny-Toth, I.I.K.; Williams, P.J.S. The Spectra of Radio Sources in the Revised 3c Catalogue. *Astrophys. J.* **1969**, *157*, 1. <https://doi.org/10.1086/150046>.
80. Genzel, R.; Pauliny-Toth, I.I.K.; Preuss, E.; Witzel, A. Measurements of the flux densities and spectra of discrete radio sources at centimeter wavelengths. V. The observations at 14.9 GHz (2 cm). *Astron. J.* **1976**, *81*, 1084–1088. <https://doi.org/10.1086/111989>.
81. Burgarella, D.; Buat, V.; Iglesias-Páramo, J. Star formation and dust attenuation properties in galaxies from a statistical ultraviolet-to-far-infrared analysis. *Mon. Not. R. Astron. Soc.* **2005**, *360*, 1413–1425. <https://doi.org/10.1111/j.1365-2966.2005.09131.x>.
82. Noll, S.; Burgarella, D.; Giovannoli, E.; Buat, V.; Marcellac, D.; Muñoz-Mateos, J.C. Analysis of galaxy spectral energy distributions from far-UV to far-IR with CIGALE: Studying a SINGS test sample. *Astron. Astrophys.* **2009**, *507*, 1793–1813. <https://doi.org/10.1051/0004-6361/200912497>.
83. Boquien, M.; Burgarella, D.; Roehlly, Y.; Buat, V.; Ciesla, L.; Corre, D.; Inoue, A.K.; Salas, H. CIGALE: A python Code Investigating GALaxy Emission. *Astron. Astrophys.* **2019**, *622*, A103. <https://doi.org/10.1051/0004-6361/201834156>.
84. Bruzual, G.; Charlot, S. Stellar population synthesis at the resolution of 2003. *Mon. Not. R. Astron. Soc.* **2003**, *344*, 1000–1028. <https://doi.org/10.1046/j.1365-8711.2003.06897.x>.
85. Chabrier, G. Galactic Stellar and Substellar Initial Mass Function. *Publ. Astron. Soc. Pac.* **2003**, *115*, 763–795. <https://doi.org/10.1086/376392>.
86. Kroupa, P. On the variation of the initial mass function. *Mon. Not. R. Astron. Soc.* **2001**, *322*, 231–246. <https://doi.org/10.1046/j.1365-8711.2001.04022.x>.
87. Madau, P.; Dickinson, M. Cosmic Star-Formation History. *Annu. Rev. Astron. Astrophys.* **2014**, *52*, 415–486. <https://doi.org/10.1146/annurev-astro-081811-125615>.
88. Calzetti, D.; Armus, L.; Bohlin, R.C.; Kinney, A.L.; Koornneef, J.; Storchi-Bergmann, T. The Dust Content and Opacity of Actively Star-forming Galaxies. *Astrophys. J.* **2000**, *533*, 682–695. <https://doi.org/10.1086/308692>.
89. Dale, D.A.; Helou, G.; Magdis, G.E.; Armus, L.; Díaz-Santos, T.; Shi, Y. A Two-parameter Model for the Infrared/Submillimeter/Radio Spectral Energy Distributions of Galaxies and Active Galactic Nuclei. *Astrophys. J.* **2014**, *784*, 83. <https://doi.org/10.1088/0004-637X/784/1/83>.
90. Draine, B.T.; Aniano, G.; Krause, O.; Groves, B.; Sandstrom, K.; Braun, R.; Leroy, A.; Klaas, U.; Linz, H.; Rix, H.W.; et al. Andromeda’s Dust. *Astrophys. J.* **2014**, *780*, 172. <https://doi.org/10.1088/0004-637X/780/2/172>.
91. Casey, C.M. Far-infrared spectral energy distribution fitting for galaxies near and far. *Mon. Not. R. Astron. Soc.* **2012**, *425*, 3094–3103. <https://doi.org/10.1111/j.1365-2966.2012.21455.x>.
92. Jones, A.P.; Köhler, M.; Ysard, N.; Bocchio, M.; Verstraete, L. The global dust modelling framework THEMIS. *Astron. Astrophys.* **2017**, *602*, A46. <https://doi.org/10.1051/0004-6361/201630225>.
93. Stalevski, M.; Fritz, J.; Baes, M.; Nakos, T.; Popović, L.Č. 3D radiative transfer modelling of the dusty tori around active galactic nuclei as a clumpy two-phase medium. *Mon. Not. R. Astron. Soc.* **2012**, *420*, 2756–2772. <https://doi.org/10.1111/j.1365-2966.2011.19775.x>.
94. Stalevski, M.; Ricci, C.; Ueda, Y.; Lira, P.; Fritz, J.; Baes, M. The dust covering factor in active galactic nuclei. *Mon. Not. R. Astron. Soc.* **2016**, *458*, 2288–2302. <https://doi.org/10.1093/mnras/stw444>.
95. Helou, G.; Soifer, B.T.; Rowan-Robinson, M. Thermal infrared and nonthermal radio : Remarkable correlation in disks of galaxies. *Astrophys. J. Lett.* **1985**, *298*, L7–L11. <https://doi.org/10.1086/184556>.
96. Calistro Rivera, G.; Lusso, E.; Hennawi, J.F.; Hogg, D.W. AGNfitter: A Bayesian MCMC Approach to Fitting Spectral Energy Distributions of AGNs. *Astrophys. J.* **2016**, *833*, 98. <https://doi.org/10.3847/1538-4357/833/1/98>.
97. Martínez-Ramírez, L.N.; Calistro Rivera, G.; Lusso, E.; Bauer, F.E.; Nardini, E.; Buchner, J.; Brown, M.J.I.; Pineda, J.C.B.; Temple, M.J.; Banerji, M.; et al. AGNFITTER-RX: Modeling the radio-to-X-ray spectral energy distributions of AGNs. *Astron. Astrophys.* **2024**, *688*, A46. <https://doi.org/10.1051/0004-6361/202449329>.
98. Schreiber, C.; Elbaz, D.; Pannella, M.; Ciesla, L.; Wang, T.; Franco, M. Dust temperature and mid-to-total infrared color distributions for star-forming galaxies at 0 < z < 4. *Astron. Astrophys.* **2018**, *609*, A30. <https://doi.org/10.1051/0004-6361/201731506>.
99. Temple, M.J.; Hewett, P.C.; Banerji, M. Modelling type 1 quasar colours in the era of Rubin and Euclid. *Mon. Not. R. Astron. Soc.* **2021**, *508*, 737–754. <https://doi.org/10.1093/mnras/stab2586>.

100. Hatziminaoglou, E.; Messias, H.; Souza, R.; Borkar, A.; Farrah, D.; Feltre, A.; Magdis, G.; Pitchford, L.K.; Pérez-Fournon, I. An ALMA Band 7 survey of SDSS/Herschel quasars in Stripe 82: I. The properties of the 870 micron counterparts. *Astron. Astrophys.* **2025**, *702*, A183. <https://doi.org/10.1051/0004-6361/202555105>.
101. Popesso, P.; Morselli, L.; Concas, A.; Schreiber, C.; Rodighiero, G.; Cresci, G.; Belli, S.; Ilbert, O.; Erfanianfar, G.; Mancini, C.; et al. The main sequence of star-forming galaxies - II. A non-evolving slope at the high-mass end. *Mon. Not. R. Astron. Soc.* **2019**, *490*, 5285–5299. <https://doi.org/10.1093/mnras/stz2635>.
102. Brinchmann, J.; Charlot, S.; White, S.D.M.; Tremonti, C.; Kauffmann, G.; Heckman, T.; Brinkmann, J. The physical properties of star-forming galaxies in the low-redshift Universe. *Mon. Not. R. Astron. Soc.* **2004**, *351*, 1151–1179. <https://doi.org/10.1111/j.1365-2966.2004.07881.x>.
103. Elbaz, D.; Dickinson, M.; Hwang, H.S.; Díaz-Santos, T.; Magdis, G.; Magnelli, B.; Le Borgne, D.; Galliano, F.; Pannella, M.; Chanial, P.; et al. GOODS-Herschel: An infrared main sequence for star-forming galaxies. *Astron. Astrophys.* **2011**, *533*, A119. <https://doi.org/10.1051/0004-6361/201117239>.
104. Speagle, J.S.; Steinhardt, C.L.; Capak, P.L.; Silverman, J.D. A Highly Consistent Framework for the Evolution of the Star-Forming “Main Sequence” from $z \sim 0$ –6. *Astrophys. J. Suppl. Ser.* **2014**, *214*, 15. <https://doi.org/10.1088/0067-0049/214/2/15>.
105. Renzini, A.; Peng, Y.J. An Objective Definition for the Main Sequence of Star-forming Galaxies. *Astrophys. J. Lett.* **2015**, *801*, L29. <https://doi.org/10.1088/2041-8205/801/2/L29>.
106. Popesso, P.; Concas, A.; Morselli, L.; Schreiber, C.; Rodighiero, G.; Cresci, G.; Belli, S.; Erfanianfar, G.; Mancini, C.; Inami, H.; et al. The main sequence of star-forming galaxies - I. The local relation and its bending. *Mon. Not. R. Astron. Soc.* **2019**, *483*, 3213–3226. <https://doi.org/10.1093/mnras/sty3210>.
107. Shimizu, T.T.; Mushotzky, R.F.; Meléndez, M.; Koss, M.; Rosario, D.J. Decreased specific star formation rates in AGN host galaxies. *Mon. Not. R. Astron. Soc.* **2015**, *452*, 1841–1860. <https://doi.org/10.1093/mnras/stv1407>.
108. Stemo, A.; Comerford, J.M.; Barrows, R.S.; Stern, D.; Assef, R.J.; Griffith, R.L. A Catalog of AGN Host Galaxies Observed with HST/ACS: Correlations between Star Formation and AGN Activity. *Astrophys. J.* **2020**, *888*, 78. <https://doi.org/10.3847/1538-4357/ab5f66>.
109. Husemann, B.; Jahnke, K.; Sánchez, S.F.; Wisotzki, L.; Nugroho, D.; Kupko, D.; Schramm, M. Integral field spectroscopy of nearby QSOs - I. ENLR size-luminosity relation, ongoing star formation and resolved gas-phase metallicities. *Mon. Not. R. Astron. Soc.* **2014**, *443*, 755–783. <https://doi.org/10.1093/mnras/stu1167>.
110. Woo, J.H.; Son, D.; Bae, H.J. Delayed or No Feedback? Gas Outflows in Type 2 AGNs. III. *Astrophys. J.* **2017**, *839*, 120. <https://doi.org/10.3847/1538-4357/aa6894>.
111. Smirnova-Pinchukova, I.; Husemann, B.; Davis, T.A.; Smith, C.M.A.; Singha, M.; Tremblay, G.R.; Klessen, R.S.; Powell, M.; Connor, T.; Baum, S.A.; et al. The Close AGN Reference Survey (CARS). No obvious signature of AGN feedback on star formation, but subtle trends. *Astron. Astrophys.* **2022**, *659*, A125. <https://doi.org/10.1051/0004-6361/202142011>.
112. Jarvis, M.E.; Harrison, C.M.; Mainieri, V.; Calistro Rivera, G.; Jethwa, P.; Zhang, Z.Y.; Alexander, D.M.; Circosta, C.; Costa, T.; De Breuck, C.; et al. High molecular gas content and star formation rates in local galaxies that host quasars, outflows, and jets. *Mon. Not. R. Astron. Soc.* **2020**, *498*, 1560–1575. <https://doi.org/10.1093/mnras/staa2196>.
113. Shangguan, J.; Ho, L.C.; Bauer, F.E.; Wang, R.; Treister, E. AGN Feedback and Star Formation of Quasar Host Galaxies: Insights from the Molecular Gas. *Astrophys. J.* **2020**, *899*, 112. <https://doi.org/10.3847/1538-4357/aba8a1>.
114. Xie, Y.; Ho, L.C.; Zhuang, M.Y.; Shangguan, J. The Infrared Emission and Vigorous Star Formation of Low-redshift Quasars. *Astrophys. J.* **2021**, *910*, 124. <https://doi.org/10.3847/1538-4357/abe404>.
115. Koutoulidis, L.; Mountrichas, G.; Georgantopoulos, I.; Poulisis, E.; Plionis, M. Host galaxy properties of X-ray active galactic nuclei in the local Universe. *Astron. Astrophys.* **2022**, *658*, A35. <https://doi.org/10.1051/0004-6361/202142466>.
116. Pitchford, L.K.; Hatziminaoglou, E.; Feltre, A.; Farrah, D.; Clarke, C.; Harris, K.A.; Hurley, P.; Oliver, S.; Page, M.; Wang, L. Extreme star formation events in quasar hosts over $0.5 < z < 4$. *Mon. Not. R. Astron. Soc.* **2016**, *462*, 4067–4077. <https://doi.org/10.1093/mnras/stw1840>.
117. Schneider, R.; Bianchi, S.; Valiante, R.; Risaliti, G.; Salvadori, S. The origin of the far-infrared continuum of $z \sim 6$ quasars. A radiative transfer model for SDSS J1148+5251. *Astron. Astrophys.* **2015**, *579*, A60. <https://doi.org/10.1051/0004-6361/201526105>.
118. McKinney, J.; Hayward, C.C.; Rosenthal, L.J.; Martínez-Galarza, J.R.; Pope, A.; Sajina, A.; Smith, H.A. Dust-enshrouded AGNs Can Dominate Host-galaxy-scale Cold Dust Emission. *Astrophys. J.* **2021**, *921*, 55. <https://doi.org/10.3847/1538-4357/ac185f>.
119. Di Mascia, F.; Carniani, S.; Gallerani, S.; Vito, F.; Pallottini, A.; Ferrara, A.; Valentini, M. Is the star-formation rate in $z \sim 6$ quasars overestimated? *Mon. Not. R. Astron. Soc.* **2023**, *518*, 3667–3674. <https://doi.org/10.1093/mnras/stac3306>.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.