



Rob J Hyndman

Functional time series

with applications in demography

5. Forecasting functional time series via PLS

Outline

- 1 Functional Partial Least Squares**
- 2 Application: French mortality rates
- 3 Application: Australian fertility rates
- 4 Forecast accuracy comparisons
- 5 Bootstrap intervals
- 6 Comparisons
- 7 References

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using
- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using

- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using

$$\hat{s}_{t+1|T}(x) = \hat{\mu}(x) + \int [s_T(x) - \hat{\mu}(x)] b(x, u) du$$

- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using

$$\hat{s}_{T+1|T}(x) = \hat{\mu}(x) + \int [s_T(x) - \hat{\mu}(x)] b(x, u) du$$

- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using

$$\hat{s}_{T+1|T}(x) = \hat{\mu}(x) + \int [s_T(x) - \hat{\mu}(x)] b(x, u) du$$

- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using

$$\hat{s}_{T+1|T}(x) = \hat{\mu}(x) + \int [s_T(x) - \hat{\mu}(x)] b(x, u) du$$

- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional PLS

- PCA components are designed to explain historical variation. They do not necessarily provide the best predictors.
- Partial least squares extracts uncorrelated latent components cores by maximizing the covariance between predictors and response.
- Response is $s_t(x)$ and predictor is $s_{t-1}(x)$.
- We want to predict $s_t(x)$ using

$$\hat{s}_{T+1|T}(x) = \hat{\mu}(x) + \int [s_T(x) - \hat{\mu}(x)] b(x, u) du$$

- This is a functional ARH(1).
- How to choose $b(x, u)$?

Functional weighted PLS

Outer relationship

$$\hat{\mathbf{f}}^*(x) = \sum_{k=1}^{\infty} \beta_k \psi_k(x), \quad \text{and} \quad \hat{\mathbf{g}}^*(x) = \sum_{k=1}^{\infty} \beta_k \phi_k(x)$$

- $\mathbf{W} = \text{diagonal}(w_1, \dots, w_T)$, $w_t = \kappa(1 - \kappa)^{T-t}$
- $\mathbf{f}^*(x) = \mathbf{W}[s_1^*(x), \dots, s_{T-1}^*(x)]'$ and $\mathbf{g}^*(x) = \mathbf{W}[s_2^*(x), \dots, s_T^*(x)]'$ are weighted decentralized functional predictors and responses
- β_k denotes common latent component scores
- $\psi_k(x)$ and $\phi_k(x)$ are latent components of predictors and responses respectively

Functional weighted PLS

Outer relationship

$$\hat{\mathbf{f}}^*(x) = \sum_{k=1}^{\infty} \beta_k \psi_k(x), \quad \text{and} \quad \hat{\mathbf{g}}^*(x) = \sum_{k=1}^{\infty} \beta_k \phi_k(x)$$

- $\mathbf{W} = \text{diagonal}(w_1, \dots, w_T)$, $w_t = \kappa(1 - \kappa)^{T-t}$
- $\mathbf{f}^*(x) = \mathbf{W}[s_1^*(x), \dots, s_{T-1}^*(x)]'$ and $\mathbf{g}^*(x) = \mathbf{W}[s_2^*(x), \dots, s_T^*(x)]'$ are weighted decentralized functional predictors and responses
- β_k denotes common latent component scores
- $\psi_k(x)$ and $\phi_k(x)$ are latent components of predictors and responses respectively

Functional weighted PLS

Outer relationship

$$\hat{\mathbf{f}}^*(x) = \sum_{k=1}^{\infty} \beta_k \psi_k(x), \quad \text{and} \quad \hat{\mathbf{g}}^*(x) = \sum_{k=1}^{\infty} \beta_k \phi_k(x)$$

- $\mathbf{W} = \text{diagonal}(w_1, \dots, w_T)$, $w_t = \kappa(1 - \kappa)^{T-t}$
- $\mathbf{f}^*(x) = \mathbf{W}[s_1^*(x), \dots, s_{T-1}^*(x)]'$ and $\mathbf{g}^*(x) = \mathbf{W}[s_2^*(x), \dots, s_T^*(x)]'$ are weighted decentralized functional predictors and responses
- β_k denotes common latent component scores
- $\psi_k(x)$ and $\phi_k(x)$ are latent components of predictors and responses respectively

Functional weighted PLS

Outer relationship

$$\hat{\mathbf{f}}^*(x) = \sum_{k=1}^{\infty} \beta_k \psi_k(x), \quad \text{and} \quad \hat{\mathbf{g}}^*(x) = \sum_{k=1}^{\infty} \beta_k \phi_k(x)$$

- $\mathbf{W} = \text{diagonal}(w_1, \dots, w_T)$, $w_t = \kappa(1 - \kappa)^{T-t}$
- $\mathbf{f}^*(x) = \mathbf{W}[s_1^*(x), \dots, s_{T-1}^*(x)]'$ and $\mathbf{g}^*(x) = \mathbf{W}[s_2^*(x), \dots, s_T^*(x)]'$ are weighted decentralized functional predictors and responses
- β_k denotes common latent component scores
- $\psi_k(x)$ and $\phi_k(x)$ are latent components of predictors and responses respectively

Functional weighted PLS

$$\beta_k = \int \mathbf{f}^*(x) w_k(x) dx = \int \mathbf{g}^*(x) m_k(x) dx,$$

Compute β_k iteratively:

1 Let $\mathbf{f}_0^*(x) = \mathbf{f}^*(x)$ and $\mathbf{g}_0^*(x) = \mathbf{g}^*(x)$

2 Obtain $w_k(x)$ iteratively, starting with $w_k^{(0)}(x) = 1$:

$$w_k^{(i)}(x) = \iint w_k^{(i-1)}(v) [\hat{\mathbf{f}}_{k-1}^*(v)]' \hat{\mathbf{g}}_{k-1}^*(u) [\hat{\mathbf{g}}_{k-1}^*(u)]' \hat{\mathbf{f}}_{k-1}^*(x) dv du$$

3 $\beta_k = \int \mathbf{f}^*(x) w_k(x) dx.$

4 $\hat{\mathbf{f}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{f}}_{k-1}^*(x)$

5 $\hat{\mathbf{g}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{g}}_{k-1}^*(x)$

Functional weighted PLS

$$\beta_k = \int \mathbf{f}^*(x) w_k(x) dx = \int \mathbf{g}^*(x) m_k(x) dx,$$

Compute β_k iteratively:

1 Let $\mathbf{f}_0^*(x) = \mathbf{f}^*(x)$ and $\mathbf{g}_0^*(x) = \mathbf{g}^*(x)$

2 Obtain $w_k(x)$ iteratively, starting with $w_k^{(0)}(x) = 1$:

$$w_k^{(i)}(x) = \iint w_k^{(i-1)}(v) [\hat{\mathbf{f}}_{k-1}^*(v)]' \hat{\mathbf{g}}_{k-1}^*(u) [\hat{\mathbf{g}}_{k-1}^*(u)]' \hat{\mathbf{f}}_{k-1}^*(x) dv du$$

3 $\beta_k = \int \mathbf{f}^*(x) w_k(x) dx.$

4 $\hat{\mathbf{f}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{f}}_{k-1}^*(x)$

5 $\hat{\mathbf{g}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{g}}_{k-1}^*(x)$

Functional weighted PLS

$$\beta_k = \int \mathbf{f}^*(x) w_k(x) dx = \int \mathbf{g}^*(x) m_k(x) dx,$$

Compute β_k iteratively:

1 Let $\mathbf{f}_0^*(x) = \mathbf{f}^*(x)$ and $\mathbf{g}_0^*(x) = \mathbf{g}^*(x)$

2 Obtain $w_k(x)$ iteratively, starting with $w_k^{(0)}(x) = 1$:

$$w_k^{(i)}(x) = \iint w_k^{(i-1)}(v) [\hat{\mathbf{f}}_{k-1}^*(v)]' \hat{\mathbf{g}}_{k-1}^*(u) [\hat{\mathbf{g}}_{k-1}^*(u)]' \hat{\mathbf{f}}_{k-1}^*(x) dv du$$

3 $\beta_k = \int \mathbf{f}^*(x) w_k(x) dx.$

4 $\hat{\mathbf{f}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{f}}_{k-1}^*(x)$

5 $\hat{\mathbf{g}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{g}}_{k-1}^*(x)$

Functional weighted PLS

$$\beta_k = \int \mathbf{f}^*(x) w_k(x) dx = \int \mathbf{g}^*(x) m_k(x) dx,$$

Compute β_k iteratively:

- 1 Let $\mathbf{f}_0^*(x) = \mathbf{f}^*(x)$ and $\mathbf{g}_0^*(x) = \mathbf{g}^*(x)$
- 2 Obtain $w_k(x)$ iteratively, starting with $w_k^{(0)}(x) = 1$:
$$w_k^{(i)}(x) = \iint w_k^{(i-1)}(v) [\hat{\mathbf{f}}_{k-1}^*(v)]' \hat{\mathbf{g}}_{k-1}^*(u) [\hat{\mathbf{g}}_{k-1}^*(u)]' \hat{\mathbf{f}}_{k-1}^*(x) dv du$$
- 3 $\beta_k = \int \mathbf{f}^*(x) w_k(x) dx.$
- 4 $\hat{\mathbf{f}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{f}}_{k-1}^*(x)$
- 5 $\hat{\mathbf{g}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{g}}_{k-1}^*(x)$

Functional weighted PLS

$$\beta_k = \int \mathbf{f}^*(x) w_k(x) dx = \int \mathbf{g}^*(x) m_k(x) dx,$$

Compute β_k iteratively:

- 1 Let $\mathbf{f}_0^*(x) = \mathbf{f}^*(x)$ and $\mathbf{g}_0^*(x) = \mathbf{g}^*(x)$
- 2 Obtain $w_k(x)$ iteratively, starting with $w_k^{(0)}(x) = 1$:
$$w_k^{(i)}(x) = \iint w_k^{(i-1)}(v) [\hat{\mathbf{f}}_{k-1}^*(v)]' \hat{\mathbf{g}}_{k-1}^*(u) [\hat{\mathbf{g}}_{k-1}^*(u)]' \hat{\mathbf{f}}_{k-1}^*(x) dv du$$
- 3 $\beta_k = \int \mathbf{f}^*(x) w_k(x) dx.$
- 4 $\hat{\mathbf{f}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{f}}_{k-1}^*(x)$
- 5 $\hat{\mathbf{g}}_k^*(x) = (\mathbf{I} - \beta_k \beta_k') \hat{\mathbf{g}}_{k-1}^*(x)$

Functional weighted PLS

Computationally equivalent approach

- Discretize $s_t^*(x)$ on a dense grid of q equally spaced points.
- Denote discretized $s_t^*(x)$ as $T \times q$ matrix \mathbf{G}^* and let $\mathbf{G} = \mathbf{W}\mathbf{G}^*$.
- Define \mathbf{G}_{k-1} and \mathbf{F}_{k-1} analogously
- $w_1(x) =$ largest eigenvector of $\mathbf{G}'\mathbf{F}$
- $m_k(x) =$ largest eigenvector of $\mathbf{G}'_{k-1}\mathbf{F}_{k-1}$
- $w_k(x) =$ largest eigenvector of $\mathbf{F}'_{k-1}\mathbf{G}_{k-1}$

Functional weighted PLS

Computationally equivalent approach

- Discretize $s_t^*(x)$ on a dense grid of q equally spaced points.
- Denote discretized $s_t^*(x)$ as $T \times q$ matrix \mathbf{G}^* and let $\mathbf{G} = \mathbf{W}\mathbf{G}^*$.
- Define \mathbf{G}_{k-1} and \mathbf{F}_{k-1} analogously
- $w_1(x) =$ largest eigenvector of $\mathbf{G}'\mathbf{F}$
- $m_k(x) =$ largest eigenvector of $\mathbf{G}'_{k-1}\mathbf{F}_{k-1}$
- $w_k(x) =$ largest eigenvector of $\mathbf{F}'_{k-1}\mathbf{G}_{k-1}$

Functional weighted PLS

Computationally equivalent approach

- Discretize $s_t^*(x)$ on a dense grid of q equally spaced points.
- Denote discretized $s_t^*(x)$ as $T \times q$ matrix \mathbf{G}^* and let $\mathbf{G} = \mathbf{W}\mathbf{G}^*$.
- Define \mathbf{G}_{k-1} and \mathbf{F}_{k-1} analogously
 - $w_1(x) =$ largest eigenvector of $\mathbf{G}'\mathbf{F}$
 - $m_k(x) =$ largest eigenvector of $\mathbf{G}'_{k-1}\mathbf{F}_{k-1}$
 - $w_k(x) =$ largest eigenvector of $\mathbf{F}'_{k-1}\mathbf{G}_{k-1}$

Functional weighted PLS

Computationally equivalent approach

- Discretize $s_t^*(x)$ on a dense grid of q equally spaced points.
- Denote discretized $s_t^*(x)$ as $T \times q$ matrix \mathbf{G}^* and let $\mathbf{G} = \mathbf{W}\mathbf{G}^*$.
- Define \mathbf{G}_{k-1} and \mathbf{F}_{k-1} analogously
- $w_1(x) =$ largest eigenvector of $\mathbf{G}'\mathbf{F}$
- $m_k(x) =$ largest eigenvector of $\mathbf{G}'_{k-1}\mathbf{F}_{k-1}$
- $w_k(x) =$ largest eigenvector of $\mathbf{F}'_{k-1}\mathbf{G}_{k-1}$

Functional weighted PLS

Computationally equivalent approach

- Discretize $s_t^*(x)$ on a dense grid of q equally spaced points.
- Denote discretized $s_t^*(x)$ as $T \times q$ matrix \mathbf{G}^* and let $\mathbf{G} = \mathbf{W}\mathbf{G}^*$.
- Define \mathbf{G}_{k-1} and \mathbf{F}_{k-1} analogously
- $w_1(x) =$ largest eigenvector of $\mathbf{G}'\mathbf{F}$
- $m_k(x) =$ largest eigenvector of $\mathbf{G}'_{k-1}\mathbf{F}_{k-1}$
- $w_k(x) =$ largest eigenvector of $\mathbf{F}'_{k-1}\mathbf{G}_{k-1}$

Functional weighted PLS

Computationally equivalent approach

- Discretize $s_t^*(x)$ on a dense grid of q equally spaced points.
- Denote discretized $s_t^*(x)$ as $T \times q$ matrix \mathbf{G}^* and let $\mathbf{G} = \mathbf{W}\mathbf{G}^*$.
- Define \mathbf{G}_{k-1} and \mathbf{F}_{k-1} analogously
- $w_1(x) =$ largest eigenvector of $\mathbf{G}'\mathbf{F}$
- $m_k(x) =$ largest eigenvector of $\mathbf{G}'_{k-1}\mathbf{F}_{k-1}$
- $w_k(x) =$ largest eigenvector of $\mathbf{F}'_{k-1}\mathbf{G}_{k-1}$

Functional weighted PLS

Functional autoregression coefficient:

$$b(x, u) = \sum_{k=1}^{\infty} w_k(u) \phi_k(x)$$

By orthogonality of β_k :

$$\phi_k(x) = (\beta_k' \beta_k)^{-1} \beta_k' \hat{\mathbf{g}}^*(x).$$

Therefore

$$\hat{b}(x, u) = \sum_{k=1}^K w_k(u) (\beta_k' \beta_k)^{-1} \beta_k' \hat{\mathbf{g}}^*(x),$$

for some finite K .

Functional weighted PLS

Functional autoregression coefficient:

$$b(x, u) = \sum_{k=1}^{\infty} w_k(u) \phi_k(x)$$

By orthogonality of β_k :

$$\phi_k(x) = (\beta_k' \beta_k)^{-1} \beta_k' \hat{\mathbf{g}}^*(x).$$

Therefore

$$\hat{b}(x, u) = \sum_{k=1}^K w_k(u) (\beta_k' \beta_k)^{-1} \beta_k' \hat{\mathbf{g}}^*(x),$$

for some finite K .

Functional weighted PLS

Functional autoregression coefficient:

$$b(x, u) = \sum_{k=1}^{\infty} w_k(u) \phi_k(x)$$

By orthogonality of β_k :

$$\phi_k(x) = (\beta_k' \beta_k)^{-1} \beta_k' \hat{\mathbf{g}}^*(x).$$

Therefore

$$\hat{b}(x, u) = \sum_{k=1}^K w_k(u) (\beta_k' \beta_k)^{-1} \beta_k' \hat{\mathbf{g}}^*(x),$$

for some finite K .

Functional weighted PLS

Forecasted curves:


$$\hat{f}_{t+1|t}(x) = \hat{\mu}(x) + \int [\hat{f}_t(u) - \hat{\mu}(u)] \hat{b}(x, u) du.$$

For $\hat{f}_{t+h|t}(x)$ where $h > 1$, apply iteratively.

Outline


- 1 Functional Partial Least Squares
- 2 Application: French mortality rates**
- 3 Application: Australian fertility rates
- 4 Forecast accuracy comparisons
- 5 Bootstrap intervals
- 6 Comparisons
- 7 References

French female mortality rates




French female mortality rates







France: female death rates (1816–2012)




French mortality rate models




French mortality rate models




French mortality rate models



French mortality rate forecasts




French mortality rate forecasts




Outline

- 1 Functional Partial Least Squares
- 2 Application: French mortality rates
- 3 Application: Australian fertility rates**
- 4 Forecast accuracy comparisons
- 5 Bootstrap intervals
- 6 Comparisons
- 7 References








Australian fertility rates









Australian fertility rates




Australian fertility rate models




Australian fertility rate models




Australian fertility rate models



Australian fertility rate forecasts



Australian fertility rate forecasts



Outline

- 1 Functional Partial Least Squares
- 2 Application: French mortality rates
- 3 Application: Australian fertility rates
- 4 Forecast accuracy comparisons**
- 5 Bootstrap intervals
- 6 Comparisons
- 7 References

Forecast accuracy comparisons

$$\text{MSE}_t = \frac{1}{p} \sum_{i=1}^p [y_t(x_i) - \hat{y}_{t|t-1}(x_i)]^2.$$

- Averaged over last m years of observed data.
- For French female mortality, $m = 30$
- For Australian fertility, $m = 20$
- For comparison, compare random walk:
 $y_{t+1|t}(x) = y_t(x).$

Forecast accuracy comparisons

$$\text{MSE}_t = \frac{1}{p} \sum_{i=1}^p [y_t(x_i) - \hat{y}_{t|t-1}(x_i)]^2.$$

- Averaged over last m years of observed data.
- For French female mortality, $m = 30$
- For Australian fertility, $m = 20$
- For comparison, compare random walk:
 $y_{t+1|t}(x) = y_t(x).$

Forecast accuracy comparisons

$$\text{MSE}_t = \frac{1}{p} \sum_{i=1}^p [y_t(x_i) - \hat{y}_{t|t-1}(x_i)]^2.$$

- Averaged over last m years of observed data.
- For French female mortality, $m = 30$
- For Australian fertility, $m = 20$
- For comparison, compare random walk:
 $y_{t+1|t}(x) = y_t(x).$


Forecast accuracy comparisons

$$\text{MSE}_t = \frac{1}{p} \sum_{i=1}^p [y_t(x_i) - \hat{y}_{t|t-1}(x_i)]^2.$$

- Averaged over last m years of observed data.
- For French female mortality, $m = 30$
- For Australian fertility, $m = 20$
- For comparison, compare random walk:
 $y_{t+1|t}(x) = y_t(x).$

Forecast accuracy comparisons

French female mortality rates




Forecast accuracy comparisons

MSE: French female mortality rates ($\times 1000$)

K	FPC	FPC_{w}	FPLSR	FPLSR_{w}	RW
1	0.5956	0.0293	0.5994	0.0607	
2	0.0537	0.0310	0.0738	0.0288	
3	0.0316	0.0310	0.0445	0.0288	
4	0.0296	0.0311	0.0428	0.0288	
5	0.0287	0.0311	0.0472	0.0297	
6	0.0425	0.0311	0.0474	0.0291	0.0437

Forecast accuracy comparisons

Australian fertility rates



Forecast accuracy comparisons

MSE: Australian fertility rates

K	FPC	FPC_w	FPLSR	$FPLSR_w$	RW
1	99.0611	16.7304	94.0311	53.8186	
2	56.3095	3.3019	54.3410	17.5883	
3	24.9330	3.2580	26.0428	10.2599	
4	15.6845	3.1995	19.7227	4.4818	
5	4.4495	3.2132	5.9299	4.0573	
6	3.4310	3.2123	4.9205	2.9046	4.9800

Computation time

- Weighted FPLSR more efficient than weighted FPC as FPC requires many univariate time series models.
- **Time to fit 100 replications:**

Method	Fertility data	Mortality data
FPC	34.1072	62.2797
FPC _w	33.1424	60.8426
FPLSR	0.4287	2.9184
FPLSR _w	0.4537	3.1602
RW	0.0000	0.0002

(Intel Xeon 2.33GHz processor)

Computation time

- Weighted FPLSR more efficient than weighted FPC as FPC requires many univariate time series models.
- **Time to fit 100 replications:**

Method	Fertility data	Mortality data
FPC	34.1072	62.2797
FPC _w	33.1424	60.8426
FPLSR	0.4287	2.9184
FPLSR _w	0.4537	3.1602
RW	0.0000	0.0002

(Intel Xeon 2.33GHz processor)

Outline

- 1 Functional Partial Least Squares
- 2 Application: French mortality rates
- 3 Application: Australian fertility rates
- 4 Forecast accuracy comparisons
- 5 Bootstrap intervals**
- 6 Comparisons
- 7 References

Sources of uncertainty

Functional PCA

- 1 smoothing error in estimating $s_t(x)$
- 2 error in estimating $\mu(x)$
- 3 error in forecasting the scores $\beta_{t,k}$
- 4 error in the model residuals $e_t(x)$
- 5 appropriateness of model

Sources of uncertainty

Functional PCA

- 1 smoothing error in estimating $s_t(x)$
- 2 error in estimating $\mu(x)$
- 3 error in forecasting the scores $\beta_{t,k}$
- 4 error in the model residuals $e_t(x)$
- 5 appropriateness of model

Sources of uncertainty

Functional PCA

- 1 smoothing error in estimating $s_t(x)$
- 2 error in estimating $\mu(x)$
- 3 error in forecasting the scores $\beta_{t,k}$
- 4 error in the model residuals $e_t(x)$
- 5 appropriateness of model

Sources of uncertainty

Functional PCA

- 1 smoothing error in estimating $s_t(x)$
- 2 error in estimating $\mu(x)$
- 3 error in forecasting the scores $\beta_{t,k}$
- 4 error in the model residuals $e_t(x)$
- 5 appropriateness of model

Sources of uncertainty

Functional PCA

- 1 smoothing error in estimating $s_t(x)$
- 2 error in estimating $\mu(x)$
- 3 error in forecasting the scores $\beta_{t,k}$
- 4 error in the model residuals $e_t(x)$
- 5 appropriateness of model

Sources of uncertainty

Functional PLS

- 1 smoothing error in estimating $s_t(x)$
- 2 error in estimating $\mu(x)$
- 3 error in estimating $b(x, u)$
- 4 error in the model residuals $e_t(x)$
- 5 appropriateness of model

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
- $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
- Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
- $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
- $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,j}, \dots, \hat{\varepsilon}_{T,j}\}$.
- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
- $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
- Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
- $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
- $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,j}, \dots, \hat{\varepsilon}_{T,j}\}$.
- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
 - $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
 - Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
 - $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
 - $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,j}, \dots, \hat{\varepsilon}_{T,j}\}$.
- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
- $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
- Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
- $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
- $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,j}, \dots, \hat{\varepsilon}_{T,j}\}$.

$$Y_{T+h|T}^{(\ell)}(x_i) = \mu(x_i) + \sum_{k=1}^K \beta_{T+h|T,k}^{(\ell)} \phi_k(x_i) + e^{(\ell)}(x_i) + \hat{\sigma}_{T+h}(x_i) \varepsilon^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
- $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
- Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
- $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
- $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,i}, \dots, \hat{\varepsilon}_{T,i}\}$.

$$y_{T+h|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \sum_{k=1}^K \beta_{T+h|T,k}^{(\ell)} \phi_k(x_i) + e^{(\ell)}(x_i) + \hat{\sigma}_{T+h}(x_i) \varepsilon^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
- $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
- Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
- $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
- $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,i}, \dots, \hat{\varepsilon}_{T,i}\}$.

$$y_{T+h|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \sum_{k=1}^K \beta_{T+h|T,k}^{(\ell)} \phi_k(x_i) + e^{(\ell)}(x_i) + \hat{\sigma}_{T+h}(x_i) \varepsilon^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPC

- Let $\hat{\xi}_{t,k} = \hat{\beta}_{t,k} - \hat{\beta}_{t|t-1,k}$ be 1-step errors of PC scores.
- $\{\xi_k^{(\ell)}\}$ sampled with replacement from $\{\hat{\xi}_{t,k}\}$.
- Simulate future sample paths of $\beta_{T+h|T,k}$ using these bootstrapped residuals: $\{\beta_{T+h|T,k}^{(\ell)}\}$.
- $\{e^{(\ell)}(x)\}$ sampled with replacement from residual functions $\{\hat{e}_1(x), \dots, \hat{e}_T(x)\}$.
- $\{\varepsilon^{(\ell)}\}$ sampled with replacement from $\{\hat{\varepsilon}_{1,i}, \dots, \hat{\varepsilon}_{T,i}\}$.

$$y_{T+h|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \sum_{k=1}^K \beta_{T+h|T,k}^{(\ell)} \phi_k(x_i) + e^{(\ell)}(x_i) + \hat{\sigma}_{T+h}(x_i) \varepsilon^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPLS

- Resample residuals and construct bootstrap samples:

$$\mathbf{f}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{f}}^*(x) + \mathbf{e}^{(\ell)}(x),$$

$$\mathbf{g}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{g}}^*(x) + \mathbf{e}^{(\ell)}(x).$$

- Construct weighted FPLSR model for each bootstrap sample.
- Compute bootstrapped forecast variants:

$$y_{T+1|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \int (\hat{f}_T(u) - \hat{\mu}(u)) \hat{b}^{(\ell)}(x_i, u) du + \hat{\sigma}_{T+1}(x_i) \varepsilon^{(\ell)}$$

Bootstrap curves for FPLS

- Resample residuals and construct bootstrap samples:

$$\mathbf{f}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{f}}^*(x) + \mathbf{e}^{(\ell)}(x),$$

$$\mathbf{g}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{g}}^*(x) + \mathbf{e}^{(\ell)}(x).$$

- Construct weighted FPLSR model for each bootstrap sample.
- Compute bootstrapped forecast variants:

$$y_{T+1|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \int (f_T(u) - \hat{\mu}(u)) \hat{b}^{(\ell)}(x_i, u) du + \hat{\sigma}_{T+1}(x_i) \hat{\varepsilon}^{(\ell)}$$

Bootstrap curves for FPLS

- Resample residuals and construct bootstrap samples:

$$\mathbf{f}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{f}}^*(x) + \mathbf{e}^{(\ell)}(x),$$

$$\mathbf{g}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{g}}^*(x) + \mathbf{e}^{(\ell)}(x).$$

- Construct weighted FPLSR model for each bootstrap sample.
- Compute bootstrapped forecast variants:

$$y_{T+1|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \int (f_T(u) - \hat{\mu}(u)) \hat{b}^{(\ell)}(x_i, u) du + \hat{\sigma}_{T+1}(x_i) \hat{\varepsilon}^{(\ell)}$$

Bootstrap curves for FPLS

- Resample residuals and construct bootstrap samples:

$$\mathbf{f}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{f}}^*(x) + \mathbf{e}^{(\ell)}(x),$$

$$\mathbf{g}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{g}}^*(x) + \mathbf{e}^{(\ell)}(x).$$

- Construct weighted FPLSR model for each bootstrap sample.
- Compute bootstrapped forecast variants:

$$y_{T+1|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \int (f_T(u) - \hat{\mu}(u)) \hat{b}^{(\ell)}(x_i, u) du + \hat{\sigma}_{T+1}(x_i) \hat{\varepsilon}^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPLS

- Resample residuals and construct bootstrap samples:

$$\mathbf{f}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{f}}^*(x) + \mathbf{e}^{(\ell)}(x),$$

$$\mathbf{g}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{g}}^*(x) + \mathbf{e}^{(\ell)}(x).$$

- Construct weighted FPLSR model for each bootstrap sample.
- Compute bootstrapped forecast variants:

$$y_{T+1|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \int (f_T(u) - \hat{\mu}(u)) \hat{b}^{(\ell)}(x_i, u) du + \hat{\sigma}_{T+1}(x_i) \hat{\varepsilon}^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Bootstrap curves for FPLS

- Resample residuals and construct bootstrap samples:

$$\mathbf{f}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{f}}^*(x) + \mathbf{e}^{(\ell)}(x),$$

$$\mathbf{g}^{(\ell)}(x) = \hat{\mu}(x) + \hat{\mathbf{g}}^*(x) + \mathbf{e}^{(\ell)}(x).$$

- Construct weighted FPLSR model for each bootstrap sample.
- Compute bootstrapped forecast variants:

$$y_{T+1|T}^{(\ell)}(x_i) = \hat{\mu}(x_i) + \int (f_T(u) - \hat{\mu}(u)) \hat{b}^{(\ell)}(x_i, u) du + \hat{\sigma}_{T+1}(x_i) \hat{\varepsilon}^{(\ell)}$$

- Prediction intervals are produced from the bootstrap variants using percentiles.

Coverage probability

Empirical coverage probability of 95% intervals

$$\frac{1}{mph} \sum_{t=T-m+1}^T \sum_{j=1}^h \sum_{i=1}^p \mathbf{1} \left(\hat{y}_{t+j|t}^{(0.025)}(\mathbf{x}_i) < y_{t+j}(\mathbf{x}_i) < \hat{y}_{t+j|t}^{(0.975)}(\mathbf{x}_i) \right)$$

- $\hat{y}_{t+j|t}^{(\alpha)}(\mathbf{x}_i) = \alpha$ -quantile from bootstrap samples
- m = smallest number of observations used to fit a model

Method	Fertility data	Mortality data
FPC	98.00%	97.19%
FPLS	96.86%	97.23%

Coverage probability

Empirical coverage probability of 95% intervals

$$\frac{1}{mph} \sum_{t=T-m+1}^T \sum_{j=1}^h \sum_{i=1}^p \mathbf{1} \left(\hat{y}_{t+j|t}^{(0.025)}(x_i) < y_{t+j}(x_i) < \hat{y}_{t+j|t}^{(0.975)}(x_i) \right)$$

- $\hat{y}_{t+j|t}^{(\alpha)}(x_i)$ = α -quantile from bootstrap samples
- m = smallest number of observations used to fit a model

Method	Fertility data	Mortality data
FPC	98.00%	97.19%
FPLS	96.86%	97.23%

Coverage probability

Empirical coverage probability of 95% intervals

$$\frac{1}{mph} \sum_{t=T-m+1}^T \sum_{j=1}^h \sum_{i=1}^p \mathbf{1} \left(\hat{y}_{t+j|t}^{(0.025)}(\mathbf{x}_i) < y_{t+j}(\mathbf{x}_i) < \hat{y}_{t+j|t}^{(0.975)}(\mathbf{x}_i) \right)$$

- $\hat{y}_{t+j|t}^{(\alpha)}(\mathbf{x}_i)$ = α -quantile from bootstrap samples
- m = smallest number of observations used to fit a model

Method	Fertility data	Mortality data
FPC	98.00%	97.19%
FPLS	96.86%	97.23%

Coverage probability

Empirical coverage probability of 95% intervals

$$\frac{1}{m p h} \sum_{t=T-m+1}^T \sum_{j=1}^h \sum_{i=1}^p \mathbf{1} \left(\hat{y}_{t+j|t}^{(0.025)}(\mathbf{x}_i) < y_{t+j}(\mathbf{x}_i) < \hat{y}_{t+j|t}^{(0.975)}(\mathbf{x}_i) \right)$$

- $\hat{y}_{t+j|t}^{(\alpha)}(\mathbf{x}_i) = \alpha$ -quantile from bootstrap samples
- m = smallest number of observations used to fit a model

Method	Fertility data	Mortality data
FPC	98.00%	97.19%
FPLS	96.86%	97.23%

Adjusted prediction intervals

- Compute $d(x)$ = difference between $(1 - \alpha/2)$ and $(\alpha/2)$ quantiles of $\{\hat{f}_{t+1}(x) - \hat{f}_{t+1|t}(x); t = m, \dots, T - 1\}$
- Let $[\ell_h(x), u_h(x)]$ be $100(1 - \alpha)\%$ h -step-ahead prediction intervals obtained from bootstrap methods.
- Ideally, $u_1(x) - \ell_1(x) = d(x)$.

Adjusted prediction interval

$$\left[0.5\{\ell_h(x) + u_h(x)\} - \{u_h(x) - \ell_h(x)\}p(x), \right. \\ \left. 0.5\{\ell_h(x) + u_h(x)\} + \{u_h(x) - \ell_h(x)\}p(x) \right]$$

where $p(x) = 0.5d(x)/[u_1(x) - \ell_1(x)]$.

Adjusted prediction intervals

- Compute $d(x)$ = difference between $(1 - \alpha/2)$ and $(\alpha/2)$ quantiles of $\{\hat{f}_{t+1}(x) - \hat{f}_{t+1|t}(x); t = m, \dots, T - 1\}$
- Let $[\ell_h(x), u_h(x)]$ be $100(1 - \alpha)\%$ h -step-ahead prediction intervals obtained from bootstrap methods.
- Ideally, $u_1(x) - \ell_1(x) = d(x)$.

Adjusted prediction interval

$$\left[0.5\{\ell_h(x) + u_h(x)\} - \{u_h(x) - \ell_h(x)\}p(x), \right. \\ \left. 0.5\{\ell_h(x) + u_h(x)\} + \{u_h(x) - \ell_h(x)\}p(x) \right]$$

where $p(x) = 0.5d(x)/[u_1(x) - \ell_1(x)]$.

Adjusted prediction intervals

- Compute $d(x)$ = difference between $(1 - \alpha/2)$ and $(\alpha/2)$ quantiles of $\{\hat{f}_{t+1}(x) - \hat{f}_{t+1|t}(x); t = m, \dots, T - 1\}$
- Let $[\ell_h(x), u_h(x)]$ be $100(1 - \alpha)\%$ h -step-ahead prediction intervals obtained from bootstrap methods.
- Ideally, $u_1(x) - \ell_1(x) = d(x)$.

Adjusted prediction interval

$$\left[0.5\{\ell_h(x) + u_h(x)\} - \{u_h(x) - \ell_h(x)\}p(x), \right. \\ \left. 0.5\{\ell_h(x) + u_h(x)\} + \{u_h(x) - \ell_h(x)\}p(x) \right]$$

where $p(x) = 0.5d(x)/[u_1(x) - \ell_1(x)]$.

Adjusted prediction intervals

- Compute $d(x)$ = difference between $(1 - \alpha/2)$ and $(\alpha/2)$ quantiles of $\{\hat{f}_{t+1}(x) - \hat{f}_{t+1|t}(x); t = m, \dots, T - 1\}$
- Let $[\ell_h(x), u_h(x)]$ be $100(1 - \alpha)\%$ h -step-ahead prediction intervals obtained from bootstrap methods.
- Ideally, $u_1(x) - \ell_1(x) = d(x)$.

Adjusted prediction interval

$$\left[0.5\{\ell_h(x) + u_h(x)\} - \{u_h(x) - \ell_h(x)\}p(x), \right. \\ \left. 0.5\{\ell_h(x) + u_h(x)\} + \{u_h(x) - \ell_h(x)\}p(x) \right]$$

where $p(x) = 0.5d(x)/[u_1(x) - \ell_1(x)]$.

- For $h = 1$, these have same coverage as in-sample intervals if distributions symmetric

Adjusted prediction intervals

- Compute $d(x)$ = difference between $(1 - \alpha/2)$ and $(\alpha/2)$ quantiles of $\{\hat{f}_{t+1}(x) - \hat{f}_{t+1|t}(x); t = m, \dots, T - 1\}$
- Let $[\ell_h(x), u_h(x)]$ be $100(1 - \alpha)\%$ h -step-ahead prediction intervals obtained from bootstrap methods.
- Ideally, $u_1(x) - \ell_1(x) = d(x)$.

Adjusted prediction interval

$$\left[0.5\{\ell_h(x) + u_h(x)\} - \{u_h(x) - \ell_h(x)\}p(x), \right. \\ \left. 0.5\{\ell_h(x) + u_h(x)\} + \{u_h(x) - \ell_h(x)\}p(x) \right]$$

where $p(x) = 0.5d(x)/[u_1(x) - \ell_1(x)]$.

- For $h = 1$, these have same coverage as in-sample intervals if distributions symmetric.

Adjusted prediction intervals

- Compute $d(x)$ = difference between $(1 - \alpha/2)$ and $(\alpha/2)$ quantiles of $\{\hat{f}_{t+1}(x) - \hat{f}_{t+1|t}(x); t = m, \dots, T - 1\}$
- Let $[\ell_h(x), u_h(x)]$ be $100(1 - \alpha)\%$ h -step-ahead prediction intervals obtained from bootstrap methods.
- Ideally, $u_1(x) - \ell_1(x) = d(x)$.

Adjusted prediction interval

$$\left[0.5\{\ell_h(x) + u_h(x)\} - \{u_h(x) - \ell_h(x)\}p(x), \right. \\ \left. 0.5\{\ell_h(x) + u_h(x)\} + \{u_h(x) - \ell_h(x)\}p(x) \right]$$


where $p(x) = 0.5d(x)/[u_1(x) - \ell_1(x)]$.

- For $h = 1$, these have same coverage as in-sample intervals if distributions symmetric.


Adjusted prediction intervals

	Fertility data		Mortality data	
	95%	adj 95%	95%	adj 95%
FPC	98.00%	95.86%	97.19%	95.91%
FPLS	96.86%	94.89%	97.23%	94.95%


Adjusted prediction intervals




Adjusted prediction intervals



Adjusted prediction intervals



Adjusted prediction intervals



Outline

- 1 Functional Partial Least Squares
- 2 Application: French mortality rates
- 3 Application: Australian fertility rates
- 4 Forecast accuracy comparisons
- 5 Bootstrap intervals
- 6 Comparisons**
- 7 References

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- latent components more suitable for prediction
- easier than variance decomposition

Both methods implemented in fpls package for R

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- latent components more suitable for prediction rather than variance decomposition.
- can be used to model non-linear relationships

Both methods implemented in `fda` package for R

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- latent components more suitable for prediction rather than variance decomposition.
- Faster as there is no need to fit univariate time series models

Both methods implemented in `fplsa` package for R

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- 1 latent components more suitable for prediction rather than variance decomposition.
- 2 Faster as there is no need to fit univariate time series models

Both methods implemented in `ftsa` package for R.

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- 1 latent components more suitable for prediction rather than variance decomposition.
- 2 Faster as there is no need to fit univariate time series models

Both methods implemented in `ftsa` package for R.

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- 1 latent components more suitable for prediction rather than variance decomposition.
- 2 Faster as there is no need to fit univariate time series models

Both methods implemented in `ftsa` package for R.

Comparisons

FPC advantages

- 1 allows more complex dynamics and higher order components;
- 2 eases interpretability of dynamic changes by separating out effects of a few orthogonal components;

FPLS advantages

- 1 latent components more suitable for prediction rather than variance decomposition.
- 2 Faster as there is no need to fit univariate time series models

Both methods implemented in `ftsa` package for R.

Outline

- 1 Functional Partial Least Squares
- 2 Application: French mortality rates
- 3 Application: Australian fertility rates
- 4 Forecast accuracy comparisons
- 5 Bootstrap intervals
- 6 Comparisons
- 7 References**

Selected references



Hyndman, Shang (2009). “Forecasting functional time series (with discussion)”. *Journal of the Korean Statistical Society* **38**(3), 199–221.



Hyndman (2014). *demography: Forecasting mortality, fertility, migration and population data*.

cran.r-project.org/package=demography



Shang, Hyndman (2013). *ftsa: Functional time series analysis*.

cran.r-project.org/package=ftsa