




Rob J Hyndman

Optimal forecast reconciliation for big time series data



Follow along using R



Requirements

Install the hts package and its dependencies.

Outline

- 1 Hierarchical and grouped time series**
- 2 BLUF: Best Linear Unbiased Forecasts
- 3 Application: Australian tourism
- 4 Temporal hierarchies
- 5 References

Labour market participation

Australia and New Zealand Standard Classification of Occupations

- 8 major groups
 - 43 sub-major groups
 - 97 minor groups
 - 359 unit groups
 - * 1023 occupations

Example: statistician

2 Professionals

22 Business, Human Resource and Marketing Professionals

224 Information and Organisation Professionals

2241 Actuaries, Mathematicians and Statisticians

224113 Statistician

Labour market participation

Australia and New Zealand Standard Classification of Occupations

- 8 major groups
 - 43 sub-major groups
 - 97 minor groups
 - 359 unit groups
 - * 1023 occupations

Example: statistician

2 Professionals


22 Business, Human Resource and Marketing Professionals

224 Information and Organisation Professionals

2241 Actuaries, Mathematicians and Statisticians

224113 Statistician

Australian tourism demand




Australian tourism demand

- Quarterly data on visitor night from 1998:Q1 – 2013:Q4
- From: *National Visitor Survey*, based on annual interviews of 120,000 Australians aged 15+, collected by Tourism Research Australia.
- Split by 7 states, 27 zones and 76 regions (a geographical hierarchy)
- Also split by purpose of travel
 - Holiday
 - Visiting friends and relatives (VFR)
 - Business
 - Other
- 304 bottom-level series



Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




Examples

- Labour turnover by occupation
- Tourism by state and region

Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




Examples

- Labour turnover by occupation
- Tourism by state and region

Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




Examples

- Labour turnover by occupation
- Tourism by state and region

Grouped time series

A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.




Examples

- Labour turnover by occupation and state
- Tourism by state and purpose of travel

Grouped time series

A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.




Examples

- Labour turnover by occupation and state
- Tourism by state and purpose of travel

Grouped time series


A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.



Examples

- Labour turnover by occupation and state
- Tourism by state and purpose of travel

Hierarchical time series




y_t : observed aggregate of all series at time t .

$y_{X,t}$: observation on series X at time t .

\mathbf{b}_t : vector of all series at bottom level in time t .

Hierarchical time series




y_t : observed aggregate of all series at time t .

$y_{X,t}$: observation on series X at time t .

b_t : vector of all series at bottom level in time t .

Hierarchical time series




y_t : observed aggregate of all series at time t .

$y_{X,t}$: observation on series X at time t .

\mathbf{b}_t : vector of all series at bottom level in time t .

$$\mathbf{y}_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}$$

Hierarchical time series




y_t : observed aggregate of all series at time t .

$y_{X,t}$: observation on series X at time t .

\mathbf{b}_t : vector of all series at bottom level in time t .

$$\mathbf{y}_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}$$

Hierarchical time series




y_t : observed aggregate of all series at time t .

$y_{X,t}$: observation on series X at time t .

\mathbf{b}_t : vector of all series at bottom level in time t .

$$\mathbf{y}_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}}_{\mathbf{b}_t}$$

Hierarchical time series



y_t : observed aggregate of all series at time t .


$y_{X,t}$: observation on series X at time t .

\mathbf{b}_t : vector of all series at bottom level in time t .

$$\mathbf{y}_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}}_{\mathbf{b}_t}$$


$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$

Hierarchical time series




$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$

Hierarchical time series



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$


Hierarchical time series



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$


$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

Grouped data




$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix}}_{\mathbf{b}_t}$$

Grouped data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix}}_{\mathbf{b}_t}$$

Grouped data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix}}_{\mathbf{b}_t}$$

$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

Hierarchical and grouped time series

Every collection of time series with aggregation constraints can be written as

$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

where

- \mathbf{y}_t is a vector of all series at time t
- \mathbf{b}_t is a vector of the most disaggregated series at time t
- \mathbf{S} is a “summing matrix” containing the aggregation constraints.

hts package for R



hts: Hierarchical and grouped time series

Methods for analysing and forecasting hierarchical and grouped time series

Version: 4.5
Depends: forecast (\geq 5.0), SparseM
Imports: parallel, utils
Published: 2015-06-29
Author: Rob J Hyndman, Earo Wang and Alan Lee
with contributions from Shanika Wickramasuriya
Maintainer: Rob J Hyndman <Rob.Hyndman at monash.edu>
BugReports: <https://github.com/robjhyndman/hts/issues>
License: GPL ($>$ 2)

Example using R


```
library(hts)
```

```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))
```

Example using R

```
library(hts)
```


```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))
```



Example using R

```
library(hts)
```

```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))
```



```
summary(y)  
smatrix(y)
```

Example using R

Example 1 illustrating the usage of the "groups" argument

```
abc <- ts(5 + matrix(sort(rnorm(1600))), ncol = 16, nrow = 100))

sex <- rep(c("female", "male"), each = 8)
state <- rep(c("NSW", "VIC", "QLD", "SA", "WA", "NT", "ACT", "TAS"), 2)

gc <- rbind(sex, state) # a matrix consists of strings.
gn <- rbind(rep(1:2, each = 8), rep(1:8, 2))
rownames(gc) <- rownames(gn) <- c("Sex", "State")

x <- gts(abc, groups = gc)
y <- gts(abc, groups = gn)

plot(x, level=3)
plot(x, level=2)
plot(x, level=1)
plot(x, level=0)
```

Example using R

Example 2 with two simple hierarchies (geography and product) to show the argument "characters"

```
bnames1 <- c("VICMelbA1", "VICMelbA2", "VICGeelA1", "VICGeelA2",  
            "VICMelbB1", "VICMelbB2", "VICGeelB1", "VICGeelB2",  
            "NSWSyndA1", "NSWSyndA2", "NSWollA1", "NSWollA2",  
            "NSWSyndB1", "NSWSyndB2", "NSWollB1", "NSWollB2")  
  
bts1 <- matrix(ts(rnorm(160))), ncol = 16)  
colnames(bts1) <- bnames1  
  
x1 <- gts(bts1, characters = list(c(3, 4), c(1, 1)),  
        gnames = c("State", "State.City", "Product", "Product.Size",  
                    "State.Product", "State.Product.Size", "State.City.Product"))  
  
plot(x1, level=1)  
plot(x1, level=7)
```


Example using R

Example 3 with a non-hierarchical grouped time series of 3 grouping variables (state, age and sex)

```
bnames2 <- c("VIC1F","VIC1M","VIC2F","VIC2M","VIC3F","VIC3M",  
            "NSW1F","NSW1M","NSW2F","NSW2M","NSW3F","NSW3M")  
bts2 <- matrix(ts(rnorm(120)), ncol = 12)  
colnames(bts2) <- bnames2  
x2 <- gts(bts2, characters = c(3, 1, 1),  
         gnames=c("State","Age","Sex",  
                  "State.Age","State.Sex","Age.Sex"))  
  
plot(x2, level=3)
```

Outline

- 1 Hierarchical and grouped time series
- 2 BLUF: Best Linear Unbiased Forecasts**
- 3 Application: Australian tourism
- 4 Temporal hierarchies
- 5 References

Forecasting notation

Let $\hat{\mathbf{y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{y}_t . (They may not add up.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

for some matrix \mathbf{P} .

Forecasting notation

Let $\hat{\mathbf{y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{y}_t . (They may not add up.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{y}}_n(h)$ to get bottom-level forecasts.

■ \mathbf{S} sums them up

Forecasting notation

Let $\hat{\mathbf{y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{y}_t . (They may not add up.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up

Forecasting notation

Let $\hat{\mathbf{y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{y}_t . (They may not add up.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up

Forecasting notation

Let $\hat{\mathbf{y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{y}_t . (They may not add up.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:

$$E[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = E[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = E[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.

$$\hat{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{b}}_n(h)$$

We want the revised forecasts to be

$$\text{unbiased: } E[\tilde{\mathbf{y}}_n(h)] = \mathbf{SP}E[\hat{\mathbf{y}}_n(h)] = \mathbf{SP}\beta_n(h)$$

Revised forecasts are unbiased iff $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Revised forecasts are unbiased iff $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\boldsymbol{\beta}_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\boldsymbol{\beta}_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\boldsymbol{\beta}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h)$.

Revised forecasts are unbiased iff $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:
$$\mathbb{E}[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Revised forecasts are unbiased iff $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Revised forecasts are unbiased iff $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Revised forecasts are unbiased iff $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let $\hat{\mathbf{b}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Revised forecasts are unbiased iff **$\mathbf{SPS} = \mathbf{S}$** .

General properties: variance

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Let error variance of h -step base forecasts $\hat{\mathbf{y}}_n(h)$ be

$$\mathbf{W}_h = \text{Var}[\mathbf{y}_{n+h} - \hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

Then the error variance of the corresponding revised forecasts is

$$\text{Var}[\mathbf{y}_{n+h} - \tilde{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{SPW}_h\mathbf{P}'\mathbf{S}'$$

General properties: variance

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

Let error variance of h -step base forecasts $\hat{\mathbf{y}}_n(h)$ be

$$\mathbf{W}_h = \text{Var}[\mathbf{y}_{n+h} - \hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

Then the error variance of the corresponding revised forecasts is

$$\text{Var}[\mathbf{y}_{n+h} - \tilde{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{SPW}_h\mathbf{P}'\mathbf{S}'$$

BLUF via trace minimization

Theorem

For any \mathbf{P} satisfying $\mathbf{SPS} = \mathbf{S}$, then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{SPW}_h\mathbf{P}'\mathbf{S}']$$

has solution $\mathbf{P} = (\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger$.

- \mathbf{W}_h^\dagger is generalized inverse of \mathbf{W}_h .
- **Problem:** \mathbf{W}_h hard to estimate, especially for $h > 1$.

BLUF via trace minimization

Theorem

For any \mathbf{P} satisfying $\mathbf{SPS} = \mathbf{S}$, then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{SPW}_h\mathbf{P}'\mathbf{S}']$$

has solution $\mathbf{P} = (\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger$.

- \mathbf{W}_h^\dagger is generalized inverse of \mathbf{W}_h .
- **Problem:** \mathbf{W}_h hard to estimate, especially for $h > 1$.

BLUF via trace minimization

Theorem

For any \mathbf{P} satisfying $\mathbf{SPS} = \mathbf{S}$, then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{SPW}_h\mathbf{P}'\mathbf{S}']$$

has solution $\mathbf{P} = (\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger$.

- \mathbf{W}_h^\dagger is generalized inverse of \mathbf{W}_h .
- **Problem:** \mathbf{W}_h hard to estimate, especially for $h > 1$.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Assume forecast errors have the same aggregation constraints as the data.
- Then $\mathbf{W}_h = \mathbf{S}\Omega_h\mathbf{S}'$ where Ω_h is covariance matrix of bottom level errors.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Assume forecast errors have the same aggregation constraints as the data.
- Then $\mathbf{W}_h = \mathbf{S}\mathbf{\Omega}_h\mathbf{S}'$ where $\mathbf{\Omega}_h$ is covariance matrix of bottom level errors.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Assume forecast errors have the same aggregation constraints as the data.
- Then $\mathbf{W}_h = \mathbf{S}\mathbf{\Omega}_h\mathbf{S}'$ where $\mathbf{\Omega}_h$ is covariance matrix of bottom level errors.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Assume forecast errors have the same aggregation constraints as the data.
- Then $\mathbf{W}_h = \mathbf{S}\mathbf{\Omega}_h\mathbf{S}'$ where $\mathbf{\Omega}_h$ is covariance matrix of bottom level errors.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Assume forecast errors have the same aggregation constraints as the data.
- Then $\mathbf{W}_h = \mathbf{S}\mathbf{\Omega}_h\mathbf{S}'$ where $\mathbf{\Omega}_h$ is covariance matrix of bottom level errors.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Assume forecast errors have the same aggregation constraints as the data.
- Then $\mathbf{W}_h = \mathbf{S}\mathbf{\Omega}_h\mathbf{S}'$ where $\mathbf{\Omega}_h$ is covariance matrix of bottom level errors.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 2: WLS

- Suppose we approximate \mathbf{W}_1 by its diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Easy to estimate, and places weight where we have best forecasts.
- Empirically, WLS gives better forecasts than OLS.
- Still need to estimate covariance matrix to get prediction intervals.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 2: WLS

- Suppose we approximate \mathbf{W}_1 by its diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Easy to estimate, and places weight where we have best forecasts.
- Empirically, WLS gives better forecasts than OLS.
- Still need to estimate covariance matrix to produce prediction intervals.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 2: WLS

- Suppose we approximate \mathbf{W}_1 by its diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Easy to estimate, and places weight where we have best forecasts.
- Empirically, WLS gives better forecasts than OLS.
- Still need to estimate covariance matrix to produce prediction intervals.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 2: WLS

- Suppose we approximate \mathbf{W}_1 by its diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Easy to estimate, and places weight where we have best forecasts.
- Empirically, WLS gives better forecasts than OLS.
- Still need to estimate covariance matrix to produce prediction intervals.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 2: WLS

- Suppose we approximate \mathbf{W}_1 by its diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Easy to estimate, and places weight where we have best forecasts.
- Empirically, WLS gives better forecasts than OLS.
- Still need to estimate covariance matrix to produce prediction intervals.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 3: GLS

- Estimate \mathbf{W}_1 using shrinkage to the diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Allows for covariances.
- Empirically, GLS gives better forecasts than WLS or OLS.
- Difficult to compute for large number of time series.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 3: GLS

- Estimate \mathbf{W}_1 using shrinkage to the diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Allows for covariances.
- Empirically, GLS gives better forecasts than WLS or OLS.
- Difficult to compute for large numbers of time series.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 3: GLS

- Estimate \mathbf{W}_1 using shrinkage to the diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Allows for covariances.
- Empirically, GLS gives better forecasts than WLS or OLS.
- Difficult to compute for large numbers of time series.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 3: GLS

- Estimate \mathbf{W}_1 using shrinkage to the diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Allows for covariances.
- Empirically, GLS gives better forecasts than WLS or OLS.
- Difficult to compute for large numbers of time series.

Optimal combination forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^\dagger\hat{\mathbf{y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 3: GLS

- Estimate \mathbf{W}_1 using shrinkage to the diagonal and assume that $\mathbf{W}_h \propto \mathbf{W}_1$.
- Allows for covariances.
- Empirically, GLS gives better forecasts than WLS or OLS.
- Difficult to compute for large numbers of time series.

Example using R

```
library(hts)
```

```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))
```

```
# Forecast 10-step-ahead using WLS combination method  
# ETS used for each series by default  
fc <- forecast(y, h=10)
```

```
# Select your own methods  
ally <- aggts(y)  
allf <- matrix(, nrow=10, ncol=ncol(ally))  
for(i in 1:ncol(ally))  
  allf[,i] <- mymethod(ally[,i], h=10)  
allf <- ts(allf, start=2004)  
# Reconcile forecasts so they add up  
fc2 <- combinef(allf, nodes=y$nodes)
```

forecast.gts function

Usage

```
forecast(object, h,  
  method = c("comb", "bu", "mo", "tdgsf", "tdgsa", "tdfp"),  
  fmethod = c("ets", "rw", "arima"),  
  weights = c("sd", "none", "nseries"),  
  positive = FALSE,  
  parallel = FALSE, num.cores = 2, ...)
```

Arguments

<code>object</code>	Hierarchical time series object of class gts.
<code>h</code>	Forecast horizon
<code>method</code>	Method for distributing forecasts within the hierarchy.
<code>fmethod</code>	Forecasting method to use
<code>positive</code>	If TRUE, forecasts are forced to be strictly positive
<code>weights</code>	Weights used for "optimal combination" method. When <code>weights = "sd"</code> , it takes account of the standard deviation of forecasts.
<code>parallel</code>	If TRUE, allow parallel processing
<code>num.cores</code>	If <code>parallel = TRUE</code> , specify how many cores are going to be used

Exercise


Use the `infantgts` data to:

- 1 Plot various levels of aggregation.
- 2 Forecast the series using `auto.arima` with the default WLS reconciliation method.
- 3 Plot the reconciled forecasts at various levels of aggregation.

Outline

- 1 Hierarchical and grouped time series
- 2 BLUF: Best Linear Unbiased Forecasts
- 3 Application: Australian tourism**
- 4 Temporal hierarchies
- 5 References


Australian tourism



Australian tourism

Hierarchy:

- States (7)
- Zones (27)
- Regions (82)



Australian tourism

Hierarchy:

- States (7)
- Zones (27)
- Regions (82)

Base forecasts


ETS (exponential smoothing) models

□ > 3%




Base forecasts

Domestic tourism forecasts: Total




Base forecasts

Domestic tourism forecasts: NSW




Base forecasts

Domestic tourism forecasts: VIC




Base forecasts

Domestic tourism forecasts: Nth.Coast.NSW




Base forecasts

Domestic tourism forecasts: Metro.QLD




Base forecasts

Domestic tourism forecasts: Sth.WA




Base forecasts

Domestic tourism forecasts: X201.Melbourne




Base forecasts

Domestic tourism forecasts: X402.Murraylands




Base forecasts

Domestic tourism forecasts: X809.Daly



Forecast evaluation


Training sets



Forecast evaluation

Training sets


Test set $h = 1$



Forecast evaluation

Training sets


Test set $h = 2$



Forecast evaluation

Training sets


Test set $h = 3$



Forecast evaluation

Training sets


Test set $h = 4$



Forecast evaluation

Training sets


Test set $h = 5$



Forecast evaluation

Training sets

Test set $h = 6$




Hierarchy: states, zones, regions

RMSE	Forecast horizon						Ave
	$h = 1$	$h = 2$	$h = 3$	$h = 4$	$h = 5$	$h = 6$	
Australia							
Base	1762.04	1770.29	1766.02	1818.82	1705.35	1721.17	1757.28
Bottom	1736.92	1742.69	1722.79	1752.74	1666.73	1687.43	1718.22
OLS	1747.60	1757.68	1751.77	1800.67	1686.00	1706.45	1741.69
WLS	1705.21	1715.87	1703.75	1729.56	1627.79	1661.24	1690.57
GLS	1704.64	1715.60	1705.31	1729.04	1626.36	1661.64	1690.43
States							
Base	399.77	404.16	401.92	407.26	395.38	401.17	401.61
Bottom	404.29	406.95	404.96	409.02	399.80	401.55	404.43
OLS	404.47	407.62	405.43	413.79	401.10	404.90	406.22
WLS	398.84	402.12	400.71	405.03	394.76	398.23	399.95
GLS	398.84	402.16	400.86	405.03	394.59	398.22	399.95
Regions							
Base	93.15	93.38	93.45	93.79	93.50	93.56	93.47
Bottom	93.15	93.38	93.45	93.79	93.50	93.56	93.47
OLS	93.28	93.53	93.64	94.17	93.78	93.88	93.71
WLS	93.02	93.32	93.38	93.72	93.39	93.53	93.39
GLS	92.98	93.27	93.34	93.66	93.34	93.46	93.34

Outline

- 1 Hierarchical and grouped time series
- 2 BLUF: Best Linear Unbiased Forecasts
- 3 Application: Australian tourism
- 4 Temporal hierarchies**
- 5 References


Temporal hierarchies



Basic idea:

- ➡ Forecast series at each available frequency.
- ➡ Optimally combine forecasts within the same year.


Temporal hierarchies



Basic idea:

- ➡ Forecast series at each available frequency.
- ➡ Optimally combine forecasts within the same year.

Monthly series




■ $k = 2, 4, 12$ nodes

■ $k = 3, 6, 12$ nodes

■ Why not $k = 2, 3, 4, 6, 12$ nodes?

Monthly series




■ $k = 2, 4, 12$ nodes

■ $k = 3, 6, 12$ nodes

■ Why not $k = 2, 3, 4, 6, 12$ nodes?

Monthly series



- $k = 2, 4, 12$ nodes
- $k = 3, 6, 12$ nodes
- Why not $k = 2, 3, 4, 6, 12$ nodes?

Monthly data

$$\underbrace{\begin{pmatrix} A \\ \text{Semi}A_1 \\ \text{Semi}A_2 \\ \text{Four}M_1 \\ \text{Four}M_2 \\ \text{Four}M_3 \\ Q_1 \\ \vdots \\ Q_4 \\ \text{Bi}M_1 \\ \vdots \\ \text{Bi}M_6 \\ M_1 \\ \vdots \\ M_{12} \end{pmatrix}}_{(28 \times 1)} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & \vdots & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & \vdots & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix}}_{\substack{I_{12} \\ S}} \underbrace{\begin{pmatrix} M_1 \\ M_2 \\ M_3 \\ M_4 \\ M_5 \\ M_6 \\ M_7 \\ M_8 \\ M_9 \\ M_{10} \\ M_{11} \\ M_{12} \end{pmatrix}}_{B_t}$$

In general

For a time series y_1, \dots, y_T , observed at frequency m , we generate aggregate series

$$y_j^{[k]} = \sum_{t=1+(j-1)k}^{jk} y_t, \quad \text{for } j = 1, \dots, \lfloor T/k \rfloor$$

- $k \in F(m) = \{\text{factors of } m\}$.
- A single unique hierarchy is only possible when there are no coprime pairs in $F(m)$.
- $M_k = m/k$ is seasonal period of aggregated series.

In general

For a time series y_1, \dots, y_T , observed at frequency m , we generate aggregate series

$$y_j^{[k]} = \sum_{t=1+(j-1)k}^{jk} y_t, \quad \text{for } j = 1, \dots, \lfloor T/k \rfloor$$

- $k \in F(m) = \{\text{factors of } m\}$.
- A single unique hierarchy is only possible when there are no coprime pairs in $F(m)$.
- $M_k = m/k$ is seasonal period of aggregated series.

In general

For a time series y_1, \dots, y_T , observed at frequency m , we generate aggregate series

$$y_j^{[k]} = \sum_{t=1+(j-1)k}^{jk} y_t, \quad \text{for } j = 1, \dots, \lfloor T/k \rfloor$$

- $k \in F(m) = \{\text{factors of } m\}$.
- A single unique hierarchy is only possible when there are no coprime pairs in $F(m)$.
- $M_k = m/k$ is seasonal period of aggregated series.

WLS weights

Hierarchy variance scaling Λ_H : diagonal.

Series variance scaling Λ_V : elements equal within aggregation level.

Structural scaling $\Lambda_S = \text{diag}(\mathbf{S}\mathbf{1})$: elements equal to # nodes at each level.

- Depends only on seasonal period m .
- Independent of data and model.
- Allows forecasts where no errors available.

Quarterly example

$$\Lambda_H = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_{S_1}^2, \hat{\sigma}_{S_2}^2, \hat{\sigma}_{Q_1}^2, \hat{\sigma}_{Q_2}^2, \hat{\sigma}_{Q_3}^2, \hat{\sigma}_{Q_4}^2)$$

$$\Lambda_V = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_S^2, \hat{\sigma}_S^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2)$$

$$\Lambda_S = \text{diag}(4, 2, 2, 1, 1, 1, 1)$$

WLS weights

Hierarchy variance scaling Λ_H : diagonal.

Series variance scaling Λ_V : elements equal within aggregation level.

Structural scaling $\Lambda_S = \text{diag}(\mathbf{S}\mathbf{1})$: elements equal to # nodes at each level.

- Depends only on seasonal period m .
- Independent of data and model.
- Allows forecasts where no errors available.

Quarterly example

$$\Lambda_H = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_{S_1}^2, \hat{\sigma}_{S_2}^2, \hat{\sigma}_{Q_1}^2, \hat{\sigma}_{Q_2}^2, \hat{\sigma}_{Q_3}^2, \hat{\sigma}_{Q_4}^2)$$

$$\Lambda_V = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_S^2, \hat{\sigma}_S^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2)$$

$$\Lambda_S = \text{diag}(4, 2, 2, 1, 1, 1, 1)$$

WLS weights

Hierarchy variance scaling Λ_H : diagonal.

Series variance scaling Λ_V : elements equal within aggregation level.

Structural scaling $\Lambda_S = \text{diag}(\mathbf{S}\mathbf{1})$: elements equal to # nodes at each level.

- Depends only on seasonal period m .
- Independent of data and model.
- Allows forecasts where no errors available.

Quarterly example

$$\Lambda_H = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_{S_1}^2, \hat{\sigma}_{S_2}^2, \hat{\sigma}_{Q_1}^2, \hat{\sigma}_{Q_2}^2, \hat{\sigma}_{Q_3}^2, \hat{\sigma}_{Q_4}^2)$$

$$\Lambda_V = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_S^2, \hat{\sigma}_S^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2)$$

$$\Lambda_S = \text{diag}(4, 2, 2, 1, 1, 1, 1)$$

WLS weights

Hierarchy variance scaling Λ_H : diagonal.

Series variance scaling Λ_V : elements equal within aggregation level.

Structural scaling $\Lambda_S = \text{diag}(\mathbf{S}\mathbf{1})$: elements equal to # nodes at each level.

- Depends only on seasonal period m .
- Independent of data and model.
- Allows forecasts where no errors available.

Quarterly example

$$\Lambda_H = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_{S_1}^2, \hat{\sigma}_{S_2}^2, \hat{\sigma}_{Q_1}^2, \hat{\sigma}_{Q_2}^2, \hat{\sigma}_{Q_3}^2, \hat{\sigma}_{Q_4}^2)$$

$$\Lambda_V = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_S^2, \hat{\sigma}_S^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2)$$

$$\Lambda_S = \text{diag}(4, 2, 2, 1, 1, 1, 1)$$

WLS weights

Hierarchy variance scaling Λ_H : diagonal.

Series variance scaling Λ_V : elements equal within aggregation level.

Structural scaling $\Lambda_S = \text{diag}(\mathbf{S}\mathbf{1})$: elements equal to # nodes at each level.

- Depends only on seasonal period m .
- Independent of data and model.
- Allows forecasts where no errors available.

Quarterly example

$$\Lambda_H = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_{S_1}^2, \hat{\sigma}_{S_2}^2, \hat{\sigma}_{Q_1}^2, \hat{\sigma}_{Q_2}^2, \hat{\sigma}_{Q_3}^2, \hat{\sigma}_{Q_4}^2)$$

$$\Lambda_V = \text{diag}(\hat{\sigma}_A^2, \hat{\sigma}_S^2, \hat{\sigma}_S^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2, \hat{\sigma}_Q^2)$$

$$\Lambda_S = \text{diag}(4, 2, 2, 1, 1, 1, 1)$$

Experimental setup:

- M3 forecasting competition (Makridakis and Hibon, 2000, *IJF*). In total 3003 series.
- 1,428 monthly series with a test sample of 12 observations each.
- 756 quarterly series with a test sample of 8 observations each.
- Forecast each series with ETS models.

Experimental setup:

- M3 forecasting competition (Makridakis and Hibon, 2000, *IJF*). In total 3003 series.
- 1,428 monthly series with a test sample of 12 observations each.
- 756 quarterly series with a test sample of 8 observations each.
- Forecast each series with ETS models.

Experimental setup:

- M3 forecasting competition (Makridakis and Hibon, 2000, *IJF*). In total 3003 series.
- 1,428 monthly series with a test sample of 12 observations each.
- 756 quarterly series with a test sample of 8 observations each.
- Forecast each series with ETS models.

Experimental setup:

- M3 forecasting competition (Makridakis and Hibon, 2000, *IJF*). In total 3003 series.
- 1,428 monthly series with a test sample of 12 observations each.
- 756 quarterly series with a test sample of 8 observations each.
- Forecast each series with ETS models.

Results: Monthly

MAE percent difference relative to base

	$\max h$	BU	WLS_H	WLS_V	WLS_S
Annual	1	-19.6	-22.0	-22.0	-25.1
Semi-annual	3	0.6	-4.0	-3.6	-5.4
Four-monthly	4	2.0	-2.4	-2.2	-3.0
Quarterly	6	2.4	-1.6	-1.7	-2.8
Bi-monthly	9	0.7	-2.9	-3.3	-4.3
Monthly	18	0.0	-2.2	-3.2	-3.9


Results: Quarterly

MAE percent difference relative to base


	max h	BU	WLS _H	WLS _V	WLS _S
Annual	1	-20.9	-22.7	-22.8	-22.7
Semi-annual	3	-4.5	-6.0	-6.2	-4.8
Quarterly	6	0.0	-0.2	-1.1	-0.3

UK Accidents and Emergency Demand


Annual ($k=52$)




Semi-annual ($k=26$)




Quarterly ($k=13$)




Monthly ($k=4$)



Bi-weekly ($k=2$)



Weekly ($k=1$)



--- base

— reconciled

UK Accidents and Emergency Demand

- 1 Type 1 Departments — Major A&E
- 2 Type 2 Departments — Single Specialty
- 3 Type 3 Departments — Other A&E/Minor Injury
- 4 Total Attendances
- 5 Type 1 Departments — Major A&E > 4 hrs
- 6 Type 2 Departments — Single Specialty > 4 hrs
- 7 Type 3 Departments — Other A&E/Minor Injury > 4 hrs
- 8 Total Attendances > 4 hrs
- 9 Emergency Admissions via Type 1 A&E
- 10 Total Emergency Admissions via A&E
- 11 Other Emergency Admissions (i.e., not via A&E)
- 12 Total Emergency Admissions
- 13 Number of patients spending > 4 hrs from decision to admission

UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Reconciled using WLS_V .
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	h	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- **Base forecasts using `auto.arima()`.**
- Reconciled using WLS_V .
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	h	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Reconciled using WLS_V .
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	h	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Reconciled using WLS_V .
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	h	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Reconciled using WLS_V .
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	h	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Reconciled using WLS_V .
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	h	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

Outline

- 1 Hierarchical and grouped time series
- 2 BLUF: Best Linear Unbiased Forecasts
- 3 Application: Australian tourism
- 4 Temporal hierarchies
- 5 References**

References



RJ Hyndman, RA Ahmed, G Athanasopoulos, and HL Shang (2011). “Optimal combination forecasts for hierarchical time series”. *Computational statistics & data analysis* **55**(9), 2579–2589.



RJ Hyndman, AJ Lee, and E Wang (2014). *Fast computation of reconciled forecasts for hierarchical and grouped time series*. Working paper 17/14. Monash University



SL Wickramasuriya, G Athanasopoulos, and RJ Hyndman (2015). *Forecasting hierarchical and grouped time series through trace minimization*. Working paper 15/15. Monash University



G Athanasopoulos, RJ Hyndman, N Kourentzes, and F Petropoulos (2015). *Forecasting with temporal hierarchies*. Working paper. Monash University



RJ Hyndman, AJ Lee, and E Wang (2015). *hts: Hierarchical and grouped time series*. R package v4.5 on CRAN.