



Rob J Hyndman

# Forecasting: Principles and Practice



## 7. Non-seasonal ARIMA models

[OTexts.com/fpp/8/](https://otexts.com/fpp/8/)

# Outline

- 1 Autoregressive models**
- 2 Moving average models
- 3 Non-seasonal ARIMA models
- 4 Estimation and order selection
- 5 ARIMA modelling in R
- 6 Forecasting

# Autoregressive models

## Autoregressive (AR) models:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + e_t,$$

where  $e_t$  is white noise. This is a multiple regression with **lagged values** of  $y_t$  as predictors.

# Autoregressive models

## Autoregressive (AR) models:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + e_t,$$

where  $e_t$  is white noise. This is a multiple regression with **lagged values** of  $y_t$  as predictors.


# Autoregressive models

## Autoregressive (AR) models:


$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + e_t,$$

where  $e_t$  is white noise. This is a multiple regression with **lagged values** of  $y_t$  as predictors.

AR(1)



AR(2)



# AR(1) model

$$y_t = 2 - 0.8y_{t-1} + e_t$$

$$e_t \sim N(0, 1)$$


$$T = 100.$$

# AR(1) model

$$y_t = 2 - 0.8y_{t-1} + e_t$$

$$e_t \sim N(0, 1)$$

$T = 100.$



# AR(1) model

$$y_t = c + \phi_1 y_{t-1} + e_t$$

- When  $\phi_1 = 0$ ,  $y_t$  is **equivalent to WN**
- When  $\phi_1 = 1$  and  $c = 0$ ,  $y_t$  is **equivalent to a RW**
- When  $\phi_1 = 1$  and  $c \neq 0$ ,  $y_t$  is **equivalent to a RW with drift**
- When  $\phi_1 < 0$ ,  $y_t$  tends to **oscillate between positive and negative values.**



# AR(1) model

$$y_t = c + \phi_1 y_{t-1} + e_t$$

- When  $\phi_1 = 0$ ,  $y_t$  is **equivalent to WN**
- When  $\phi_1 = 1$  and  $c = 0$ ,  $y_t$  is **equivalent to a RW**
- When  $\phi_1 = 1$  and  $c \neq 0$ ,  $y_t$  is **equivalent to a RW with drift**
- When  $\phi_1 < 0$ ,  $y_t$  tends to **oscillate between positive and negative values.**

# AR(1) model

$$y_t = c + \phi_1 y_{t-1} + e_t$$

- When  $\phi_1 = 0$ ,  $y_t$  is **equivalent to WN**
- When  $\phi_1 = 1$  and  $c = 0$ ,  $y_t$  is **equivalent to a RW**
- When  $\phi_1 = 1$  and  $c \neq 0$ ,  $y_t$  is **equivalent to a RW with drift**
- When  $\phi_1 < 0$ ,  $y_t$  tends to **oscillate between positive and negative values**.

# AR(1) model

$$y_t = c + \phi_1 y_{t-1} + e_t$$

- When  $\phi_1 = 0$ ,  $y_t$  is **equivalent to WN**
- When  $\phi_1 = 1$  and  $c = 0$ ,  $y_t$  is **equivalent to a RW**
- When  $\phi_1 = 1$  and  $c \neq 0$ ,  $y_t$  is **equivalent to a RW with drift**
- When  $\phi_1 < 0$ ,  $y_t$  tends to **oscillate between positive and negative values.**

# AR(2) model

$$y_t = 8 + 1.3y_{t-1} - 0.7y_{t-2} + e_t$$


$$e_t \sim N(0, 1)$$

$$T = 100.$$

# AR(2) model

$$y_t = 8 + 1.3y_{t-1} - 0.7y_{t-2} + e_t$$

$$e_t \sim N(0, 1)$$
$$T = 100.$$



# Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

## General condition for stationarity

Complex roots of  $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  lie outside the unit circle on the complex plane.

■ For  $p = 1$ :  $-1 < \phi_1 < 1$ .

■ For  $p = 2$ :

$$-1 < \phi_2 < 1 \quad \phi_2 + \phi_1 < 1 \quad \phi_2 - \phi_1 < 1.$$

■ More complicated conditions hold for  $p \geq 3$ .

# Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

## General condition for stationarity

Complex roots of  $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  lie outside the unit circle on the complex plane.

- For  $p = 1$ :  $-1 < \phi_1 < 1$ .
- For  $p = 2$ :  
 $-1 < \phi_2 < 1$        $\phi_2 + \phi_1 < 1$        $\phi_2 - \phi_1 < 1$ .
- More complicated conditions hold for  $p \geq 3$ .
- Estimation software takes care of this.

# Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

## General condition for stationarity

Complex roots of  $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  lie outside the unit circle on the complex plane.

- For  $p = 1$ :  $-1 < \phi_1 < 1$ .
- For  $p = 2$ :  
 $-1 < \phi_2 < 1$        $\phi_2 + \phi_1 < 1$        $\phi_2 - \phi_1 < 1$ .
- More complicated conditions hold for  $p \geq 3$ .
- Estimation software takes care of this.



# Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

## General condition for stationarity

Complex roots of  $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  lie outside the unit circle on the complex plane.

- For  $p = 1$ :  $-1 < \phi_1 < 1$ .
- For  $p = 2$ :  
 $-1 < \phi_2 < 1$        $\phi_2 + \phi_1 < 1$        $\phi_2 - \phi_1 < 1$ .
- More complicated conditions hold for  $p \geq 3$ .
- Estimation software takes care of this.

# Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

## General condition for stationarity

Complex roots of  $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$  lie outside the unit circle on the complex plane.

- For  $p = 1$ :  $-1 < \phi_1 < 1$ .
- For  $p = 2$ :  
 $-1 < \phi_2 < 1$        $\phi_2 + \phi_1 < 1$        $\phi_2 - \phi_1 < 1$ .
- More complicated conditions hold for  $p \geq 3$ .
- Estimation software takes care of this.

# Outline

- 1 Autoregressive models
- 2 Moving average models**
- 3 Non-seasonal ARIMA models
- 4 Estimation and order selection
- 5 ARIMA modelling in R
- 6 Forecasting

# Moving Average (MA) models

## Moving Average (MA) models:

$$y_t = c + e_t + \theta_1 e_{t-1} + \theta_2 e_{t-2} + \cdots + \theta_q e_{t-q},$$

where  $e_t$  is white noise. This is a multiple regression with **past errors** as predictors. *Don't confuse this with moving average smoothing!*

# Moving Average (MA) models

## Moving Average (MA) models:

$$y_t = c + e_t + \theta_1 e_{t-1} + \theta_2 e_{t-2} + \cdots + \theta_q e_{t-q},$$



where  $e_t$  is white noise. This is a multiple regression with **past errors** as predictors. *Don't confuse this with moving average smoothing!*

# Moving Average (MA) models

## Moving Average (MA) models:

$$y_t = c + e_t + \theta_1 e_{t-1} + \theta_2 e_{t-2} + \cdots + \theta_q e_{t-q},$$

where  $e_t$  is white noise. This is a multiple regression with **past errors** as predictors. *Don't confuse this with moving average smoothing!*



# MA(1) model

$$y_t = 20 + e_t + 0.8e_{t-1}$$

$$e_t \sim N(0, 1)$$


$$T = 100.$$

# MA(1) model

$$y_t = 20 + e_t + 0.8e_{t-1}$$

$$e_t \sim N(0, 1)$$

$T = 100.$





# MA(2) model

$$y_t = e_t - e_{t-1} + 0.8e_{t-2}$$


$$e_t \sim N(0, 1)$$

$$T = 100.$$

# MA(2) model

$$y_t = e_t - e_{t-1} + 0.8e_{t-2}$$

$e_t \sim N(0, 1)$   
 $T = 100.$



# Invertibility

- Any  $MA(q)$  process can be written as an  $AR(\infty)$  process if we impose some constraints on the MA parameters.
- Then the MA model is called “invertible”.
- Invertible models have some mathematical properties that make them easier to use in practice.
- Invertibility of an ARIMA model is equivalent to forecastability of an ETS model.

# Invertibility

- Any  $MA(q)$  process can be written as an  $AR(\infty)$  process if we impose some constraints on the MA parameters.
- Then the MA model is called “invertible”.
- Invertible models have some mathematical properties that make them easier to use in practice.
- Invertibility of an ARIMA model is equivalent to forecastability of an ETS model.

# Invertibility

- Any  $MA(q)$  process can be written as an  $AR(\infty)$  process if we impose some constraints on the MA parameters.
- Then the MA model is called “invertible”.
- Invertible models have some mathematical properties that make them easier to use in practice.
- Invertibility of an ARIMA model is equivalent to forecastability of an ETS model.

# Invertibility

- Any  $MA(q)$  process can be written as an  $AR(\infty)$  process if we impose some constraints on the MA parameters.
- Then the MA model is called “invertible”.
- Invertible models have some mathematical properties that make them easier to use in practice.
- Invertibility of an ARIMA model is equivalent to forecastability of an ETS model.

# Invertibility

## General condition for invertibility

Complex roots of  $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q$  lie outside the unit circle on the complex plane.

■ For  $q = 1$ :  $-1 < \theta_1 < 1$ .

■ For  $q = 2$ :

$$-1 < \theta_2 < 1 \quad \theta_2 + \theta_1 > -1 \quad \theta_1 - \theta_2 < 1.$$

■ More complicated conditions hold for  $q \geq 3$ .

■ For  $q \geq 3$ ,  $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q = 0$

# Invertibility

## General condition for invertibility

Complex roots of  $1 + \theta_1 z + \theta_2 z^2 + \cdots + \theta_q z^q$  lie outside the unit circle on the complex plane.

- For  $q = 1$ :  $-1 < \theta_1 < 1$ .

- For  $q = 2$ :

$$-1 < \theta_2 < 1 \quad \theta_2 + \theta_1 > -1 \quad \theta_1 - \theta_2 < 1.$$

- More complicated conditions hold for  $q \geq 3$ .

- Estimation software takes care of this.



# Invertibility

## General condition for invertibility

Complex roots of  $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q$  lie outside the unit circle on the complex plane.

- For  $q = 1$ :  $-1 < \theta_1 < 1$ .
- For  $q = 2$ :  
 $-1 < \theta_2 < 1$        $\theta_2 + \theta_1 > -1$        $\theta_1 - \theta_2 < 1$ .
- More complicated conditions hold for  $q \geq 3$ .
- Estimation software takes care of this.

# Invertibility

## General condition for invertibility

Complex roots of  $1 + \theta_1 z + \theta_2 z^2 + \cdots + \theta_q z^q$  lie outside the unit circle on the complex plane.

- For  $q = 1$ :  $-1 < \theta_1 < 1$ .
- For  $q = 2$ :  
 $-1 < \theta_2 < 1 \quad \theta_2 + \theta_1 > -1 \quad \theta_1 - \theta_2 < 1$ .
- More complicated conditions hold for  $q \geq 3$ .
- Estimation software takes care of this.

# Invertibility

## General condition for invertibility

Complex roots of  $1 + \theta_1 z + \theta_2 z^2 + \cdots + \theta_q z^q$  lie outside the unit circle on the complex plane.

- For  $q = 1$ :  $-1 < \theta_1 < 1$ .
- For  $q = 2$ :  
 $-1 < \theta_2 < 1 \quad \theta_2 + \theta_1 > -1 \quad \theta_1 - \theta_2 < 1$ .
- More complicated conditions hold for  $q \geq 3$ .
- Estimation software takes care of this.

# Outline

- 1 Autoregressive models
- 2 Moving average models
- 3 Non-seasonal ARIMA models**
- 4 Estimation and order selection
- 5 ARIMA modelling in R
- 6 Forecasting

# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$**  and **lagged errors**.
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$  and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$  and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$  and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

- Combine ARMA model with differencing.



# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$  and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

- Combine ARMA model with **differencing.**
- $(1 - B)^d y_t$  follows an ARMA model.

# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$  and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

- Combine ARMA model with **differencing.**
- $(1 - B)^d y_t$  follows an ARMA model.

# ARIMA models

## Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} \\ + \theta_1 e_{t-1} + \cdots + \theta_q e_{t-q} + e_t.$$

- Predictors include both **lagged values of  $y_t$  and lagged errors.**
- Conditions on coefficients ensure stationarity.
- Conditions on coefficients ensure invertibility.

## Autoregressive Integrated Moving Average models

- Combine ARMA model with **differencing.**
- $(1 - B)^d y_t$  follows an ARMA model.

# ARIMA models

## Autoregressive Integrated Moving Average models

### ARIMA( $p, d, q$ ) model

AR:  $p$  = order of the autoregressive part  
I:  $d$  = degree of first differencing involved  
MA:  $q$  = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- AR( $p$ ): ARIMA( $p,0,0$ )
- MA( $q$ ): ARIMA(0,0, $q$ )

# ARIMA models

## Autoregressive Integrated Moving Average models

### ARIMA( $p, d, q$ ) model

AR:  $p$  = order of the autoregressive part  
I:  $d$  = degree of first differencing involved  
MA:  $q$  = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- AR( $p$ ): ARIMA( $p,0,0$ )
- MA( $q$ ): ARIMA(0,0, $q$ )

# ARIMA models

## Autoregressive Integrated Moving Average models

### ARIMA( $p, d, q$ ) model

AR:  $p$  = order of the autoregressive part  
I:  $d$  = degree of first differencing involved  
MA:  $q$  = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- AR( $p$ ): ARIMA( $p,0,0$ )
- MA( $q$ ): ARIMA(0,0, $q$ )

# ARIMA models

## Autoregressive Integrated Moving Average models

### ARIMA( $p, d, q$ ) model

AR:  $p$  = order of the autoregressive part  
I:  $d$  = degree of first differencing involved  
MA:  $q$  = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- $AR(p)$ : ARIMA( $p,0,0$ )
- $MA(q)$ : ARIMA(0,0, $q$ )

# ARIMA models

## Autoregressive Integrated Moving Average models

### ARIMA( $p, d, q$ ) model

AR:  $p$  = order of the autoregressive part  
I:  $d$  = degree of first differencing involved  
MA:  $q$  = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- AR( $p$ ): ARIMA( $p,0,0$ )
- MA( $q$ ): ARIMA(0,0, $q$ )



# Backshift notation for ARIMA

## ■ ARMA model:

$$y_t = c + \phi_1 B y_t + \cdots + \phi_p B^p y_t + e_t + \theta_1 B e_t + \cdots + \theta_q B^q e_t$$

or  $(1 - \phi_1 B - \cdots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \cdots + \theta_q B^q) e_t$

## ■ ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + (1 + \theta_1 B) e_t$$

# Backshift notation for ARIMA

## ■ ARMA model:

$$y_t = c + \phi_1 B y_t + \cdots + \phi_p B^p y_t + e_t + \theta_1 B e_t + \cdots + \theta_q B^q e_t$$

or  $(1 - \phi_1 B - \cdots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \cdots + \theta_q B^q) e_t$

## ■ ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + (1 + \theta_1 B) e_t$$

# Backshift notation for ARIMA

## ■ ARMA model:

$$y_t = c + \phi_1 B y_t + \cdots + \phi_p B^p y_t + e_t + \theta_1 B e_t + \cdots + \theta_q B^q e_t$$

or  $(1 - \phi_1 B - \cdots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \cdots + \theta_q B^q) e_t$

## ■ ARIMA(1,1,1) model:

$$(1 - \phi_1 B) \underset{\substack{\uparrow \\ \text{First} \\ \text{difference}}}{(1 - B)} y_t = c + (1 + \theta_1 B) e_t$$

# Backshift notation for ARIMA

## ■ ARMA model:

$$y_t = c + \phi_1 B y_t + \cdots + \phi_p B^p y_t + e_t + \theta_1 B e_t + \cdots + \theta_q B^q e_t$$

or  $(1 - \phi_1 B - \cdots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \cdots + \theta_q B^q) e_t$

## ■ ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + (1 + \theta_1 B) e_t$$

↑  
AR(1)

# Backshift notation for ARIMA

## ■ ARMA model:

$$y_t = c + \phi_1 B y_t + \cdots + \phi_p B^p y_t + e_t + \theta_1 B e_t + \cdots + \theta_q B^q e_t$$

or  $(1 - \phi_1 B - \cdots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \cdots + \theta_q B^q) e_t$

## ■ ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + \underset{\substack{\uparrow \\ \text{MA}(1)}}{(1 + \theta_1 B)} e_t$$

# Backshift notation for ARIMA

## ■ ARMA model:

$$y_t = c + \phi_1 B y_t + \cdots + \phi_p B^p y_t + e_t + \theta_1 B e_t + \cdots + \theta_q B^q e_t$$

or  $(1 - \phi_1 B - \cdots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \cdots + \theta_q B^q) e_t$


## ■ ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + (1 + \theta_1 B) e_t$$

Written out:

$$y_t = c + y_{t-1} + \phi_1 y_{t-1} - \phi_1 y_{t-2} + \theta_1 e_{t-1} + e_t$$

# US personal consumption



# US personal consumption

```
> fit <- auto.arima(usconsumption[,1],  
  max.P=0,max.Q=0,D=0)
```

ARIMA(0,0,3) with non-zero mean

Coefficients:

|      | ma1    | ma2    | ma3    | intercept |
|------|--------|--------|--------|-----------|
|      | 0.2542 | 0.2260 | 0.2695 | 0.7562    |
| s.e. | 0.0767 | 0.0779 | 0.0692 | 0.0844    |

sigma^2 estimated as 0.3856: log likelihood=-154.73  
AIC=319.46 AICc=319.84 BIC=334.96



# US personal consumption

```
> fit <- auto.arima(usconsumption[,1],  
  max.P=0,max.Q=0,D=0)
```

ARIMA(0,0,3) with non-zero mean

Coefficients:

|      | ma1    | ma2    | ma3    | intercept |
|------|--------|--------|--------|-----------|
|      | 0.2542 | 0.2260 | 0.2695 | 0.7562    |
| s.e. | 0.0767 | 0.0779 | 0.0692 | 0.0844    |

sigma^2 estimated as 0.3856: log likelihood=-154.73  
AIC=319.46 AICc=319.84 BIC=334.96


**ARIMA(0,0,3) or MA(3) model:**

$$y_t = 0.756 + e_t + 0.254e_{t-1} + 0.226e_{t-2} + 0.269e_{t-3},$$

where  $e_t$  is white noise with standard deviation  $0.62 = \sqrt{0.3856}$ .


# US personal consumption

Forecasts from ARIMA(0,0,3) with non-zero mean



# US personal consumption

Forecasts from ARIMA(0,0,3) with non-zero mean



`plot(forecast(fit,h=10),include=80)`

# Understanding ARIMA models

- If  $c = 0$  and  $d = 0$ , the long-term forecasts will go to zero.
- If  $c = 0$  and  $d = 1$ , the long-term forecasts will go to a non-zero constant.
- If  $c = 0$  and  $d = 2$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 0$ , the long-term forecasts will go to the mean of the data.
- If  $c \neq 0$  and  $d = 1$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 2$ , the long-term forecasts will follow a quadratic trend.

# Understanding ARIMA models

- If  $c = 0$  and  $d = 0$ , the long-term forecasts will go to zero.
- If  $c = 0$  and  $d = 1$ , the long-term forecasts will go to a non-zero constant.
- If  $c = 0$  and  $d = 2$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 0$ , the long-term forecasts will go to the mean of the data.
- If  $c \neq 0$  and  $d = 1$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 2$ , the long-term forecasts will follow a quadratic trend.

# Understanding ARIMA models

- If  $c = 0$  and  $d = 0$ , the long-term forecasts will go to zero.
- If  $c = 0$  and  $d = 1$ , the long-term forecasts will go to a non-zero constant.
- If  $c = 0$  and  $d = 2$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 0$ , the long-term forecasts will go to the mean of the data.
- If  $c \neq 0$  and  $d = 1$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 2$ , the long-term forecasts will follow a quadratic trend.

# Understanding ARIMA models

- If  $c = 0$  and  $d = 0$ , the long-term forecasts will go to zero.
- If  $c = 0$  and  $d = 1$ , the long-term forecasts will go to a non-zero constant.
- If  $c = 0$  and  $d = 2$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 0$ , the long-term forecasts will go to the mean of the data.
- If  $c \neq 0$  and  $d = 1$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 2$ , the long-term forecasts will follow a quadratic trend.

# Understanding ARIMA models

- If  $c = 0$  and  $d = 0$ , the long-term forecasts will go to zero.
- If  $c = 0$  and  $d = 1$ , the long-term forecasts will go to a non-zero constant.
- If  $c = 0$  and  $d = 2$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 0$ , the long-term forecasts will go to the mean of the data.
- If  $c \neq 0$  and  $d = 1$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 2$ , the long-term forecasts will follow a quadratic trend.



# Understanding ARIMA models

- If  $c = 0$  and  $d = 0$ , the long-term forecasts will go to zero.
- If  $c = 0$  and  $d = 1$ , the long-term forecasts will go to a non-zero constant.
- If  $c = 0$  and  $d = 2$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 0$ , the long-term forecasts will go to the mean of the data.
- If  $c \neq 0$  and  $d = 1$ , the long-term forecasts will follow a straight line.
- If  $c \neq 0$  and  $d = 2$ , the long-term forecasts will follow a quadratic trend.

# Understanding ARIMA models

## Forecast variance and $d$

- The higher the value of  $d$ , the more rapidly the prediction intervals increase in size.
- For  $d = 0$ , the long-term forecast standard deviation will go to the standard deviation of the historical data.

## Cyclic behaviour

For cyclic behaviour, the model must be able to capture the periodicity of the data. This is achieved by including a seasonal component in the model.

# Understanding ARIMA models

## Forecast variance and $d$

- The higher the value of  $d$ , the more rapidly the prediction intervals increase in size.
- For  $d = 0$ , the long-term forecast standard deviation will go to the standard deviation of the historical data.

## Cyclic behaviour

- For cyclic forecasts,  $p > 2$  and some restrictions on coefficients are required.
- The model must be able to capture the cyclic pattern.

# Understanding ARIMA models

## Forecast variance and $d$

- The higher the value of  $d$ , the more rapidly the prediction intervals increase in size.
- For  $d = 0$ , the long-term forecast standard deviation will go to the standard deviation of the historical data.

## Cyclic behaviour

- For cyclic forecasts,  $p > 2$  and some restrictions on coefficients are required.
- If  $p = 2$ , we need  $\phi_1^2 + 4\phi_2 < 0$ . Then average cycle of length

$$(2\pi) / [\arccos(-\phi_1(1 - \phi_2)/(4\phi_2))].$$

# Understanding ARIMA models

## Forecast variance and $d$

- The higher the value of  $d$ , the more rapidly the prediction intervals increase in size.
- For  $d = 0$ , the long-term forecast standard deviation will go to the standard deviation of the historical data.

## Cyclic behaviour

- For cyclic forecasts,  $p > 2$  and some restrictions on coefficients are required.
- If  $p = 2$ , we need  $\phi_1^2 + 4\phi_2 < 0$ . Then average cycle of length

$$(2\pi) / [\arccos(-\phi_1(1 - \phi_2)/(4\phi_2))].$$

# Understanding ARIMA models

## Forecast variance and $d$

- The higher the value of  $d$ , the more rapidly the prediction intervals increase in size.
- For  $d = 0$ , the long-term forecast standard deviation will go to the standard deviation of the historical data.

## Cyclic behaviour

- For cyclic forecasts,  $p > 2$  and some restrictions on coefficients are required.
- If  $p = 2$ , we need  $\phi_1^2 + 4\phi_2 < 0$ . Then average cycle of length

$$(2\pi) / [\arccos(-\phi_1(1 - \phi_2)/(4\phi_2))].$$

# Partial autocorrelations

**Partial autocorrelations** measure relationship between  $y_t$  and  $y_{t-k}$ , when the effects of other time lags —  $1, 2, 3, \dots, k-1$  — are removed.

$\alpha_k$  =  $k$ th partial autocorrelation coefficient  
= equal to the estimate of  $b_k$  in regression:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives  $\alpha_k$  for different values of  $k$ .

# Partial autocorrelations

**Partial autocorrelations** measure relationship between  $y_t$  and  $y_{t-k}$ , when the effects of other time lags —  $1, 2, 3, \dots, k-1$  — are removed.

$\alpha_k$  =  $k$ th partial autocorrelation coefficient  
= equal to the estimate of  $b_k$  in regression:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives  $\alpha_k$  for different values of  $k$ .
- There are more efficient ways of calculating  $\alpha_k$ .

$$\alpha_1 = \rho_1$$

$$\alpha_2 = \rho_2 - \rho_1^2$$



# Partial autocorrelations

**Partial autocorrelations** measure relationship between  $y_t$  and  $y_{t-k}$ , when the effects of other time lags —  $1, 2, 3, \dots, k-1$  — are removed.

$\alpha_k$  =  $k$ th partial autocorrelation coefficient  
= equal to the estimate of  $b_k$  in regression:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives  $\alpha_k$  for different values of  $k$ .
- There are more efficient ways of calculating  $\alpha_k$ .
- $\alpha_1 = \rho_1$
- same critical values of  $\pm 1.96/\sqrt{T}$  as for ACF.

# Partial autocorrelations

**Partial autocorrelations** measure relationship between  $y_t$  and  $y_{t-k}$ , when the effects of other time lags —  $1, 2, 3, \dots, k-1$  — are removed.

$\alpha_k$  =  $k$ th partial autocorrelation coefficient  
= equal to the estimate of  $b_k$  in regression:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives  $\alpha_k$  for different values of  $k$ .
- There are more efficient ways of calculating  $\alpha_k$ .
- $\alpha_1 = \rho_1$
- same critical values of  $\pm 1.96/\sqrt{T}$  as for ACF.

# Partial autocorrelations

**Partial autocorrelations** measure relationship between  $y_t$  and  $y_{t-k}$ , when the effects of other time lags —  $1, 2, 3, \dots, k-1$  — are removed.

$\alpha_k$  =  $k$ th partial autocorrelation coefficient  
= equal to the estimate of  $b_k$  in regression:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives  $\alpha_k$  for different values of  $k$ .
- There are more efficient ways of calculating  $\alpha_k$ .
- $\alpha_1 = \rho_1$
- same critical values of  $\pm 1.96/\sqrt{T}$  as for ACF.

# Partial autocorrelations


**Partial autocorrelations** measure relationship between  $y_t$  and  $y_{t-k}$ , when the effects of other time lags —  $1, 2, 3, \dots, k-1$  — are removed.

$\alpha_k$  =  $k$ th partial autocorrelation coefficient  
= equal to the estimate of  $b_k$  in regression:



$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives  $\alpha_k$  for different values of  $k$ .
- There are more efficient ways of calculating  $\alpha_k$ .
- $\alpha_1 = \rho_1$
- same critical values of  $\pm 1.96/\sqrt{T}$  as for ACF.

# Example: US consumption



# Example: US consumption



# ACF and PACF interpretation

**ARIMA( $p,d,0$ )** model if ACF and PACF plots of differenced data show:

- the ACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $p$  in PACF, but none beyond lag  $p$ .

**ARIMA( $0,d,q$ )** model if ACF and PACF plots of differenced data show:

- the PACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $q$  in ACF, but none beyond lag  $q$ .

# ACF and PACF interpretation

**ARIMA( $p,d,0$ )** model if ACF and PACF plots of differenced data show:

- the ACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $p$  in PACF, but none beyond lag  $p$ .

**ARIMA( $0,d,q$ )** model if ACF and PACF plots of differenced data show:

- the PACF is exponentially decaying or sinusoidal;
- the ACF has a significant spike at lag  $q$ , but none beyond lag  $q$ .



# ACF and PACF interpretation

**ARIMA( $p,d,0$ )** model if ACF and PACF plots of differenced data show:

- the ACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $p$  in PACF, but none beyond lag  $p$ .

**ARIMA( $0,d,q$ )** model if ACF and PACF plots of differenced data show:

- the PACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $q$  in ACF, but none beyond lag  $q$ .

# ACF and PACF interpretation

**ARIMA( $p, d, 0$ )** model if ACF and PACF plots of differenced data show:

- the ACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $p$  in PACF, but none beyond lag  $p$ .

**ARIMA( $0, d, q$ )** model if ACF and PACF plots of differenced data show:

- the PACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $q$  in ACF, but none beyond lag  $q$ .

# ACF and PACF interpretation

**ARIMA( $p, d, 0$ )** model if ACF and PACF plots of differenced data show:


- the ACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $p$  in PACF, but none beyond lag  $p$ .

**ARIMA( $0, d, q$ )** model if ACF and PACF plots of differenced data show:


- the PACF is exponentially decaying or sinusoidal;
- there is a significant spike at lag  $q$  in ACF, but none beyond lag  $q$ .

# Example: Mink trapping


Annual number of minks trapped



# Example: Mink trapping



# Example: Mink trapping



# Outline

- 1 Autoregressive models
- 2 Moving average models
- 3 Non-seasonal ARIMA models
- 4 Estimation and order selection**
- 5 ARIMA modelling in R
- 6 Forecasting

# Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters  $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$ .

- MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2.$$

- The `Arima()` command allows CLS or MLE estimation.
- Non-linear optimization must be used in either case.
- Different software will give different estimates



# Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters  $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$ .

- MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2.$$

- The `Arima()` command allows CLS or MLE estimation.
- Non-linear optimization must be used in either case.
- Different software will give different estimates.

# Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters  $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$ .

- MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2.$$

- The `Arima()` command allows CLS or MLE estimation.
- Non-linear optimization must be used in either case.
- Different software will give different estimates.

# Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters  $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$ .

- MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2.$$

- The `Arima()` command allows CLS or MLE estimation.
- Non-linear optimization must be used in either case.
- Different software will give different estimates.

# Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters  $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$ .

- MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2.$$

- The `Arima()` command allows CLS or MLE estimation.
- Non-linear optimization must be used in either case.
- Different software will give different estimates.

# Information criteria

## Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where  $L$  is the likelihood of the data,

$k = 1$  if  $c \neq 0$  and  $k = 0$  if  $c = 0$ .

## Corrected AIC:

$$\text{AIC}_c = \text{AIC} + \frac{2(p + q + k + 1)(p + q + k + 2)}{T - p - q - k - 2}.$$

## Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + \log(T)(p + q + k - 1).$$

Good models are obtained by minimizing either the AIC,  $\text{AIC}_c$  or BIC. Our preference is to use the  $\text{AIC}_c$ .

# Information criteria

## Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where  $L$  is the likelihood of the data,

$k = 1$  if  $c \neq 0$  and  $k = 0$  if  $c = 0$ .

## Corrected AIC:

$$\text{AIC}_c = \text{AIC} + \frac{2(p + q + k + 1)(p + q + k + 2)}{T - p - q - k - 2}.$$

## Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + \log(T)(p + q + k - 1).$$

Good models are obtained by minimizing either the AIC,  $\text{AIC}_c$  or BIC. Our preference is to use the  $\text{AIC}_c$ .

# Information criteria

## Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where  $L$  is the likelihood of the data,

$k = 1$  if  $c \neq 0$  and  $k = 0$  if  $c = 0$ .

## Corrected AIC:

$$\text{AIC}_c = \text{AIC} + \frac{2(p + q + k + 1)(p + q + k + 2)}{T - p - q - k - 2}.$$

## Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + \log(T)(p + q + k - 1).$$

Good models are obtained by minimizing either the AIC,  $\text{AIC}_c$  or BIC. Our preference is to use the  $\text{AIC}_c$ .

# Information criteria

## Akaike's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where  $L$  is the likelihood of the data,

$k = 1$  if  $c \neq 0$  and  $k = 0$  if  $c = 0$ .

## Corrected AIC:

$$\text{AIC}_c = \text{AIC} + \frac{2(p + q + k + 1)(p + q + k + 2)}{T - p - q - k - 2}.$$

## Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + \log(T)(p + q + k - 1).$$

Good models are obtained by minimizing either the AIC,  $\text{AIC}_c$  or BIC. Our preference is to use the  $\text{AIC}_c$ .



# Outline

- 1 Autoregressive models
- 2 Moving average models
- 3 Non-seasonal ARIMA models
- 4 Estimation and order selection
- 5 ARIMA modelling in R**
- 6 Forecasting

# How does auto.arima() work?

## A non-seasonal ARIMA process

$$\phi(B)(1 - B)^d y_t = c + \theta(B)\varepsilon_t$$

Need to select appropriate orders:  $p, q, d$

# How does auto.arima() work?

## A non-seasonal ARIMA process

$$\phi(B)(1 - B)^d y_t = c + \theta(B)\varepsilon_t$$

Need to select appropriate orders:  $p, q, d$

## Hyndman and Khandakar (JSS, 2008) algorithm:

- Select no. differences  $d$  and  $D$  via unit root tests.
- Select  $p, q$  by minimising AICc.
- Use stepwise search to traverse model space.

# How does auto.arima() work?

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1)$$

where  $L$  is the maximised likelihood fitted to the *differenced* data,  $k = 1$  if  $c \neq 0$  and  $k = 0$  otherwise.

**Step 1:** Select current model (with smallest AIC) from:

ARIMA(2,  $d$ , 2)

ARIMA(0,  $d$ , 0)

ARIMA(1,  $d$ , 0)

ARIMA(0,  $d$ , 1)

**Step 2:** Consider variations of current model:

- vary one of  $p, q$ , from current model by  $\pm 1$
- $p, q$  both vary from current model by  $\pm 1$
- Include/exclude  $c$  from current model

Model with lowest AIC becomes current model.

Repeat Step 2 until no lower AIC can be found.

# How does auto.arima() work?

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1)$$

where  $L$  is the maximised likelihood fitted to the *differenced* data,  $k = 1$  if  $c \neq 0$  and  $k = 0$  otherwise.

**Step 1:** Select current model (with smallest AIC) from:

ARIMA(2,  $d$ , 2)

ARIMA(0,  $d$ , 0)

ARIMA(1,  $d$ , 0)

ARIMA(0,  $d$ , 1)

**Step 2:** Consider variations of current model:

- vary one of  $p, q$ , from current model by  $\pm 1$
- $p, q$  both vary from current model by  $\pm 1$
- Include/exclude  $c$  from current model

Model with lowest AIC becomes current model.

**Repeat Step 2 until no lower AIC can be found.**

# How does auto.arima() work?

$$AIC = -2 \log(L) + 2(p + q + k + 1)$$

where  $L$  is the maximised likelihood fitted to the *differenced* data,  $k = 1$  if  $c \neq 0$  and  $k = 0$  otherwise.

**Step 1:** Select current model (with smallest AIC) from:

ARIMA(2,  $d$ , 2)

ARIMA(0,  $d$ , 0)

ARIMA(1,  $d$ , 0)

ARIMA(0,  $d$ , 1)

**Step 2:** Consider variations of current model:

- vary one of  $p, q$ , from current model by  $\pm 1$
- $p, q$  both vary from current model by  $\pm 1$
- Include/exclude  $c$  from current model

Model with lowest AIC becomes current model.

Repeat Step 2 until no lower AIC can be found.

# How does auto.arima() work?

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1)$$

where  $L$  is the maximised likelihood fitted to the *differenced* data,  $k = 1$  if  $c \neq 0$  and  $k = 0$  otherwise.

**Step 1:** Select current model (with smallest AIC) from:

ARIMA(2,  $d$ , 2)

ARIMA(0,  $d$ , 0)

ARIMA(1,  $d$ , 0)

ARIMA(0,  $d$ , 1)

**Step 2:** Consider variations of current model:

- vary one of  $p, q$ , from current model by  $\pm 1$
- $p, q$  both vary from current model by  $\pm 1$
- Include/exclude  $c$  from current model

Model with lowest AIC becomes current model.

**Repeat Step 2 until no lower AIC can be found.**

# Choosing your own model

```
tsdisplay(internet)
adf.test(internet)
kpss.test(internet)
kpss.test(diff(internet))
tsdisplay(diff(internet))
fit <- Arima(internet,order=c(3,1,0))
fit2 <- auto.arima(internet)
Acf(residuals(fit))
Box.test(residuals(fit), fitdf=3, lag=10,
         type="Ljung")
tsdiag(fit)
forecast(fit)
plot(forecast(fit))
```



# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an  $AR(p)$  or  $MA(q)$  model appropriate?
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure


- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data.
- 4 The automated algorithm only takes care of steps 3–5. So even if you use it, you will still need to take care of the other steps yourself.
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.




# Modelling procedure

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of  $t$ . The automated algorithm only takes care of
- 4 Steps 3–5. So even if you use it, you will still need to take care of the other steps yourself.
- 5 Try your chosen model(s), and use the  $AIC_c$  to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

# Modelling procedure



# Seasonally adjusted electrical equipment



# Seasonally adjusted electrical equipment

- 1 Time plot shows sudden changes, particularly big drop in 2008/2009 due to global economic environment. Otherwise nothing unusual and no need for data adjustments.
- 2 No evidence of changing variance, so no Box-Cox transformation.
- 3 Data are clearly non-stationary, so we take first differences.


# Seasonally adjusted electrical equipment

- 1 Time plot shows sudden changes, particularly big drop in 2008/2009 due to global economic environment. Otherwise nothing unusual and no need for data adjustments.
- 2 No evidence of changing variance, so no Box-Cox transformation.
- 3 Data are clearly non-stationary, so we take first differences.




## Seasonally adjusted electrical equipment

- 1 Time plot shows sudden changes, particularly big drop in 2008/2009 due to global economic environment. Otherwise nothing unusual and no need for data adjustments.
- 2 No evidence of changing variance, so no Box-Cox transformation.
- 3 Data are clearly non-stationary, so we take first differences.

# Seasonally adjusted electrical equipment



# Seasonally adjusted electrical equipment



> `tsdisplay(diff(eeadj))`



- 4 PACF is suggestive of AR(3). So initial candidate model is ARIMA(3,1,0). No other obvious candidates.
- 5 Fit ARIMA(3,1,0) model along with variations: ARIMA(4,1,0), ARIMA(2,1,0), ARIMA(3,1,1), etc. ARIMA(3,1,1) has smallest  $AIC_c$  value.

- 4 PACF is suggestive of AR(3). So initial candidate model is ARIMA(3,1,0). No other obvious candidates.
- 5 Fit ARIMA(3,1,0) model along with variations: ARIMA(4,1,0), ARIMA(2,1,0), ARIMA(3,1,1), etc. ARIMA(3,1,1) has smallest  $AIC_c$  value.

# Seasonally adjusted electrical equipment

```
> fit <- Arima(eeadj, order=c(3,1,1))
```

```
> summary(fit)
```

Series: eeadj

ARIMA(3,1,1)

Coefficients:

|      | ar1    | ar2    | ar3    | ma1     |
|------|--------|--------|--------|---------|
|      | 0.0519 | 0.1191 | 0.3730 | -0.4542 |
| s.e. | 0.1840 | 0.0888 | 0.0679 | 0.1993  |

sigma<sup>2</sup> estimated as 9.532: log likelihood=-484.08

AIC=978.17    AICc=978.49    BIC=994.4

- 6 ACF plot of residuals from ARIMA(3,1,1) model look like white noise.

```
Acf(residuals(fit))  
Box.test(residuals(fit), lag=24,  
         fitdf=4, type="Ljung")
```

- 6 ACF plot of residuals from ARIMA(3,1,1) model look like white noise.

```
Acf(residuals(fit))  
Box.test(residuals(fit), lag=24,  
         fitdf=4, type="Ljung")
```


- 6 ACF plot of residuals from ARIMA(3,1,1) model look like white noise.

```
Acf(residuals(fit))
```

```
Box.test(residuals(fit), lag=24,  
         fitdf=4, type="Ljung")
```


# Seasonally adjusted electrical equipment

Forecasts from ARIMA(3,1,1)



# Seasonally adjusted electrical equipment

Forecasts from ARIMA(3,1,1)



```
> plot(forecast(fit))
```



# Outline

- 1 Autoregressive models
- 2 Moving average models
- 3 Non-seasonal ARIMA models
- 4 Estimation and order selection
- 5 ARIMA modelling in R
- 6 Forecasting**

# Point forecasts

- 1 Rearrange ARIMA equation so  $y_t$  is on LHS.
- 2 Rewrite equation by replacing  $t$  by  $T + h$ .
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with  $h = 1$ . Repeat for  $h = 2, 3, \dots$

# Point forecasts

- 1 Rearrange ARIMA equation so  $y_t$  is on LHS.
- 2 Rewrite equation by replacing  $t$  by  $T + h$ .
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with  $h = 1$ . Repeat for  $h = 2, 3, \dots$

# Point forecasts

- 1 Rearrange ARIMA equation so  $y_t$  is on LHS.
- 2 Rewrite equation by replacing  $t$  by  $T + h$ .
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with  $h = 1$ . Repeat for  $h = 2, 3, \dots$

# Point forecasts

- 1 Rearrange ARIMA equation so  $y_t$  is on LHS.
- 2 Rewrite equation by replacing  $t$  by  $T + h$ .
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with  $h = 1$ . Repeat for  $h = 2, 3, \dots$

# Point forecasts

- 1 Rearrange ARIMA equation so  $y_t$  is on LHS.
- 2 Rewrite equation by replacing  $t$  by  $T + h$ .
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with  $h = 1$ . Repeat for  $h = 2, 3, \dots$

# Point forecasts

## ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)e_t,$$

$$\begin{aligned} [1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4] y_t \\ = (1 + \theta_1 B)e_t, \end{aligned}$$

$$\begin{aligned} y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} \\ + \phi_3 y_{t-4} = e_t + \theta_1 e_{t-1}. \end{aligned}$$

$$\begin{aligned} y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3 y_{t-4} + e_t + \theta_1 e_{t-1}. \end{aligned}$$

# Point forecasts

## ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)e_t,$$

$$\begin{aligned} [1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4] y_t \\ = (1 + \theta_1 B)e_t, \end{aligned}$$

$$\begin{aligned} y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} \\ + \phi_3 y_{t-4} = e_t + \theta_1 e_{t-1}. \end{aligned}$$

$$\begin{aligned} y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3 y_{t-4} + e_t + \theta_1 e_{t-1}. \end{aligned}$$



# Point forecasts

## ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)e_t,$$

$$\begin{aligned} [1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4] y_t \\ = (1 + \theta_1 B)e_t, \end{aligned}$$

$$\begin{aligned} y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} \\ + \phi_3 y_{t-4} = e_t + \theta_1 e_{t-1}. \end{aligned}$$

$$\begin{aligned} y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3 y_{t-4} + e_t + \theta_1 e_{t-1}. \end{aligned}$$

# Point forecasts

## ARIMA(3,1,1) forecasts: Step 1

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)e_t,$$

$$\begin{aligned} [1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4] y_t \\ = (1 + \theta_1 B)e_t, \end{aligned}$$

$$\begin{aligned} y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} \\ + \phi_3 y_{t-4} = e_t + \theta_1 e_{t-1}. \end{aligned}$$

$$\begin{aligned} y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3 y_{t-4} + e_t + \theta_1 e_{t-1}. \end{aligned}$$

# Point forecasts ( $h=1$ )

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + e_t + \theta_1e_{t-1}.$$

## ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + e_{T+1} + \theta_1e_T.$$

## ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+1|T} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + \theta_1\hat{e}_T.$$

# Point forecasts ( $h=1$ )

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + e_t + \theta_1e_{t-1}.$$

## ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + e_{T+1} + \theta_1e_T.$$

## ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+1|T} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + \theta_1\hat{e}_T.$$

# Point forecasts ( $h=1$ )

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + e_t + \theta_1e_{t-1}.$$

## ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + e_{T+1} + \theta_1e_T.$$

## ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+1|T} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} \\ - \phi_3y_{T-3} + \theta_1\hat{e}_T.$$

# Point forecasts ( $h=2$ )

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + e_t + \theta_1e_{t-1}.$$

## ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2} + e_{T+2} + \theta_1e_{T+1}.$$

## ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+2|T} = (1 + \phi_1)\hat{y}_{T+1|T} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2}.$$

# Point forecasts ( $h=2$ )

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + e_t + \theta_1e_{t-1}.$$

## ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2} + e_{T+2} + \theta_1e_{T+1}.$$

## ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+2|T} = (1 + \phi_1)\hat{y}_{T+1|T} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2}.$$

# Point forecasts ( $h=2$ )

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} \\ - \phi_3y_{t-4} + e_t + \theta_1e_{t-1}.$$

## ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2} + e_{T+2} + \theta_1e_{T+1}.$$

## ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+2|T} = (1 + \phi_1)\hat{y}_{T+1|T} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} \\ - \phi_3y_{T-2}.$$



# Forecast intervals

## 95% forecast interval

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where  $v_{T+h|T}$  is estimated forecast variance.

- $v_{T+1|T} = \hat{\sigma}^2$  for all ARIMA models regardless of parameters and orders.
- Multi-step forecast intervals for ARIMA(0,0,q):

$$y_t = e_t + \sum_{i=1}^q \theta_i e_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[ 1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

# Forecast intervals

## 95% forecast interval

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where  $v_{T+h|T}$  is estimated forecast variance.

- $v_{T+1|T} = \hat{\sigma}^2$  for all ARIMA models regardless of parameters and orders.
- Multi-step forecast intervals for ARIMA(0,0,q):

$$y_t = e_t + \sum_{i=1}^q \theta_i e_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[ 1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

# Forecast intervals

## 95% forecast interval

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where  $v_{T+h|T}$  is estimated forecast variance.

- $v_{T+1|T} = \hat{\sigma}^2$  for all ARIMA models regardless of parameters and orders.
- Multi-step forecast intervals for ARIMA(0,0,q):

$$y_t = e_t + \sum_{i=1}^q \theta_i e_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[ 1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

# Forecast intervals

## 95% forecast interval

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where  $v_{T+h|T}$  is estimated forecast variance.

- Multi-step forecast intervals for ARIMA(0,0,q):

$$y_t = e_t + \sum_{i=1}^q \theta_i e_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[ 1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

- AR(1): Rewrite as MA( $\infty$ ) and use above result.
- Other models beyond scope of this subject.

# Forecast intervals

## 95% forecast interval

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where  $v_{T+h|T}$  is estimated forecast variance.

- Multi-step forecast intervals for ARIMA(0,0,q):

$$y_t = e_t + \sum_{i=1}^q \theta_i e_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[ 1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

- AR(1): Rewrite as MA( $\infty$ ) and use above result.
- Other models beyond scope of this subject.

# Forecast intervals

## 95% forecast interval

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where  $v_{T+h|T}$  is estimated forecast variance.

- Multi-step forecast intervals for ARIMA(0,0,q):

$$y_t = e_t + \sum_{i=1}^q \theta_i e_{t-i}.$$

$$v_{T|T+h} = \hat{\sigma}^2 \left[ 1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

- AR(1): Rewrite as MA( $\infty$ ) and use above result.
- Other models beyond scope of this subject.

# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.

# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
  - Prediction intervals can be difficult to calculate by hand
  - Calculations assume residuals are **uncorrelated** and **normally distributed.**
  - Prediction intervals tend to be too narrow.
- Prediction intervals are based on the assumption that the model assumptions are correct. If the model assumptions are incorrect, the prediction intervals will be too narrow.
- Prediction intervals are based on the assumption that the model assumptions are correct. If the model assumptions are incorrect, the prediction intervals will be too narrow.
- Prediction intervals are based on the assumption that the model assumptions are correct. If the model assumptions are incorrect, the prediction intervals will be too narrow.
- Prediction intervals are based on the assumption that the model assumptions are correct. If the model assumptions are incorrect, the prediction intervals will be too narrow.



# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
  - the uncertainty in the parameter estimates has not been accounted for.
  - the ARIMA model assumes historical patterns will not change during the forecast period.
  - the ARIMA model assumes uncorrelated future

# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
  - the uncertainty in the parameter estimates has not been accounted for.
  - the ARIMA model assumes historical patterns will not change during the forecast period.
  - the ARIMA model assumes uncorrelated future errors

# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
  - the uncertainty in the parameter estimates has not been accounted for.
  - the ARIMA model assumes historical patterns will not change during the forecast period.
  - the ARIMA model assumes uncorrelated future errors

# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
  - the uncertainty in the parameter estimates has not been accounted for.
  - the ARIMA model assumes historical patterns will not change during the forecast period.
  - the ARIMA model assumes uncorrelated future errors

# Prediction intervals

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
  - the uncertainty in the parameter estimates has not been accounted for.
  - the ARIMA model assumes historical patterns will not change during the forecast period.
  - the ARIMA model assumes uncorrelated future errors