



Rob J Hyndman

Forecasting using



12. Advanced methods

[OTexts.com/fpp/2/5/](https://otexts.com/fpp/2/5/)

Outline

- 1 Cross-validation**
- 2 Time series with complex seasonality
- 3 Forecasting proportions
- 4 Some case studies
- 5 Forecasting resources

Cross-validation

Standard cross-validation

A more sophisticated version of training/test sets.

- Select one observation for test set, and use remaining observations in training set.
Compute error on test observation.
- Repeat using each possible observation as the test set.
- Compute accuracy measure over all errors.
- Does not work normally for time series because we cannot use future observations to build a model.

Cross-validation

Standard cross-validation

A more sophisticated version of training/test sets.

- Select one observation for test set, and use remaining observations in training set.
Compute error on test observation.
- Repeat using each possible observation as the test set.
- Compute accuracy measure over all errors.
- Does not work normally for time series because we cannot use future observations to build a model.

Cross-validation

Standard cross-validation

A more sophisticated version of training/test sets.

- Select one observation for test set, and use remaining observations in training set.
Compute error on test observation.
- Repeat using each possible observation as the test set.
- Compute accuracy measure over all errors.
- Does not work normally for time series because we cannot use future observations to build a model.

Cross-validation

Standard cross-validation

A more sophisticated version of training/test sets.

- Select one observation for test set, and use remaining observations in training set.
Compute error on test observation.
- Repeat using each possible observation as the test set.
- Compute accuracy measure over all errors.
- Does not work normally for time series because we cannot use future observations to build a model.

Time series cross-validation

Assume k is the minimum number of observations for a training set.

- Select observation $k + i$ for test set, and use observations at times $1, 2, \dots, k + i - 1$ to estimate model. Compute error on forecast for time $k + i$.
- Repeat for $i = 0, 1, \dots, T - k$ where T is total number of observations.
- Compute accuracy measure over all errors.

Also called **rolling forecasting origin** because the origin ($k + i - 1$) at which forecast is based rolls forward in time.

Time series cross-validation

Assume k is the minimum number of observations for a training set.

- Select observation $k + i$ for test set, and use observations at times $1, 2, \dots, k + i - 1$ to estimate model. Compute error on forecast for time $k + i$.
- Repeat for $i = 0, 1, \dots, T - k$ where T is total number of observations.
- Compute accuracy measure over all errors.

Also called **rolling forecasting origin** because the origin ($k + i - 1$) at which forecast is based rolls forward in time.

Time series cross-validation

Assume k is the minimum number of observations for a training set.

- Select observation $k + i$ for test set, and use observations at times $1, 2, \dots, k + i - 1$ to estimate model. Compute error on forecast for time $k + i$.
- Repeat for $i = 0, 1, \dots, T - k$ where T is total number of observations.
- Compute accuracy measure over all errors.

Also called **rolling forecasting origin** because the origin ($k + i - 1$) at which forecast is based rolls forward in time.

Time series cross-validation

Assume k is the minimum number of observations for a training set.

- Select observation $k + i$ for test set, and use observations at times $1, 2, \dots, k + i - 1$ to estimate model. Compute error on forecast for time $k + i$.
- Repeat for $i = 0, 1, \dots, T - k$ where T is total number of observations.
- Compute accuracy measure over all errors.

Also called **rolling forecasting origin** because the origin ($k + i - 1$) at which forecast is based rolls forward in time.

Time series cross-validation


Assume k is the minimum number of observations for a training set.

- Select observation $k + i$ for test set, and use observations at times $1, 2, \dots, k + i - 1$ to estimate model. Compute error on forecast for time $k + i$.
- Repeat for $i = 0, 1, \dots, T - k$ where T is total number of observations.
- Compute accuracy measure over all errors.

Also called **rolling forecasting origin** because the origin ($k + i - 1$) at which forecast is based rolls forward in time.


Example: Pharmaceutical sales

Antidiabetic drug sales



Example: Pharmaceutical sales

Log Antidiabetic drug sales



Example: Pharmaceutical sales

Which of these models is best?

- 1 Linear model with trend and seasonal dummies applied to log data.
- 2 ARIMA model applied to log data
- 3 ETS model applied to original data

Example: Pharmaceutical sales

Which of these models is best?

- 1 Linear model with trend and seasonal dummies applied to log data.
- 2 ARIMA model applied to log data
- 3 ETS model applied to original data

Example: Pharmaceutical sales

Which of these models is best?

- 1 Linear model with trend and seasonal dummies applied to log data.
- 2 ARIMA model applied to log data
- 3 ETS model applied to original data

■ Set $k = 48$ as minimum training set.

■ For each k , fit the three models and calculate the cross-validated error.

■ For each k , calculate the mean cross-validated error and its standard error.

■ Plot the mean cross-validated error and its standard error against k .

Example: Pharmaceutical sales

Which of these models is best?

- 1 Linear model with trend and seasonal dummies applied to log data.
- 2 ARIMA model applied to log data
- 3 ETS model applied to original data

- Set $k = 48$ as minimum training set.
- Forecast 12 steps ahead based on data to time $k + i - 1$ for $i = 1, 2, \dots, 156$.
- Compare MAE values for each forecast horizon.

Example: Pharmaceutical sales

Which of these models is best?

- 1 Linear model with trend and seasonal dummies applied to log data.
 - 2 ARIMA model applied to log data
 - 3 ETS model applied to original data
-
- Set $k = 48$ as minimum training set.
 - Forecast 12 steps ahead based on data to time $k + i - 1$ for $i = 1, 2, \dots, 156$.
 - Compare MAE values for each forecast horizon.

Example: Pharmaceutical sales

Which of these models is best?


- 1 Linear model with trend and seasonal dummies applied to log data.
 - 2 ARIMA model applied to log data
 - 3 ETS model applied to original data
-
- Set $k = 48$ as minimum training set.
 - Forecast 12 steps ahead based on data to time $k + i - 1$ for $i = 1, 2, \dots, 156$.
 - Compare MAE values for each forecast horizon.

Example: Pharmaceutical sales


Which of these models is best?

- 1 Linear model with trend and seasonal dummies applied to log data.
 - 2 ARIMA model applied to log data
 - 3 ETS model applied to original data
-
- Set $k = 48$ as minimum training set.
 - Forecast 12 steps ahead based on data to time $k + i - 1$ for $i = 1, 2, \dots, 156$.
 - Compare MAE values for each forecast horizon.

Example: Pharmaceutical sales



Example: Pharmaceutical sales



Example: Pharmaceutical sales

```
k <- 48
n <- length(a10)
mae1 <- mae2 <- mae3 <- matrix(NA,n-k-12,12)
for(i in 1:(n-k-12))
{
  xshort <- window(a10,end=1995+(5+i)/12)
  xnext <- window(a10,start=1995+(6+i)/12,end=1996+(5+i)/12)
  fit1 <- tslm(xshort ~ trend + season, lambda=0)
  fcast1 <- forecast(fit1,h=12)
  fit2 <- auto.arima(xshort,D=1, lambda=0)
  fcast2 <- forecast(fit2,h=12)
  fit3 <- ets(xshort)
  fcast3 <- forecast(fit3,h=12)
  mae1[i,] <- abs(fcast1[['mean']]-xnext)
  mae2[i,] <- abs(fcast2[['mean']]-xnext)
  mae3[i,] <- abs(fcast3[['mean']]-xnext)
}
plot(1:12,colMeans(mae1),type="l",col=2,xlab="horizon",ylab="MAE",
     ylim=c(0.58,1.0))
lines(1:12,colMeans(mae2),type="l",col=3)
lines(1:12,colMeans(mae3),type="l",col=4)
legend("topleft",legend=c("LM","ARIMA","ETS"),col=2:4,lty=1)
```

Variations on time series cross validation

- Keep training window of fixed length.

```
xshort <- window(a10,start=i+1/12,end=1995+(5+i)/12)
```

- Compute one-step forecasts in out-of-sample period.


```
for(i in 1:(n-k))  
{  
  xshort <- window(a10,end=1995+(5+i)/12)  
  xlong <- window(a10,start=1995+(6+i)/12)  
  fit2 <- auto.arima(xshort,D=1, lambda=0)  
  fit2a <- Arima(xlong,model=fit2)  
  fit3 <- ets(xshort)  
  fit3a <- ets(xlong,model=fit3)  
  mae2a[i,] <- abs(residuals(fit3a))  
  mae3a[i,] <- abs(residuals(fit2a))  
}
```


Outline

- 1 Cross-validation
- 2 Time series with complex seasonality**
- 3 Forecasting proportions
- 4 Some case studies
- 5 Forecasting resources


Examples

US finished motor gasoline products




Examples

Number of calls to large American bank (7am–9pm)



Examples

Turkish electricity demand



TBATS

Trigonometric terms for seasonality

Box-Cox transformations for heterogeneity

ARMA errors for short-term dynamics

Trend (possibly damped)

Seasonal (including multiple and
non-integer periods)

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t & \text{if } \omega = 0. \end{cases}$$

$$y_t^{(\omega)} = \ell_{t-1} + \phi b_{t-1} + \sum_{i=1}^M s_{t-m_i}^{(i)} + d_t$$

$$\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha d_t$$

$$b_t = (1 - \phi)b + \phi b_{t-1} + \beta d_t$$

$$d_t = \sum_{i=1}^p \phi_i d_{t-i} + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$$

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)} \quad \begin{aligned} s_{j,t}^{(i)} &= s_{j,t-1}^{(i)} \cos \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \sin \lambda_j^{(i)} + \gamma_1^{(i)} d_t \\ s_{j,t}^{(i)} &= -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t \end{aligned}$$

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t & \text{if } \omega = 0. \end{cases}$$

Box-Cox transformation

$$y_t^{(\omega)} = \ell_{t-1} + \phi b_{t-1} + \sum_{i=1}^M s_{t-m_i}^{(i)} + d_t$$

$$\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha d_t$$

$$b_t = (1 - \phi)b + \phi b_{t-1} + \beta d_t$$

$$d_t = \sum_{i=1}^p \phi_i d_{t-i} + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$$

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)} \quad \begin{aligned} s_{j,t}^{(i)} &= s_{j,t-1}^{(i)} \cos \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \sin \lambda_j^{(i)} + \gamma_1^{(i)} d_t \\ s_{j,t}^{(i)} &= -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t \end{aligned}$$

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t & \text{if } \omega = 0. \end{cases}$$

Box-Cox transformation

$$y_t^{(\omega)} = \ell_{t-1} + \phi b_{t-1} + \sum_{i=1}^M s_{t-m_i}^{(i)} + d_t$$

M seasonal periods

$$\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha d_t$$

$$b_t = (1 - \phi)b + \phi b_{t-1} + \beta d_t$$

$$d_t = \sum_{i=1}^p \phi_i d_{t-i} + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$$

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)} \quad \begin{aligned} s_{j,t}^{(i)} &= s_{j,t-1}^{(i)} \cos \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \sin \lambda_j^{(i)} + \gamma_1^{(i)} d_t \\ s_{j,t}^{(i)} &= -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t \end{aligned}$$

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t & \text{if } \omega = 0. \end{cases}$$

Box-Cox transformation

$$y_t^{(\omega)} = \ell_{t-1} + \phi b_{t-1} + \sum_{i=1}^M s_{t-m_i}^{(i)} + d_t$$

M seasonal periods

$$\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha d_t$$

global and local trend

$$b_t = (1 - \phi)b + \phi b_{t-1} + \beta d_t$$

$$d_t = \sum_{i=1}^p \phi_i d_{t-i} + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$$

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)} \quad \begin{aligned} s_{j,t}^{(i)} &= s_{j,t-1}^{(i)} \cos \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \sin \lambda_j^{(i)} + \gamma_1^{(i)} d_t \\ s_{j,t}^{(i)} &= -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t \end{aligned}$$

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t & \text{if } \omega = 0. \end{cases}$$

Box-Cox transformation

$$y_t^{(\omega)} = \ell_{t-1} + \phi b_{t-1} + \sum_{i=1}^M s_{t-m_i}^{(i)} + d_t$$

M seasonal periods

$$\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha d_t$$

global and local trend

$$b_t = (1 - \phi)b + \phi b_{t-1} + \beta d_t$$

$$d_t = \sum_{i=1}^p \phi_i d_{t-i} + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$$

ARMA error

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)} \quad \begin{aligned} s_{j,t}^{(i)} &= s_{j,t-1}^{(i)} \cos \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \sin \lambda_j^{(i)} + \gamma_1^{(i)} d_t \\ s_{j,t}^{(i)} &= -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t \end{aligned}$$

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t & \text{if } \omega = 0. \end{cases}$$

Box-Cox transformation

$$y_t^{(\omega)} = \ell_{t-1} + \phi b_{t-1} + \sum_{i=1}^M s_{t-m_i}^{(i)} + d_t$$

M seasonal periods

$$\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha d_t$$

global and local trend

$$b_t = (1 - \phi)b + \phi b_{t-1} + \beta d_t$$

$$d_t = \sum_{i=1}^p \phi_i d_{t-i} + \sum_{j=1}^q \theta_j \varepsilon_{t-j} + \varepsilon_t$$

ARMA error

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)} \quad s_{j,t}^{(i)} = s_{j,t-1}^{(i)} \cos \lambda_j^{(i)} \quad s_{j,t}^{(i)} = -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t$$

Fourier-like seasonal terms

TBATS model

y_t = observation at time t

$$y_t^{(\omega)} = \begin{cases} (y_t^\omega - 1)/\omega & \text{if } \omega \neq 0; \\ \log y_t^\omega & \text{if } \omega = 0; \end{cases}$$

$$y_t^{(\omega)} = \ell_{t-1}$$

$$\ell_t = \ell_{t-1}$$

$$b_t = (1 -$$

$$d_t = \sum_{i=1}^p$$

$$s_t^{(i)} = \sum_{j=1}^{k_i} s_{j,t}^{(i)}$$

$$s_{j,t}^{(i)} = s_{j,t-1}^{(i)} c$$

$$s_{j,t}^{(i)} = -s_{j,t-1}^{(i)} \sin \lambda_j^{(i)} + s_{j,t-1}^{*(i)} \cos \lambda_j^{(i)} + \gamma_2^{(i)} d_t$$

TBATS

Trigonometric

Box-Cox

ARMA

Trend

Seasonal

Box-Cox transformation

M seasonal periods

global and local trend


ARMA error

Fourier-like seasonal terms

Examples

Forecasts from TBATS(0.999, {2,2}, 1, {<52.1785714285714,8>})

```
fit <- tbats(gasoline)
fcast <- forecast(fit)
plot(fcast)
```




Examples

Forecasts from TBATS(1, {3,1}, 0.987, {<169,5>, <845,3>})

```
fit <- tbats(callcentre)  
fcast <- forecast(fit)  
plot(fcast)
```


Number of call arrivals



Examples

Forecasts from TBATS(0, {5,3}, 0.997, {<7,3>, <354.37,12>, <365.25,4>})

```
fit <- tbats(turk)  
fcast <- forecast(fit)  
plot(fcast)
```



Outline

- 1 Cross-validation
- 2 Time series with complex seasonality
- 3 Forecasting proportions**
- 4 Some case studies
- 5 Forecasting resources

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:

$$f(u) = \log\left(\frac{u}{1-u}\right)$$

- Then build model and back-transform the forecasts.

More complicated:

Let y_t be a binary variable (e.g., water quality above some threshold value on day t).

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:

$$f(u) = \log\left(\frac{u}{1-u}\right)$$

- Then build model and back-transform the forecasts.

More complicated:

- Let y_t be a binary variable (e.g., water quality above some threshold value on day t).

Use a generalized linear model with a logit link function, e.g., $\text{Pr}(y_t = 1 | x_t) = \text{logit}^{-1}(x_t)$, where x_t is a vector of predictors and $\text{logit}^{-1}(u) = \exp(u) / (1 + \exp(u))$. The generalized linear model is implemented using the `glm` function in R. The `glm` function also provides a `predict` method for computing the predicted probabilities.

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:
$$f(u) = \log\left(\frac{u}{1-u}\right)$$
- Then build model and back-transform the forecasts.

More complicated:

- Let y_t be a binary variable (e.g., water quality above some threshold value on day t).
- Use a generalized linear model with a latent autocorrelated process. e.g., $\Pr(y_t) = f^{-1}(\alpha + \beta x_t + u_t)$ where x_t is a vector of predictors and u_t is a correlated process.
- Theory and methodology not well developed and very little software available.
- R and other packages allows some simple correlation

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:
$$f(u) = \log\left(\frac{u}{1-u}\right)$$
- Then build model and back-transform the forecasts.

More complicated:

- Let y_t be a binary variable (e.g., water quality above some threshold value on day t).
- Use a generalized linear model with a latent autocorrelated process. e.g., $\Pr(y_t) = f^{-1}(\alpha + \beta \mathbf{x}_t + u_t)$ where \mathbf{x}_t is a vector of predictors and u_t is a correlated process.
- Theory and methodology not well developed and very little software available.
- `build` and `gee` packages allows some simple correlation structures.

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:
$$f(u) = \log\left(\frac{u}{1-u}\right)$$
- Then build model and back-transform the forecasts.

More complicated:

- Let y_t be a binary variable (e.g., water quality above some threshold value on day t).
- Use a generalized linear model with a latent autocorrelated process. e.g., $\Pr(y_t) = f^{-1}(\alpha + \beta \mathbf{x}_t + u_t)$ where \mathbf{x}_t is a vector of predictors and u_t is a correlated process.
- Theory and methodology not well developed and very little software available.
- `bird` and `gee` packages allows some simple correlation structures.

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:
$$f(u) = \log\left(\frac{u}{1-u}\right)$$
- Then build model and back-transform the forecasts.

More complicated:

- Let y_t be a binary variable (e.g., water quality above some threshold value on day t).
- Use a generalized linear model with a latent autocorrelated process. e.g., $\Pr(y_t) = f^{-1}(\alpha + \beta \mathbf{x}_t + u_t)$ where \mathbf{x}_t is a vector of predictors and u_t is a correlated process.
- Theory and methodology not well developed and very little software available.
- `bird` and `gee` packages allows some simple correlation structures.

Forecasting proportions

Simple approach

- Use a logit transformation on proportions:
$$f(u) = \log\left(\frac{u}{1-u}\right)$$
- Then build model and back-transform the forecasts.

More complicated:

- Let y_t be a binary variable (e.g., water quality above some threshold value on day t).
- Use a generalized linear model with a latent autocorrelated process. e.g., $\Pr(y_t) = f^{-1}(\alpha + \beta \mathbf{x}_t + u_t)$ where \mathbf{x}_t is a vector of predictors and u_t is a correlated process.
- Theory and methodology not well developed and very little software available.
- **build and gee packages allows some simple correlation structures.**

Outline

- 1 Cross-validation
- 2 Time series with complex seasonality
- 3 Forecasting proportions
- 4 Some case studies**
- 5 Forecasting resources

CASE STUDY 1: Paperware company

Problem: Want forecasts of each of hundreds of items. Series can be stationary, trended or seasonal. They currently have a large forecasting program written in-house but it doesn't seem to produce sensible forecasts. They want me to tell them what is wrong and fix it.



CASE STUDY 1: Paperware company

Problem: Want forecasts of each of hundreds of items. Series can be stationary, trended or seasonal. They currently have a large forecasting program written in-house but it doesn't seem to produce sensible forecasts. They want me to tell them what is wrong and fix it.

Additional information

- Program written in COBOL making numerical calculations limited. It is not possible to do any optimisation.



CASE STUDY 1: Paperware company

Problem: Want forecasts of each of hundreds of items. Series can be stationary, trended or seasonal. They currently have a large forecasting program written in-house but it doesn't seem to produce sensible forecasts. They want me to tell them what is wrong and fix it.

Additional information

- Program written in COBOL making numerical calculations limited. It is not possible to do any optimisation.
- Their programmer has little experience in numerical computing.



CASE STUDY 1: Paperware company

Problem: Want forecasts of each of hundreds of items. Series can be stationary, trended or seasonal. They currently have a large forecasting program written in-house but it doesn't seem to produce sensible forecasts. They want me to tell them what is wrong and fix it.



Additional information

- Program written in COBOL making numerical calculations limited. It is not possible to do any optimisation.
- Their programmer has little experience in numerical computing.
- They employ no statisticians and want the program to produce forecasts automatically.

CASE STUDY 1: Paperware company

Methods currently used

- A** 12 month average
- C** 6 month average
- E** straight line regression over last 12 months
- G** straight line regression over last 6 months
- H** average slope between last year's and this year's values.
(Equivalent to differencing at lag 12 and taking mean.)
- I** Same as H except over 6 months.
- K** I couldn't understand the explanation.

CASE STUDY 2: PBS



CASE STUDY 2: PBS

The **Pharmaceutical Benefits Scheme** (PBS) is the Australian government drugs subsidy scheme.

- Many drugs bought from pharmacies are subsidised to allow more equitable access to modern drugs.
- The cost to government is determined by the number and types of drugs purchased. Currently nearly 1% of GDP.
- The total cost is budgeted based on forecasts of drug usage.

CASE STUDY 2: PBS

The **Pharmaceutical Benefits Scheme** (PBS) is the Australian government drugs subsidy scheme.

- Many drugs bought from pharmacies are subsidised to allow more equitable access to modern drugs.
- The cost to government is determined by the number and types of drugs purchased. Currently nearly 1% of GDP.
- The total cost is budgeted based on forecasts of drug usage.

CASE STUDY 2: PBS

The **Pharmaceutical Benefits Scheme** (PBS) is the Australian government drugs subsidy scheme.

- Many drugs bought from pharmacies are subsidised to allow more equitable access to modern drugs.
- The cost to government is determined by the number and types of drugs purchased. Currently nearly 1% of GDP.
- The total cost is budgeted based on forecasts of drug usage.

CASE STUDY 2: PBS

The **Pharmaceutical Benefits Scheme** (PBS) is the Australian government drugs subsidy scheme.

- Many drugs bought from pharmacies are subsidised to allow more equitable access to modern drugs.
- The cost to government is determined by the number and types of drugs purchased. Currently nearly 1% of GDP.
- The total cost is budgeted based on forecasts of drug usage.

CASE STUDY 2: PBS



ABC News Online
AUSTRALIAN BROADCASTING CORPORATION



NewsRadio
Streaming audio news

LISTEN: [WMP](#) | [Real](#)

Select a Topic
from the list below

Click "Refresh" or "Reload"
on your browser for the latest edition.

This Bulletin: Wed, May 30 2001 6:22 PM AES

POLITICS

Opp demands drug price restriction after PBS budget blow-out

The Federal Opposition has called for tighter controls on drug prices after the Pharmaceutical Benefits Scheme (PBS) budget blew out by almost \$800 million.

The money was spent on two new drugs including the controversial anti-smoking aid Zyban, which dropped in price from \$220 to \$22 after it was listed on the PBS.

**the
Public Record**
For full election coverage

FEATURES

**the
Public Record**
Federal Election 2001

[For a fresh perspective on the federal election, reach into ABC Online's campaign weblog, The Poll Vault.](#)

Audio News Online

SPECIALS

[Federal Election](#)

CASE STUDY 2: PBS

- In 2001: \$4.5 billion budget, under-forecasted by \$800 million.
- Thousands of products. Seasonal demand.
- Subject to covert marketing, volatile products, uncontrollable expenditure.
- Although monthly data available for 10 years, data are aggregated to annual values, and only the first three years are used in estimating the forecasts.
- All forecasts being done with the FORECAST function in MS-Excel!

Problem: How to do the forecasting better?

CASE STUDY 2: PBS

- In 2001: \$4.5 billion budget, under-forecasted by \$800 million.
- Thousands of products. Seasonal demand.
- Subject to covert marketing, volatile products, uncontrollable expenditure.
- Although monthly data available for 10 years, data are aggregated to annual values, and only the first three years are used in estimating the forecasts.
- All forecasts being done with the FORECAST function in MS-Excel!

Problem: How to do the forecasting better?

CASE STUDY 2: PBS

- In 2001: \$4.5 billion budget, under-forecasted by \$800 million.
- Thousands of products. Seasonal demand.
- Subject to covert marketing, volatile products, uncontrollable expenditure.
- Although monthly data available for 10 years, data are aggregated to annual values, and only the first three years are used in estimating the forecasts.
- All forecasts being done with the FORECAST function in MS-Excel!

Problem: How to do the forecasting better?

CASE STUDY 2: PBS

- In 2001: \$4.5 billion budget, under-forecasted by \$800 million.
- Thousands of products. Seasonal demand.
- Subject to covert marketing, volatile products, uncontrollable expenditure.
- Although monthly data available for 10 years, data are aggregated to annual values, and only the first three years are used in estimating the forecasts.
- All forecasts being done with the FORECAST function in MS-Excel!

Problem: How to do the forecasting better?

CASE STUDY 2: PBS

- In 2001: \$4.5 billion budget, under-forecasted by \$800 million.
- Thousands of products. Seasonal demand.
- Subject to covert marketing, volatile products, uncontrollable expenditure.
- Although monthly data available for 10 years, data are aggregated to annual values, and only the first three years are used in estimating the forecasts.
- All forecasts being done with the FORECAST function in MS-Excel!

Problem: How to do the forecasting better?

CASE STUDY 2: PBS

- In 2001: \$4.5 billion budget, under-forecasted by \$800 million.
- Thousands of products. Seasonal demand.
- Subject to covert marketing, volatile products, uncontrollable expenditure.
- Although monthly data available for 10 years, data are aggregated to annual values, and only the first three years are used in estimating the forecasts.
- All forecasts being done with the FORECAST function in MS-Excel!


Problem: How to do the forecasting better?

CASE STUDY 3: Airline




CASE STUDY 3: Airline


First class passengers: Melbourne–Sydney



Business class passengers: Melbourne–Sydney




Economy class passengers: Melbourne–Sydney




CASE STUDY 3: Airline

First class passengers: Melbourne–Sydney




Not the real data!
Or is it?

First class passengers: Melbourne–Sydney



Year

Economy class passengers: Melbourne–Sydney



Year

CASE STUDY 3: Airline

Problem: how to forecast passenger traffic on major routes.

Additional information

- They can provide a large amount of data on previous routes.
- Traffic is affected by school holidays, special events such as the Grand Prix, advertising campaigns, competition behaviour, etc.
- They have a highly capable team of people who are able to do most of the computing.

CASE STUDY 3: Airline

Problem: how to forecast passenger traffic on major routes.

Additional information

- They can provide a large amount of data on previous routes.
- Traffic is affected by school holidays, special events such as the Grand Prix, advertising campaigns, competition behaviour, etc.
- They have a highly capable team of people who are able to do most of the computing.

CASE STUDY 3: Airline

Problem: how to forecast passenger traffic on major routes.

Additional information

- They can provide a large amount of data on previous routes.
- Traffic is affected by school holidays, special events such as the Grand Prix, advertising campaigns, competition behaviour, etc.
- They have a highly capable team of people who are able to do most of the computing.

CASE STUDY 3: Airline

Problem: how to forecast passenger traffic on major routes.

Additional information

- They can provide a large amount of data on previous routes.
- Traffic is affected by school holidays, special events such as the Grand Prix, advertising campaigns, competition behaviour, etc.
- They have a highly capable team of people who are able to do most of the computing.

Outline

- 1 Cross-validation
- 2 Time series with complex seasonality
- 3 Forecasting proportions
- 4 Some case studies
- 5 Forecasting resources**

Useful resources

■ Organization:

- International Institute of Forecasters.

■ Conferences:

- International Symposium on Forecasting. June 2014, Rotterdam.

■ Journals:

- International Journal of Forecasting
- Foresight

Useful resources

■ Organization:

- International Institute of Forecasters.

■ Conferences:

- International Symposium on Forecasting. June 2014, Rotterdam.

■ Journals:

- International Journal of Forecasting
- Foresight



VOLUME 23, NUMBER 3

ISSN 0169-2070

international journal of forecasting



International Institute of Forecasters

Useful resources

■ Organization:


- International Institute of Forecasters.

■ Conferences:

- International Symposium on Forecasting. June 2014, Rotterdam.

■ Journals:

- International Journal of Forecasting
- Foresight



Useful resources

■ Organization:

- International Institute of Forecasters.


■ Conferences:

- International Symposium on Forecasting. June 2014, Rotterdam.

■ Journals:

- International Journal of Forecasting
- Foresight

Links to all of the above at
www.forecasters.org



Final comments

- Revolution Analytics will send you course completion certificates.
- Please vote on the students who have made most contribution to class. You will receive a voting form by email. I will announce the results on Piazza.
- I will continue to answer questions on Piazza until Christmas.
- A discussion forum for forecasting planned for OTexts. Should be launched in early 2014.
- The FPP book will be available in print on Amazon by February 2014. Maybe a kindle version will follow.

Final comments

- Revolution Analytics will send you course completion certificates.
- Please vote on the students who have made most contribution to class. You will receive a voting form by email. I will announce the results on Piazza.
- I will continue to answer questions on Piazza until Christmas.
- A discussion forum for forecasting planned for OTexts. Should be launched in early 2014.
- The FPP book will be available in print on Amazon by February 2014. Maybe a kindle version will follow.

Final comments

- Revolution Analytics will send you course completion certificates.
- Please vote on the students who have made most contribution to class. You will receive a voting form by email. I will announce the results on Piazza.
- I will continue to answer questions on Piazza until Christmas.
- A discussion forum for forecasting planned for OTexts. Should be launched in early 2014.
- The FPP book will be available in print on Amazon by February 2014. Maybe a kindle version will follow.

Final comments

- Revolution Analytics will send you course completion certificates.
- Please vote on the students who have made most contribution to class. You will receive a voting form by email. I will announce the results on Piazza.
- I will continue to answer questions on Piazza until Christmas.
- A discussion forum for forecasting planned for OTexts. Should be launched in early 2014.
- The FPP book will be available in print on Amazon by February 2014. Maybe a kindle version will follow.

Final comments

- Revolution Analytics will send you course completion certificates.
- Please vote on the students who have made most contribution to class. You will receive a voting form by email. I will announce the results on Piazza.
- I will continue to answer questions on Piazza until Christmas.
- A discussion forum for forecasting planned for OTexts. Should be launched in early 2014.
- The FPP book will be available in print on Amazon by February 2014. Maybe a kindle version will follow.