

Assignment 3

Assignment 4

AI1110: Probability and Random Variables

Indian Institute of Technology Hyderabad

Rudransh Mishra
AI21BTECH11025

29 May 2022

PROBABILITY, RANDOM VARIABLES, AND STOCHASTIC
PROCESSES

Athanasios Papoulis

L^AT_EX

Question

Example 1-3

We are given a circle C of radius r and we wish to determine the probability p that the length l of a "randomly selected" cord AB is greater than the length $r\sqrt{3}$ of the inscribed equilateral triangle.

Solution

We shall show that this problem can be given at least three reasonable solutions.

- 1 If the center M of the cord A B lies inside the circle of radius $\frac{r}{2}$ then $l > r\sqrt{3}$. It is reasonable, therefore, to consider as favorable outcomes all points inside the inner circle and as possible outcomes all points inside the outer circle. Using as measure of their numbers the corresponding areas $\pi\frac{r^2}{4}$ and πr^2 , we conclude that

$$P = \frac{\pi\frac{r^2}{4}}{\pi r^2} = \frac{1}{4} \quad (1)$$

Solution(contd.)

- ② We now assume that the end A of the cord AB is fixed. This reduces the number of possibilities but it has no effect on the value of p because the number of favorable locations of B is reduced proportionately. If B is on the 120° arc directly opposite to the point A, then $1 > r\sqrt{3}$. The favorable outcomes are now the points on this arc and the total outcomes all points on the circumference of the circle C. Using as their measurements the corresponding lengths $\frac{2\pi r}{3}$ and $2\pi r$, we obtain

$$P = \frac{\frac{2\pi r}{3}}{2\pi r} = \frac{1}{3} \quad (2)$$

Solution(contd.)

- ③ We assume finally that the direction of AB is perpendicular to the diameter of the circle. As in (2) this restriction has no effect on the value of P. If the center M of AB is less than $\frac{r}{2}$ distance from the center, then $1 > r\sqrt{3}$. Favorable outcomes are now the points on the line till $\frac{r}{2}$ distance on either side of the center and possible outcomes all points on the diameter. Using as their measures the respective lengths r and $2r$. we obtain

$$P = \frac{r}{2r} = \frac{1}{2} \quad (3)$$

Solution(contd.)

We have thus found not one but three different solutions for the same problem! One might remark that these solutions correspond to three different experiments. This is true but not obvious and, in any case, it demonstrates the ambiguities associated with the classical definition, and the need for a clear specification of the outcomes of an experiment and the meaning of the terms "possible" and "favorable."

Validity

Validity. We shall now discuss the value of the classical definition in the determination of probabilistic data and as a working hypothesis.

- 1 In many applications, the assumption that there are N equally likely alternatives is well established through long experience. For example, "If a ball is selected at random from a box containing m black and n white balls, the probability that it is white equals $n/(m + n)$," or, "If a call occurs at random in the time interval $(0, T)$, the probability that it occurs in the interval (t_1, t_2) equals $\frac{t_2 - t_1}{T}$ ".

Such conclusions are of course, valid and useful; however, their Validity rests on the meaning of the word random. The conclusion of the last example that "the unknown probability equals $\frac{t_2 - t_1}{T}$ " is not a consequence of the "randomness" of the call. The two statements are merely equivalent and they follow not from a prior reasoning but from past records of telephone calls.

Validity(contd.)

- 2 In a number of applications it is impossible to determine the probabilities of various events by repeating the underlying experiment a sufficient number of times. In such cases, we have no choice but to assume that certain alternatives are equally likely and to determine the desired probabilities from (1-7). This means that we use the classical definition as a working hypothesis. The hypothesis is accepted if its observable consequences agree with experience, otherwise it is rejected.

Python Code

```
import random
R=random.random()
pi=3.1415
#Method1
P=(pi*((R/2)**2))/(pi*(R**2))
print(" The first probability is :", P)
#Method2
P=((2*pi)/3)*(R)/(2*pi*R)
print(" The second probability is :", P)
#Method3
P=((R/2)/(R))
print(" The third probability is :", P)
```