

# Assignment 6

## AI1110: Probability and Random Variables

### Indian Institute of Technology Hyderabad

Rudransh Mishra  
AI21BTECH11025

15 June 2022

PROBABILITY, RANDOM VARIABLES, AND STOCHASTIC PROCESSES  
Athanasios Papoulis

**Example 9-38** Show that if  $R(r)$  is the inverse Fourier transform of a function  $S(\omega)$  and  $S(\omega) \geq 0$ , then, for any  $a_i$ ,

$$\sum_{i,k} a_i a_k^* R(\tau_i - \tau_k) \geq 0 \quad (1)$$

**Proof.** We know that,

$$\int_{-\infty}^{\infty} S(\omega) \left| \sum_i a_i e^{j\omega\tau_i} \right| d(\omega) \geq 0 \quad (2)$$

And,

$$\int_{-\infty}^{\infty} S(\omega) \left| \sum_i a_i e^{j\omega\tau_i} \right| d(\omega) \quad (3)$$

$$= \int_{-\infty}^{\infty} S(\omega) \left| \sum_{i,k} a_i a_k^* e^{j\omega(\tau_i - \tau_k)} \right| d(\omega) \quad (4)$$

Now, we know that,

$$\int_{-\infty}^{\infty} S(\omega) \left| \sum_{i,k} a_i a_k^* e^{j\omega(\tau_i - \tau_k)} \right| d(\omega) \quad (5)$$

$$= \sum_{i,k} a_i a_k^* R(\tau_i - \tau_k) \quad (6)$$

From (2) and (6),

$$\sum_{i,k} a_i a_k^* R(\tau_i - \tau_k) \geq 0 \quad (7)$$

and the proof is complete.

—X-X-X-X-X-X—