

Assignment 7

AI1110: Probability and Random Variables Indian Institute of Technology Hyderabad

Rudransh Mishra
AI21BTECH11025

19 June 2022

PROBABILITY, RANDOM VARIABLES, AND STOCHASTIC
PROCESSES

Athanasios Papoulis

L^AT_EX

Example 15.2

Consider a population that is able to produce new offspring of like kind. For each member let $p_k, k = 0, 1, 2, \dots$ represent the probability of creating k new members. The direct descendants of the n th generation form the $(n + 1)$ st generation. The members of each generation are independent of each other. Suppose X_n represents the size of the n th generation. It is clear that X_n depends only on X_{n-1} since $X_n = \sum_{i=1}^{X_{n-1}} Y_i$, where Y_i represents the number of offspring of the i th member of the $(n - 1)$ generation, and the manner in which the value of X_{n-1} was reached is of no consequence. Thus X_n represents a Markov chain.

Example(cont.)

Nuclear chain reactions, survival of family surnames, gene mutations, and waiting lines in a queueing system are all examples of branching processes. In a nuclear chain reaction, a particle such as a neutron scores a hit with probability p , creating m new particles, and $q = 1 - p$ represents the probability that it remains inactive with no descendants. In that case, the only possible number of descendants is zero and m with probabilities q and p . If P is close to one, the number of particles is likely to increase indefinitely, leading to an explosion, whereas if P is close to zero the process may never start.