

Cluster-forming $t > 3.98$ ($p < 1e-4$) · cluster-extent FWE $p < .05$
 model7 · $n = 58$ · surviving extent, % of mask: RPE +37.5 / -0.1 uRPE +1.0 / -8.6 Surprise +0.4 / -0.0
 model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.




Cluster-forming $t > 3.98$ ($p < 1e-4$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +37.5 / -0.1 uRPE +1.0 / -8.6 Surprise +0.4 / -0.0

model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

b Axial sections




Same cuts in every row, so extent can be compared straight down a column. Rows are the six maps on their own; the last row is all of them together.

Cluster-forming $t > 3.98$ ($p < 1e-4$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +37.5 / -0.1 uRPE +1.0 / -8.6 Surprise +0.4 / -0.0

model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

c Coronal sections





Same cuts in every row, so extent can be compared straight down a column. Rows are the six maps on their own; the last row is all of them together.

Cluster-forming $t > 3.98$ ($p < 1e-4$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +37.5 / -0.1 uRPE +1.0 / -8.6 Surprise +0.4 / -0.0

model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

d Which contrasts claim each piece of tissue



A voxel's category is the set of contrasts that call it significant, so "only" means no other contrast claims it. The remaining 5 combinations hold 133 voxels and are charted but not mapped.