


No cluster-forming threshold · TFCE, two-tailed FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +48.7 / -0.0 uRPE +0.0 / -26.6 Surprise +0.0 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

a




Rows

One cluster found by one of the six contrasts. The bracket says whether the signed-RPE map covers it ($\geq 5\%$). The number keys the row to its peak on the map pages.


Covered by map

Share of the ROI's voxels inside that map's surviving clusters.



Effect in ROI

Group one-sample t of that signal's contrast value, averaged over the ROI.




Bigger marker = a larger effect.

Holm-corrected over all ROI × signal tests.

Are they different?

One signal against another inside the ROI, paired across subjects.



TOST bound in the table is $dz = 0.368$, the smallest effect this design can


No cluster-forming threshold · TFCE, two-tailed FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +48.7 / -0.0 uRPE +0.0 / -26.6 Surprise +0.0 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

b Axial sections · $z = -20$ to $+50$ mm

one row per map, then all of them together



Same axial cuts in every row, so extent can be compared straight down a column.


No cluster-forming threshold · TFCE, two-tailed FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +48.7 / -0.0 uRPE +0.0 / -26.6 Surprise +0.0 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

c Coronal sections · $y = -88$ to $+42$ mm

one row per map, then all of them together



Same coronal cuts in every row, so extent can be compared straight down a column.



No cluster-forming threshold · TFCE, two-tailed FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +48.7 / -0.0 uRPE +0.0 / -26.6 Surprise +0.0 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

d Which contrasts claim each piece of tissue

every combination, then the six biggest mapped



A voxel's category is the set of contrasts that call it significant, so "only" means no other contrast claims it.