


Cluster-forming $t > 2.39$ ($p < .01$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +64.9 / -0.0 uRPE +2.2 / -30.3 Surprise +1.0 / -0.0

model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

a




Rows

One cluster found by one of the six contrasts. The bracket says whether the signed-RPE map covers it ($\geq 5\%$). The number keys the row to its peak on the map pages.


Covered by map

Share of the ROI's voxels inside that map's surviving clusters.



Effect in ROI


Group one-sample t of that signal's contrast value, averaged over the ROI.



Bigger marker = a larger effect.
Holm-corrected over all ROI $\times$ signal tests.

Are they different?

One signal against another inside the ROI, paired across subjects.



TOST bound in the table is $dz = 0.368$, the smallest effect this design can find with 80%.

Cluster-forming $t > 2.39$ ($p < .01$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +64.9 / -0.0 uRPE +2.2 / -30.3 Surprise +1.0 / -0.0


model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

b Axial sections · $z = -20$ to $+50$ mm

one row per map, then all of them together

RPE positive

38,852 voxels · 64.9% of mask




RPE negative

no surviving clusters




uRPE positive

1,299 voxels · 2.2% of mask




uRPE negative

18,112 voxels · 30.3% of mask




Surprise positive

616 voxels · 1.0% of mask




Surprise negative

no surviving clusters



All together

numbers = ROIs of page 1



Same axial cuts in every row, so extent can be compared straight down a column.

Cluster-forming $t > 2.39$ ($p < .01$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +64.9 / -0.0 uRPE +2.2 / -30.3 Surprise +1.0 / -0.0


model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

c Coronal sections · $y = -88$ to $+42$ mm

one row per map, then all of them together


RPE positive

38,852 voxels · 64.9% of mask




RPE negative

no surviving clusters




uRPE positive

1,299 voxels · 2.2% of mask




uRPE negative

18,112 voxels · 30.3% of mask




Surprise positive

616 voxels · 1.0% of mask




Surprise negative

no surviving clusters



All together

numbers = ROIs of page 1



Same coronal cuts in every row, so extent can be compared straight down a column.

Cluster-forming $t > 2.39$ ($p < .01$) · cluster-extent FWE $p < .05$

model7 · $n = 58$ · surviving extent, % of mask: RPE +64.9 / -0.0 uRPE +2.2 / -30.3 Surprise +1.0 / -0.0

model7 — one GLM carrying all three signals: choice (surprise, value), feedback (uRPE first, then signed RPE). Because SPM orthogonalises serially within an event, uRPE keeps all variance it shares with signed RPE, and the RPE map is the residual. Apples-to-apples across signals.

d Which contrasts claim each piece of tissue

every combination, then the six biggest mapped

