

Cluster-forming $t > 3.24$ ($p < .001$) · cluster-extent FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +52.5 / -0.6 uRPE +2.0 / -16.7 Surprise +0.9 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

a

		Covered by map						Effect in ROI			Are they different?		
		RPE		uRPE		Surprise		RPE	uRPE	Surprise	uRPE vs Surprise	RPE vs uRPE	RPE vs Surprise
		+	-	+	-	+	-						
RPE+ & uRPE- & Surprise+ * — 5 clusters													
1	DLPFC L	found by uRPE -		·	·								
2	Angular gyrus L (surprise)	found by Surprise +		·	·								
3	Precuneus L	found by Surprise +		·	·								
4	DLPFC L (2)	found by Surprise +		·	·								
5	Angular gyrus R	found by Surprise +		·	·								
RPE+ & uRPE- * — 4 clusters													
6	Caudate / putamen R	found by RPE +		·	·		·						
7	Caudate / putamen R (2)	found by uRPE -		·	·		·						
8	Middle temporal gyrus R	found by uRPE -		·	·		·						
9	Inferior parietal lobule L	found by uRPE -		·	·		·						
RPE- & uRPE+ * — 2 clusters													
10	Insula R	found by RPE - and uRPE +	·			·	·						
11	dmPFC R	found by RPE - and uRPE +	·			·	·						
RPE+ & uRPE+ — 1 cluster													
12	Inferior parietal lobule R	found by uRPE +		·		·	·						
only uRPE+ — 1 cluster													
13	Insula L	found by uRPE +	·	·		·	·						

Profile groups — 5 of them across 13 ROIs

RPE+ & uRPE- & Surprise+ * (5) RPE+ & uRPE- * (4) RPE- & uRPE+ * (2) RPE+ & uRPE+ (1) only uRPE+ (1)

* the same tissue responds positively to one signal and negatively to another — these groups are listed first.
Membership is map coverage: a map counts if it covers at least a tenth of the ROI. It is not the difference tests — a signal can be in the tag and still be “undetermined” against another.

Rows	Covered by map	Effect in ROI	Are they different?
One cluster found by one of the six contrasts. Rows are boxed by profile — the set of maps covering at least a tenth of them — and the box is drawn in that profile's colour. The number keys the row to its peak on the map pages.	Share of the ROI's voxels inside that map's surviving clusters.	Group one-sample t of that signal's contrast value, averaged over the ROI.	One signal against another inside the ROI, paired across subjects.
100% of the ROI	100% of the ROI	Positive, and significant	Different; points to the larger
50% of the ROI	50% of the ROI	Negative, and significant	Equivalent ($BF_{01} > 3$)
10% of the ROI	10% of the ROI	Open = not significant	Undetermined
Open square = negative tail	Open square = negative tail	Bigger marker = a larger effect. Holm-corrected over all ROI × signal tests.	TOST bound in the table is $d_z = 0.368$, the smallest effect this design can find with 80% power.
Dot = below 1%, no overlap	Dot = below 1%, no overlap		


Cluster-forming $t > 3.24$ ($p < .001$) · cluster-extent FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +52.5 / -0.6 uRPE +2.0 / -16.7 Surprise +0.9 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

b Axial sections · $z = -20$ to $+50$ mm

one row per map, then all of them together



Same axial cuts in every row, so extent can be compared straight down a column.


Cluster-forming $t > 3.24$ ($p < .001$) · cluster-extent FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +52.5 / -0.6 uRPE +2.0 / -16.7 Surprise +0.9 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

c Coronal sections · $y = -88$ to $+42$ mm

one row per map, then all of them together



Same coronal cuts in every row, so extent can be compared straight down a column.

Cluster-forming $t > 3.24$ ($p < .001$) · cluster-extent FWE $p < .05$

model2 · $n = 58$ · surviving extent, % of mask: RPE +52.5 / -0.6 uRPE +2.0 / -16.7 Surprise +0.9 / -0.0

model2 — the GLM the paper reports: choice (surprise, value), feedback (signed RPE only). It has no uRPE regressor at all, so the uRPE rows and maps here come from model7, exactly as the manuscript mixes them. Its RPE is the full effect, not a residual.

d Which contrasts claim each piece of tissue

every combination, then the six biggest mapped

