

Problem Set 9 for lecture Mining Massive Datasets

Due January 8, 2018, 11:59 pm

Exercise 1

(4 points)

Suppose there are 100 items, numbered 1 to 100, and also 100 baskets, also numbered 1 to 100. Item i is in basket b if and only if i divides b with no remainder. Thus, item 1 is in all the baskets, item 2 is in all fifty of the even-numbered baskets, and so on. Basket 12 consists of items $\{1, 2, 3, 4, 6, 12\}$, since these are all the integers that divide 12. Answer the following questions (without programming) and explain how you have obtained the solution:

- a) If the support threshold is 5, which items are frequent?
- b) If the support threshold is 5, which pairs of items are frequent?
- c) What is the sum of the sizes of all the baskets?
- d) What is the confidence of the following association rules $R_1 = \{5, 7\} \rightarrow 2$ and $R_2 = \{2, 3, 4\} \rightarrow 5$.

Exercise 2

(3 points)

Using the same setup of the Exercise 1, apply the A-Priori Algorithm (“on paper”, i.e. without programming) with support threshold 5. Consider $k = 3$, (frequent items, pairs and triples) and submit as your solution the results of each pass of the algorithm.

Exercise 3

(3 points)

Consider the collection of twelve baskets depicted below. Each contains three of the six items 1 through 6.

$\{1, 2, 3\}$ $\{2, 3, 4\}$ $\{3, 4, 5\}$ $\{4, 5, 6\}$
 $\{1, 3, 5\}$ $\{2, 4, 6\}$ $\{1, 3, 4\}$ $\{2, 4, 5\}$
 $\{3, 5, 6\}$ $\{1, 2, 4\}$ $\{2, 3, 5\}$ $\{3, 4, 6\}$

Suppose the support threshold is 4. On the first pass of the PCY Algorithm we use a hash table with 11 buckets, and the set $\{i, j\}$ is hashed to bucket $i \times j \bmod 11$. Answer the following questions (without programming) and explain how you have obtained your results.

- a) By any method, compute the support for each item and each pair of items.
- b) Which pairs hash to which buckets?
- c) Which buckets are frequent?
- d) Which pairs are counted on the second pass of the PCY Algorithm?

Exercise 4**(Bonus, 3 points)**

Let there be I items in a market-basket data set of B baskets. Suppose that every basket contains exactly K items. As a function of I , B , and K :

- a) How much space does the triangular-matrix method take to store the counts of all pairs of items, assuming four bytes per array element?
- b) What is the largest possible number of pairs with a nonzero count?
- c) Under what circumstances can we be certain that the triples method will use less space than the triangular array?