

The Basics of Bergman Spaces

Jacob White*
University of Nebraska Omaha

The Space $A^2(\Omega)$

Definition 1. Let $\Omega \subseteq \mathbb{C}$ be a **domain** (i.e., an open, connected set). Define

$$A^2(\Omega) = \left\{ f \text{ holomorphic on } \Omega : \int_{\Omega} |f(z)|^2 dA(z) < \infty \right\} \subseteq L^2(\Omega)$$

where dA is the ordinary two-dimensional area measure. Then, $A^2(\Omega)$ is a complex vector space, called the **Bergman space**.

- The **Bergman norm** is given by

$$\|f\|_{A^2(\Omega)} = \left[\int_{\Omega} |f(z)|^2 dA(z) \right]^{1/2}.$$

- The standard inner product on $A^2(\Omega)$ is defined by

$$\langle f, g \rangle = \int_{\Omega} f(z) \overline{g(z)} dA(z).$$

The next lemma provides an important estimate for functions in $A^2(\Omega)$.

Lemma 1. Let $K \subseteq \Omega$ be compact. There is a constant, $C_K > 0$, depending on K , such that

$$\sup_{z \in K} |f(z)| \leq C_K \|f\|_{A^2(\Omega)}$$

for all $f \in A^2(\Omega)$.

Before we prove it, we'll review a preliminary result.

*Contact: jwhite3@unomaha.edu... please let me know if you have questions, feedback, or ideas on how to improve these notes!
You can also make pull requests/issue requests/whatever how that works on the github repo for this document

Observation 1.

Proof.

