

The recent development of dependency graphs

Rui-Ray Zhang


`rui.zhang@monash.edu`

School of Mathematics, Monash University

Dependency Graphs

Graph G is a dependency graph for random variables $\mathbf{X} = (X_1, \dots, X_n)$ if


- Vertex set $V(G) = [n] = \{1, \dots, n\}$.



Dependency Graphs

Graph G is a dependency graph for random variables $\mathbf{X} = (X_1, \dots, X_n)$ if

- Vertex set $V(G) = [n] = \{1, \dots, n\}$.



- If disjoint subsets $I, J \subset [n]$ are non-adjacent in G , then $\{X_i\}_{i \in I}$ and $\{X_j\}_{j \in J}$ are independent.
 - ▶ In the above example, $\{X_1, X_2\}$ and $\{X_5, X_6\}$ are independent.
- There are ones with weaker assumptions, such as the one used in Lovász local lemma.
- The dependency graph for variables may not necessarily be unique, and the sparser ones are the more interesting ones.

Classics

- Concentration under graph-dependence.
 - ▶ Janson's concentration bound (2004)

Classics

- Concentration under graph-dependence.
 - ▶ Janson's concentration bound (2004)
- The probability of non-existence of certain structures under graph-dependence.
 - ▶ Lovász local lemma (1975).
 - ▶ Chen-Stein's method (1975).

Classics

- Concentration under graph-dependence.
 - ▶ Janson's concentration bound (2004)
- The probability of non-existence of certain structures under graph-dependence.
 - ▶ Lovász local lemma (1975).
 - ▶ Chen-Stein's method (1975).
 - ▶ Janson's inequality (1990).
 - ▶ Suen's inequality (1990).

Classics

- Concentration under graph-dependence.
 - ▶ Janson's concentration bound (2004)
- The probability of non-existence of certain structures under graph-dependence.
 - ▶ Lovász local lemma (1975).
 - ▶ Chen-Stein's method (1975).
 - ▶ Janson's inequality (1990).
 - ▶ Suen's inequality (1990).
- Limiting distribution via dependency graph.
 - ▶ Janson's normality criterion via cumulants (1988)

Concentration bounds

A McDiarmid-type bounded difference inequality for graph-dependent variables.

Theorem (Z., Liu, Wang, Wang, 2019)

If we have

- 1 f is c -Lipschitz,
- 2 F is a dependency graph for \mathbf{X} and $F = \{T_i\}_{i \in [k]}$ is a forest,

Concentration bounds

A McDiarmid-type bounded difference inequality for graph-dependent variables.

Theorem (Z., Liu, Wang, Wang, 2019)

If we have

- 1 f is \mathbf{c} -Lipschitz,
- 2 F is a dependency graph for \mathbf{X} and $F = \{T_i\}_{i \in [k]}$ is a forest,

then for every $t > 0$,

$$\mathbb{P}(f(\mathbf{X}) - \mathbb{E}[f(\mathbf{X})] \geq t) \leq \exp\left(-\frac{2t^2}{\sum_{i=1}^k c_{\min,i}^2 + \sum_{\{i,j\} \in E(F)} (c_i + c_j)^2}\right),$$

where $c_{\min,i} = \min\{c_j : j \in V(T_i)\}$ is the minimum entry of \mathbf{c} in each tree T_i .

Concentration bounds

Janson, 2004

Let random variables $\{X_i\}_{i \in V(G)}$ be G -dependent such that every X_i takes values in an interval of length $c_i \geq 0$. Then, for every $t > 0$,

$$\mathbb{P}\left(\sum_{i \in V(G)} X_i - \mathbb{E}\left[\sum_{i \in V(G)} X_i\right] \geq t\right) \leq \exp\left(-\frac{2t^2}{\chi_f(G) \sum_{i \in V(G)} c_i^2}\right),$$

where $\chi_f(G)$ is the fractional chromatic number of G .

Concentration bounds

Janson, 2004

Let random variables $\{X_i\}_{i \in V(G)}$ be G -dependent such that every X_i takes values in an interval of length $c_i \geq 0$. Then, for every $t > 0$,

$$\mathbb{P}\left(\sum_{i \in V(G)} X_i - \mathbb{E}\left[\sum_{i \in V(G)} X_i\right] \geq t\right) \leq \exp\left(-\frac{2t^2}{\chi_f(G) \sum_{i \in V(G)} c_i^2}\right),$$

where $\chi_f(G)$ is the fractional chromatic number of G .

Z., 2022

Under the same setting,

$$\mathbb{P}\left(\sum_{i \in V(G)} X_i - \mathbb{E}\left[\sum_{i \in V(G)} X_i\right] \geq t\right) \leq \exp\left(-\frac{2t^2}{D}\right),$$

where D is optimised over fractional forest vertex covers of G .

Concentration bounds

Janson, 2004

Let random variables $\{X_i\}_{i \in V(G)}$ be G -dependent such that every X_i takes values in an interval of length $c_i \geq 0$. Then, for every $t > 0$,

$$\mathbb{P}\left(\sum_{i \in V(G)} X_i - \mathbb{E}\left[\sum_{i \in V(G)} X_i\right] \geq t\right) \leq \exp\left(-\frac{2t^2}{\chi_f(G) \sum_{i \in V(G)} c_i^2}\right),$$

where $\chi_f(G)$ is the fractional chromatic number of G .

Z., 2022

Under the same setting,


$$\mathbb{P}\left(\sum_{i \in V(G)} X_i - \mathbb{E}\left[\sum_{i \in V(G)} X_i\right] \geq t\right) \leq \exp\left(-\frac{2t^2}{D}\right),$$

where D is optimised over fractional forest vertex covers of G .

- It is no worse than Janson's, better when G is sparse.
- It generalises to certain decomposable Lipschitz functions, extending McDiarmid's.

Example


Let $\{X_i\}_{i \in [9]}$ be random indicators with the dependency graph G , and $X = \sum_{i \in [9]} X_i$.



$$\mathbb{P}(X - \mathbb{E}[X] \geq t) \leq ?$$


Example

Let $\{X_i\}_{i \in [9]}$ be random indicators with the dependency graph G , and $X = \sum_{i \in [9]} X_i$.



$$\mathbb{P}(X - \mathbb{E}[X] \geq t) \leq ?$$


- Janson's idea: to cover vertices with weighted independent sets such that the sum of weights for each vertex equals 1



$$\mathbb{P}(X - \mathbb{E}[X] \geq t) \leq \exp\left(-\frac{2t^2}{3 \times 9}\right)$$


Example

Let $\{X_i\}_{i \in [9]}$ be random indicators with the dependency graph G , and $X = \sum_{i \in [9]} X_i$.




$$\mathbb{P}(X - \mathbb{E}[X] \geq t) \leq ?$$

- Janson's idea: to cover vertices with weighted independent sets such that the sum of weights for each vertex equals 1



$$\mathbb{P}(X - \mathbb{E}[X] \geq t) \leq \exp\left(-\frac{2t^2}{3 \times 9}\right)$$

- New idea: to cover vertices with weighted induced forests such that the sum of weights for each vertex equals 1



$$\mathbb{P}(X - \mathbb{E}[X] \geq t) \leq \exp\left(-\frac{8t^2}{81}\right)$$

The probability of non-occurrences

Let $\{X_i\}_i$ be G -dependent indicators for events and $X = \sum_i X_i$ count the occurrences of events (e.g., existence of certain combinatorial structure, say, Δ in $\mathcal{G}(n, p)$).

The probability of non-occurrences

Let $\{X_i\}_i$ be G -dependent indicators for events and $X = \sum_i X_i$ count the occurrences of events (e.g., existence of certain combinatorial structure, say, Δ in $\mathcal{G}(n, p)$).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[\prod_{i \in S} X_i \right],$$

for $S \subseteq V(G)$.

The probability of non-occurrences


Let $\{X_i\}_i$ be G -dependent indicators for events and $X = \sum_i X_i$ count the occurrences of events (e.g., existence of certain combinatorial structure, say, Δ in $\mathcal{G}(n, p)$).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[\prod_{i \in S} X_i \right],$$

for $S \subseteq V(G)$.

Then we have a factorization property



$$\begin{aligned} & \mu(1,3)\mu(2)\mu(4,7)\mu(5,6) \\ &= \mu(1)\mu(2)\mu(3)\mu(4)\mu(5,6)\mu(7). \end{aligned}$$

The probability of non-occurrences

Let $\{X_i\}_i$ be G -dependent indicators for events and $X = \sum_i X_i$ count the occurrences of events (e.g., existence of certain combinatorial structure, say, Δ in $\mathcal{G}(n, p)$).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[\prod_{i \in S} X_i \right],$$

for $S \subseteq V(G)$.

By the principle of inclusion–exclusion and the factorization, we have

$$\mathbb{P}(X = 0) = \mathbb{P} \left(\sum_i X_i = 0 \right)$$

The probability of non-occurrences

Let $\{X_i\}_i$ be G -dependent indicators for events and $X = \sum_i X_i$ count the occurrences of events (e.g., existence of certain combinatorial structure, say, Δ in $\mathcal{G}(n, p)$).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[\prod_{i \in S} X_i \right],$$

for $S \subseteq V(G)$.

By the principle of inclusion–exclusion and the factorization, we have

$$\mathbb{P}(X = 0) = \mathbb{P}\left(\sum_i X_i = 0\right) = \sum_{S \subseteq V(G)} (-1)^{|S|} \mu(S)$$

The probability of non-occurrences

Let $\{X_i\}_i$ be G -dependent indicators for events and $X = \sum_i X_i$ count the occurrences of events (e.g., existence of certain combinatorial structure, say, Δ in $\mathcal{G}(n, p)$).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[\prod_{i \in S} X_i \right],$$

for $S \subseteq V(G)$.

By the principle of inclusion–exclusion and the factorization, we have

$$\mathbb{P}(X = 0) = \mathbb{P} \left(\sum_i X_i = 0 \right) = \sum_{S \subseteq V(G)} (-1)^{|S|} \mu(S) = \sum_{S \subseteq V(G)} (-1)^{|S|} \mu(C_1) \mu(C_2) \dots \mu(C_k),$$

where $C_1 \cup \dots \cup C_k = S$ and C_1, \dots, C_k induce maximal pairwise non-adjacent 'polymers' (that is, connected subgraphs; also called 'animals', suggested by Dobrushin, 1996).

The exponential formula reduces the sum to connected subgraphs
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

The exponential formula reduces the sum to connected subgraphs
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

The exponential formula reduces the sum to connected subgraphs
 (idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs): $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$.
- Clusters: $\dots, \text{connected union of polymers}, \dots$

The exponential formula reduces the sum to connected subgraphs
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs): $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$.
- **Clusters**: $\dots, \text{connected union of polymers}, \dots$
- Ursell function ϕ relates to chromatic polynomials.

The exponential formula reduces the sum to connected subgraphs
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs): $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$.
- **Clusters**: $\dots, \text{connected union of polymers}, \dots$
- Ursell function ϕ relates to chromatic polynomials.
- Cluster expansion (Mayer, 1941) of partition function of a hard-core model with activity of each polymer C being $(-1)^{|C|} \mu(C)$.

The exponential formula reduces the sum to connected subgraphs
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs): $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$.
- **Clusters**: $\dots, \text{connected union of polymers}, \dots$
- Ursell function ϕ relates to chromatic polynomials.
- Cluster expansion (Mayer, 1941) of partition function of a hard-core model with activity of each polymer C being $(-1)^{|C|} \mu(C)$.
- Scott and Sokal (2005) observed that
 - ▶ the Lovász local lemma (1975) condition (equivalently, Dobrushin's criterion, 1996) gives an absolute convergence criterion;

The exponential formula reduces the sum to connected subgraphs
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs): $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$.
- **Clusters**: $\dots, \text{connected union of polymers}, \dots$
- Ursell function ϕ relates to chromatic polynomials.
- Cluster expansion (Mayer, 1941) of partition function of a hard-core model with activity of each polymer C being $(-1)^{|C|} \mu(C)$.
- Scott and Sokal (2005) observed that
 - ▶ the Lovász local lemma (1975) condition (equivalently, Dobrushin's criterion, 1996) gives an absolute convergence criterion;
 - ▶ the tight instance for lower bound is when every polymer is K_1 (also by Shearer, 1985).

$$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$$

X counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if $\mathbb{E} \left[\triangle \right] = np^2 = o(1)$, then X is asymptotically Poisson.

$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

X counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if $\mathbb{E} \left[\text{triangle} \right] = np^2 = o(1)$, then X is asymptotically Poisson.
- Janson-Łuczak-Ruciński, 1990 (July 1987, Poznan):

$$\exp \left(- \frac{\mathbb{E} \left[\text{triangle} \right]}{1 - p^3} \right) \stackrel{\text{FKG}}{\leq} \mathbb{P}(X = 0) \stackrel{\text{Janson's inequality}}{\leq} \exp \left(- \mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{diamond} \right] \right),$$

$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

X counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if $\mathbb{E} \left[\text{triangle} \right] = np^2 = o(1)$, then X is asymptotically Poisson.
- Janson-Łuczak-Ruciński, 1990 (July 1987, Poznan):

$$\exp \left(- \frac{\mathbb{E} \left[\text{triangle} \right]}{1 - p^3} \right) \stackrel{\text{FKG}}{\leq} \mathbb{P}(X = 0) \stackrel{\text{Janson's inequality}}{\leq} \exp \left(- \mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{diamond} \right] \right),$$

if $\mathbb{E} \left[\text{diamond} \right] = o(1)$, then

$$\mathbb{P}(X = 0) = \exp \left(- \mathbb{E} \left[\text{triangle} \right] + o(1) \right).$$

$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

X counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if $\mathbb{E} \left[\text{triangle} \right] = np^2 = o(1)$, then X is asymptotically Poisson.
- Janson-Łuczak-Ruciński, 1990 (July 1987, Poznan):

$$\exp \left(- \frac{\mathbb{E} \left[\text{triangle} \right]}{1 - p^3} \right) \stackrel{\text{FKG}}{\leq} \mathbb{P}(X = 0) \stackrel{\text{Janson's inequality}}{\leq} \exp \left(- \mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{two triangles sharing an edge} \right] \right),$$

if $\mathbb{E} \left[\text{two triangles sharing an edge} \right] = o(1)$, then

$$\mathbb{P}(X = 0) = \exp \left(- \mathbb{E} \left[\text{triangle} \right] + o(1) \right).$$

- Wormald, 1996: if $p = o(n^{-2/3})$, then

$$\mathbb{P}(X = 0) = \exp \left(- \mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{two triangles sharing an edge} \right] - \mathbb{E} \left[\text{three triangles sharing a vertex} \right] - \mathbb{E} \left[\text{two triangles sharing two edges} \right] + o(1) \right).$$

$$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$$

$$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$$

$$\begin{aligned} \log \mathbb{P}(X=0) &\stackrel{\text{formally}}{=} -\mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{path of length 4} \right] - \mathbb{E} \left[\text{pentagon} \right] - \mathbb{E} \left[\text{house} \right] - \mathbb{E} \left[\text{triangle with a pendant edge} \right] \\ &+ \mathbb{E} \left[\text{complex graph 1} \right] + \mathbb{E} \left[\text{complex graph 2} \right] + \mathbb{E} \left[\text{complex graph 3} \right] + \mathbb{E} \left[\text{complex graph 4} \right] + \mathbb{E} \left[\text{complex graph 5} \right] + \mathbb{E} \left[\text{triangle} \right] \\ &- \tilde{\mathbb{E}} \left[\text{triangle with red edges} \right] - \tilde{\mathbb{E}} \left[\text{path of length 4 with red edges} \right] + \tilde{\mathbb{E}} \left[\text{pentagon with red edges} \right] + \dots \end{aligned}$$

$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

$$\begin{aligned} \log \mathbb{P}(X=0) &\stackrel{\text{formally}}{=} -\mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{path of length 4} \right] - \mathbb{E} \left[\text{cycle of length 5} \right] - \mathbb{E} \left[\text{cycle of length 6} \right] - \mathbb{E} \left[\text{triangle with a pendant edge} \right] \\ &+ \mathbb{E} \left[\text{cycle of length 7} \right] + \mathbb{E} \left[\text{cycle of length 8} \right] + \mathbb{E} \left[\text{cycle of length 9} \right] + \mathbb{E} \left[\text{cycle of length 10} \right] + \mathbb{E} \left[\text{cycle of length 11} \right] + \mathbb{E} \left[\text{triangle with two pendant edges} \right] \\ &- \tilde{\mathbb{E}} \left[\text{triangle with one red edge} \right] - \tilde{\mathbb{E}} \left[\text{triangle with two red edges} \right] + \tilde{\mathbb{E}} \left[\text{triangle with three red edges} \right] + \dots \end{aligned}$$

Stark and Wormald, 2018; Mousset, Noever, Panagiotou, Samotij, 2020

If $p = o(n^{-7/11})$, then

$$\begin{aligned} \mathbb{P}(X=0) &= \exp \left(-\mathbb{E} \left[\text{triangle} \right] + \mathbb{E} \left[\text{path of length 4} \right] - \mathbb{E} \left[\text{cycle of length 5} \right] - \mathbb{E} \left[\text{cycle of length 6} \right] - \mathbb{E} \left[\text{triangle with a pendant edge} \right] \right. \\ &\left. + \mathbb{E} \left[\text{cycle of length 7} \right] + \mathbb{E} \left[\text{cycle of length 8} \right] + \mathbb{E} \left[\text{cycle of length 9} \right] + \mathbb{E} \left[\text{cycle of length 10} \right] + \mathbb{E} \left[\text{cycle of length 11} \right] + \mathbb{E} \left[\text{triangle with two pendant edges} \right] + o(1) \right). \end{aligned}$$

- Mousset, Noever, Panagiotou, Samotij used [an alternating cumulant series](#), whose terms after simplifications are responding to partitions in cluster expansion.

- Mousset, Noever, Panagiotou, Samotij used [an alternating cumulant series](#), whose terms after simplifications are responding to partitions in cluster expansion.

Joint cumulant (1929)

For any multiset of random variables $\{X_i\}_{i \in S}$,

$$\kappa(S) = \sum_{\pi \in \Pi(S)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{P \in \pi} \mu(P).$$

Also called *semi-invariant* (1889) in probability, *Ursell function* (1927) in physics.

- Mousset, Noever, Panagiotou, Samotij used **an alternating cumulant series**, whose terms after simplifications are responding to partitions in cluster expansion.

Joint cumulant (1929)

For any multiset of random variables $\{X_i\}_{i \in S}$,

$$\kappa(S) = \sum_{\pi \in \Pi(S)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{P \in \pi} \mu(P).$$

Also called *semi-invariant* (1889) in probability, *Ursell function* (1927) in physics.

For G -dependent variables and $S \subseteq V(G)$,

if S **does not induce a connected subgraph** in G , then $\kappa(S) = 0$.

- Isaev, McKay, Z. (2022+):
accurate asymptotic enumeration of Eulerian orientations, digraphs, oriented graphs.

- Mousset, Noever, Panagiotou, Samotij used an alternating cumulant series, whose terms after simplifications are responding to partitions in cluster expansion.

Joint cumulant (1929)

For any multiset of random variables $\{X_i\}_{i \in S}$,

$$\kappa(S) = \sum_{\pi \in \Pi(S)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{P \in \pi} \mu(P).$$

Also called *semi-invariant* (1889) in probability, *Ursell function* (1927) in physics.

For G -dependent variables and $S \subseteq V(G)$,

if S does not induce a connected subgraph in G , then $\kappa(S) = 0$.

- Isaev, McKay, Z. (2022+):
accurate asymptotic enumeration of Eulerian orientations, digraphs, oriented graphs.
- Stark and Wormald (2018) also have results for $\mathcal{G}_{n,m}$:
if $d = m \binom{n}{2}^{-1} = o(n^{-7/11})$, then

$$\mathbb{P}(X=0) = \exp\left(-\frac{1}{6}n^3d^3 - \frac{1}{8}n^4d^6 - \frac{1}{2}n^2d^3 + o(1)\right).$$

Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex.

Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$\mathbb{P}(\text{a random hypergraph is linear})$

$= \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).$

Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$\mathbb{P}(\text{a random hypergraph is linear})$

$= \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).$

McKay and Tian, 2020

If $p = o(n^{-3/2})$, then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 p^2 + \frac{2}{3}n^5 p^3 + o(1)\right).$$

Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$$\begin{aligned}\mathbb{P}(\text{a random hypergraph is linear}) \\ = \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).\end{aligned}$$

McKay and Tian, 2020

If $p = o(n^{-3/2})$, then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4p^2 + \frac{2}{3}n^5p^3 + o(1)\right).$$

Z. 2022; Wormald and Z., 2022+

If $p = o(n^{-7/5})$, then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4p^2 + \frac{2}{3}n^5p^3 - \frac{55}{24}n^6p^4 + \frac{3}{2}n^3p^2 + o(1)\right).$$

- In Z. 2022, the truncation error is by Mousset, Noever, Panagiotou, Samotij's results.
- Coefficients match McKay's conjecture based upon numerical simulation.

Asymptotic linearity of hypergraphs

McKay and Tian, 2020

If $d = \binom{n}{3}^{-1} m = o(n^{-3/2})$, then via switching method,

$$\mathbb{P}(H_3(n, m) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 d^2 - \frac{1}{12}n^5 d^3 + o(1)\right).$$

Asymptotic linearity of hypergraphs

McKay and Tian, 2020

If $d = \binom{n}{3}^{-1} m = o\left(n^{-3/2}\right)$, then via switching method,

$$\mathbb{P}(H_3(n, m) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 d^2 - \frac{1}{12}n^5 d^3 + o(1)\right).$$

Wormald and Z. 2022+

If $d = \binom{n}{3}^{-1} m = o\left(n^{-7/5}\right)$, then via perturbation method,

$$\mathbb{P}(H_3(n, m) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 d^2 - \frac{1}{12}n^5 d^3 - \frac{1}{24}n^6 d^4 + \frac{3}{2}n^3 d^2 + o(1)\right).$$

Dependency digraphs

Dependency digraphs

A digraph $\mathbf{D} = ([d], E)$ is a dependency digraph for the events $\{A_i\}_{i \in [d]}$ if A_i is mutually independent of all the non-adjacent $\{A_j : (i, j) \notin E\}$ for every $i \in [d]$.


Dependency digraphs

Dependency digraphs

A digraph $\mathbf{D} = ([d], E)$ is a dependency digraph for the events $\{A_i\}_{i \in [d]}$ if A_i is mutually independent of all the non-adjacent $\{A_j : (i, j) \notin E\}$ for every $i \in [d]$.

Example

Let $\{A_i\}_{i \in [5]}$ be events with the following dependency digraph.




A_1 is independent of A_4 .

A_2 is independent of $\{A_1, A_3, A_5\}$.

A natural refinement is

- adjacent vertices \rightarrow the pairs of 'strongly dependent' events,
 $D_i \subseteq [d]$ denotes the 'strongly dependent neighbours' of event A_i .
- non-adjacent vertices \rightarrow the pairs of 'weakly dependent' events.




$$\mathbb{P}(A_4 | A_1) \approx \mathbb{P}(A_4)$$

$$\mathbb{P}(A_1 \cup A_3 \cup A_5 | A_2) \approx \mathbb{P}(A_1 \cup A_3 \cup A_5)$$

A natural refinement is

- adjacent vertices \rightarrow the pairs of 'strongly dependent' events,
 $D_i \subseteq [d]$ denotes the 'strongly dependent neighbours' of event A_i .
- non-adjacent vertices \rightarrow the pairs of 'weakly dependent' events.



$$\mathbb{P}(A_4 | A_1) \approx \mathbb{P}(A_4)$$

$$\mathbb{P}(A_1 \cup A_3 \cup A_5 | A_2) \approx \mathbb{P}(A_1 \cup A_3 \cup A_5)$$

If we change ' \approx ' to ' \geq ', then it is the 'negative dependency graph' by Erdős and Spencer (1991) to obtain lopsided Lovász local lemma.

New dependency digraphs

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} on $[d]$:

New dependency digraphs

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} on $[d]$:

- φ -mixing coefficient is used to measure 'weak dependencies':

$$\varphi = \max_{i \in [d]} \left| \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \mid A_i \right) - \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \right) \right|.$$

New dependency digraphs

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} on $[d]$:

- φ -mixing coefficient is used to measure 'weak dependencies':

$$\varphi = \max_{i \in [d]} \left| \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \mid A_i \right) - \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \right) \right|.$$

- The influence of 'strongly dependent' events is

$$\Delta_1 = \sum_{i \in [d]} \mathbb{P} \left(A_i \cap \bigcup_{j \in [i-1] \cap D_i} A_j \right) \prod_{k \in [d] \setminus [i]} \mathbb{P}(\overline{A_k})$$

$$\Delta_2 = \sum_{i \in [d]} \mathbb{P}(A_i) \mathbb{P} \left(\bigcup_{j \in [i-1] \cap D_i} A_j \right) \prod_{k \in [d] \setminus [i]} \mathbb{P}(\overline{A_k}).$$

New dependency digraphs

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} on $[d]$:

- φ -mixing coefficient is used to measure 'weak dependencies'

$$\varphi = \max_{i \in [d]} \left| \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \mid A_i \right) - \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \right) \right|.$$

- The influence of 'strongly dependent' events is

$$\Delta_1 \leq \Delta'_1 = \sum_{i \in [d]} \sum_{j \in [i-1] \cap D_i} \mathbb{P}(A_i \cap A_j),$$
$$\Delta_2 \leq \Delta'_2 = \sum_{i \in [d]} \sum_{j \in [i-1] \cap D_i} \mathbb{P}(A_i) \mathbb{P}(A_j).$$

New dependency digraphs

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} on $[d]$:

- φ -mixing coefficient is used to measure 'weak dependencies'

$$\varphi = \max_{i \in [d]} \left| \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \mid A_i \right) - \mathbb{P} \left(\bigcup_{j \in [i-1] \setminus D_i} A_j \right) \right|.$$

- The influence of 'strongly dependent' events is

$$\Delta_1 \leq \Delta'_1 = \sum_{i \in [d]} \sum_{j \in [i-1] \cap D_i} \mathbb{P}(A_i \cap A_j),$$
$$\Delta_2 \leq \Delta'_2 = \sum_{i \in [d]} \sum_{j \in [i-1] \cap D_i} \mathbb{P}(A_i) \mathbb{P}(A_j).$$

- Chen–Stein Poisson approximation involves similar terms as φ , Δ'_1 , and Δ'_2 .
Chen (1975) also considered the φ -mixing condition.
- Suen's inequality (1990) also involves Δ'_1 and Δ'_2 .

Theorem (Isaev, Rodionov, Z., Zhukovskii; 2020, 2022+)

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} with vertex set $[d]$, we have

$$\left| \mathbb{P} \left(\bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left(1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.

Theorem (Isaev, Rodionov, Z., Zhukovskii; 2020, 2022+)

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} with vertex set $[d]$, we have

$$\left| \mathbb{P} \left(\bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left(1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If $\varphi = o(1)$, $\Delta_1 = o(1)$, and $\Delta_2 = o(1)$, then we have the asymptotic independence.

Theorem (Isaev, Rodionov, Z., Zhukovskii; 2020, 2022+)

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} with vertex set $[d]$, we have

$$\left| \mathbb{P} \left(\bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left(1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If $\varphi = o(1)$, $\Delta_1 = o(1)$, and $\Delta_2 = o(1)$, then we have the asymptotic independence.
 - ▶ 'Two Moments Suffice for Poisson Approximations: The Chen–Stein Method' by Arratia, Goldstein and Gordon (1989); also Chen (1975).

Theorem (Isaev, Rodionov, Z., Zhukovskii; 2020, 2022+)

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} with vertex set $[d]$, we have

$$\left| \mathbb{P} \left(\bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left(1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If $\varphi = o(1)$, $\Delta_1 = o(1)$, and $\Delta_2 = o(1)$, then we have the asymptotic independence.
 - ▶ 'Two Moments Suffice for Poisson Approximations: The Chen–Stein Method' by Arratia, Goldstein and Gordon (1989); also Chen (1975).
- Max codegrees, max clique-extension counts of random (hyper)graphs, etc. are asymptotically Gumbel.

Theorem (Isaev, Rodionov, Z., Zhukovskii; 2020, 2022+)

For any events $\{A_i\}_{i \in [d]}$ and digraph \mathbf{D} with vertex set $[d]$, we have

$$\left| \mathbb{P} \left(\bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left(1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If $\varphi = o(1)$, $\Delta_1 = o(1)$, and $\Delta_2 = o(1)$, then we have the asymptotic independence.
 - ▶ 'Two Moments Suffice for Poisson Approximations: The Chen–Stein Method' by Arratia, Goldstein and Gordon (1989); also Chen (1975).
- Max codegrees, max clique-extension counts of random (hyper)graphs, etc. are asymptotically Gumbel.
- Lu and Székely (2009) also introduced a notion of 'positive' dependency graph to obtain upper bounds for Poisson approximation for random matchings.

Open problem

- To establish the absolute convergence of cluster expansion.

Open problem

- To establish the absolute convergence of cluster expansion.
- To design a suitable model for $\mathcal{G}(n, m)$;
e.g., Féray's weighted dependency graphs for normality of subgraph counts.

Thanks!