

# The pursuit of more accurate asymptotics <sup>1</sup> via clusters <sup>2</sup> (the three-year milestone).

Rui-Ray Zhang

[rui.zhang@monash.edu](mailto:rui.zhang@monash.edu)

School of Mathematics, Monash University

Supervisors: Nick Wormald and Mikhail Isaev.

---

<sup>1</sup>Asymptotics are describing limiting behaviours as  $n \rightarrow \infty$ .

<sup>2</sup>Clusters are connected structures.

This talk covers

- The asymptotic probability of non-existence:

This talk covers

- The asymptotic probability of non-existence:
  - ▶ asymptotic linearity of binomial random hypergraphs via cluster expansion,

This talk covers

- The asymptotic probability of non-existence:
  - ▶ asymptotic linearity of binomial random hypergraphs via cluster expansion,
  - ▶ asymptotic linearity of random hypergraphs via perturbation method (with Wormald),

This talk covers

- The asymptotic probability of non-existence:
  - ▶ asymptotic linearity of binomial random hypergraphs via cluster expansion,
  - ▶ asymptotic linearity of random hypergraphs via perturbation method (with Wormald),
  - ▶ asymptotic independence under mixing via dependency digraphs (with Isaev, Rodionov, Zhukovskii),

This talk covers

- The asymptotic probability of non-existence:
  - ▶ asymptotic linearity of binomial random hypergraphs via cluster expansion,
  - ▶ asymptotic linearity of random hypergraphs via perturbation method (with Wormald),
  - ▶ asymptotic independence under mixing via dependency digraphs (with Isaev, Rodionov, Zhukovskii),
- The asymptotic enumeration:

This talk covers

- The asymptotic probability of non-existence:
  - ▶ asymptotic linearity of binomial random hypergraphs via cluster expansion,
  - ▶ asymptotic linearity of random hypergraphs via perturbation method (with Wormald),
  - ▶ asymptotic independence under mixing via dependency digraphs (with Isaev, Rodionov, Zhukovskii),
- The asymptotic enumeration:
  - ▶ asymptotic enumeration of regular tournaments, Eulerian digraphs, and Eulerian oriented graphs via cumulants (with Isaev, McKay).

## Dependency graphs

A graph  $G$  is a dependency graph for random variables  $\{X_i\}_{i \in V(G)}$  if variables  $\{X_i\}_{i \in I}$  and  $\{X_j\}_{j \in J}$  are independent for any disjoint non-adjacent  $I, J \subset V(G)$ .




## Dependency graphs

A graph  $G$  is a dependency graph for random variables  $\{X_i\}_{i \in V(G)}$  if variables  $\{X_i\}_{i \in I}$  and  $\{X_j\}_{j \in J}$  are independent for any disjoint non-adjacent  $I, J \subset V(G)$ .

## Example

Let  $\{X_i\}_{i \in [6]}$  be random variables with the following dependency graph  $G$ .



Variables  $\{X_1, X_2\}$  and  $\{X_5, X_6\}$  are independent, since disjoint vertex sets  $\{1, 2\}$  and  $\{5, 6\}$  are not adjacent in  $G$ .

## The probability of non-occurrences

Let  $\{X_i\}_i$  be  $G$ -dependent indicators for events and  $X = \sum_i X_i$  count the occurrences of events (e.g., existence of certain combinatorial structure, say,  $\Delta$  in  $\mathcal{G}(n, p)$ ).

## The probability of non-occurrences

Let  $\{X_i\}_i$  be  $G$ -dependent indicators for events and  $X = \sum_i X_i$  count the occurrences of events (e.g., existence of certain combinatorial structure, say,  $\Delta$  in  $\mathcal{G}(n, p)$ ).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[ \prod_{i \in S} X_i \right],$$

for  $S \subseteq V(G)$ .

## The probability of non-occurrences

Let  $\{X_i\}_i$  be  $G$ -dependent indicators for events and  $X = \sum_i X_i$  count the occurrences of events (e.g., existence of certain combinatorial structure, say,  $\Delta$  in  $\mathcal{G}(n, p)$ ).

Let the probability of joint occurrence denote by

$$\mu(S) = \mathbb{E} \left[ \prod_{i \in S} X_i \right],$$

for  $S \subseteq V(G)$ . By the principle of inclusion–exclusion and the factorization, we have

$$\mathbb{P}(X = 0) = \mathbb{P} \left( \sum_i X_i = 0 \right) = \sum_{S \subseteq V(G)} (-1)^{|S|} \mu(S) = \sum_{S \subseteq V(G)} (-1)^{|S|} \mu(C_1) \mu(C_2) \dots \mu(C_k),$$

where  $C_1 \cup \dots \cup C_k = S$  and  $C_1, \dots, C_k$  induce maximal pairwise non-adjacent ‘polymers’ (that is, connected subgraphs; also called ‘animals’, suggested by Dobrushin, 1996).

The exponential formula reduces the sum to connected subgraphs  
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

### Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

The exponential formula reduces the sum to connected subgraphs  
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

### Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs):  $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$ .
- **Clusters**:  $\dots, \text{connected union of polymers}, \dots$
- Ursell function  $\phi$  relates to chromatic polynomials.

The exponential formula reduces the sum to connected subgraphs  
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

### Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs):  $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$ .
- **Clusters**:  $\dots, \text{connected union of polymers}, \dots$
- Ursell function  $\phi$  relates to chromatic polynomials.
- Cluster expansion (Mayer, 1941) of partition function of a hard-core model with activity of each polymer  $C$  being  $(-1)^{|C|} \mu(C)$ .

The exponential formula reduces the sum to connected subgraphs  
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

### Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs):  $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$ .
- **Clusters**:  $\dots, \text{connected union of polymers}, \dots$
- Ursell function  $\phi$  relates to chromatic polynomials.
- Cluster expansion (Mayer, 1941) of partition function of a hard-core model with activity of each polymer  $C$  being  $(-1)^{|C|} \mu(C)$ .
- Scott and Sokal (2005) observed that
  - ▶ the Lovász local lemma (1975) condition (equivalently, Dobrushin's criterion, 1996) gives an absolute convergence criterion;



The exponential formula reduces the sum to connected subgraphs  
(idea: any simple graph is the vertex-disjoint union of connected simple graphs), we have

### Formal cluster expansion (Z. 2022)

$$\log(\mathbb{P}(X=0)) \stackrel{\text{formally}}{=} \sum_{k \geq 1} \sum_{(C_1, \dots, C_k)} \phi(C_1, \dots, C_k) \prod_{C \in (C_1, \dots, C_k)} (-1)^{|C|} \mu(C),$$

where the sum is over **ordered multisets of polymers, whose union is connected**.

- Polymers (connected subgraphs):  $\{\dots, \text{triangle}, \text{V-shape}, \text{square}, \dots\}$ .
- **Clusters**:  $\dots, \text{connected union of polymers}, \dots$
- Ursell function  $\phi$  relates to chromatic polynomials.
- Cluster expansion (Mayer, 1941) of partition function of a hard-core model with activity of each polymer  $C$  being  $(-1)^{|C|} \mu(C)$ .
- Scott and Sokal (2005) observed that
  - ▶ the Lovász local lemma (1975) condition (equivalently, Dobrushin's criterion, 1996) gives an absolute convergence criterion;
  - ▶ the tight instance for lower bound is when every polymer is  $K_1$  (also by Shearer, 1985).

$$\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$$

$X$  counts triangles in  $\mathcal{G}(n, p)$

- Ruciński, 1988: if  $\mathbb{E} \left[ \triangle \right] = np^2 = o(1)$ , then  $X$  is asymptotically Poisson.

## $\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

### $X$ counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if  $\mathbb{E} \left[ \text{triangle} \right] = np^2 = o(1)$ , then  $X$  is asymptotically Poisson.
- Janson-Łuczak-Ruciński, 1990 (July 1987, Poznan):

$$\exp \left( - \frac{\mathbb{E} \left[ \text{triangle} \right]}{1 - p^3} \right) \stackrel{\text{FKG}}{\leq} \mathbb{P}(X = 0) \stackrel{\text{Janson's inequality}}{\leq} \exp \left( - \mathbb{E} \left[ \text{triangle} \right] + \mathbb{E} \left[ \text{diamond} \right] \right),$$

# $\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

## $X$ counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if  $\mathbb{E} \left[ \text{triangle} \right] = np^2 = o(1)$ , then  $X$  is asymptotically Poisson.
- Janson-Łuczak-Ruciński, 1990 (July 1987, Poznan):

$$\exp \left( - \frac{\mathbb{E} \left[ \text{triangle} \right]}{1 - p^3} \right) \stackrel{\text{FKG}}{\leq} \mathbb{P}(X = 0) \stackrel{\text{Janson's inequality}}{\leq} \exp \left( - \mathbb{E} \left[ \text{triangle} \right] + \mathbb{E} \left[ \text{4-cycle} \right] \right),$$

if  $\mathbb{E} \left[ \text{4-cycle} \right] = o(1)$ , then

$$\mathbb{P}(X = 0) = \exp \left( - \mathbb{E} \left[ \text{triangle} \right] + o(1) \right).$$

# $\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

## $X$ counts triangles in $\mathcal{G}(n, p)$

- Ruciński, 1988: if  $\mathbb{E} \left[ \text{triangle} \right] = np^2 = o(1)$ , then  $X$  is asymptotically Poisson.
- Janson-Łuczak-Ruciński, 1990 (July 1987, Poznan):

$$\exp \left( - \frac{\mathbb{E} \left[ \text{triangle} \right]}{1 - p^3} \right) \stackrel{\text{FKG}}{\leq} \mathbb{P}(X = 0) \stackrel{\text{Janson's inequality}}{\leq} \exp \left( - \mathbb{E} \left[ \text{triangle} \right] + \mathbb{E} \left[ \text{two triangles sharing an edge} \right] \right),$$

if  $\mathbb{E} \left[ \text{two triangles sharing an edge} \right] = o(1)$ , then

$$\mathbb{P}(X = 0) = \exp \left( - \mathbb{E} \left[ \text{triangle} \right] + o(1) \right).$$

- Wormald, 1996: if  $p = o(n^{-2/3})$ , then

$$\mathbb{P}(X = 0) = \exp \left( - \mathbb{E} \left[ \text{triangle} \right] + \mathbb{E} \left[ \text{two triangles sharing an edge} \right] - \mathbb{E} \left[ \text{three triangles sharing a vertex} \right] - \mathbb{E} \left[ \text{three triangles sharing an edge} \right] + o(1) \right).$$

# $\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

$$\begin{aligned} \log \mathbb{P}(X=0) &\stackrel{\text{formally}}{=} -\mathbb{E} \left[ \text{triangle} \right] + \mathbb{E} \left[ \text{path of length 4} \right] - \mathbb{E} \left[ \text{pentagon} \right] - \mathbb{E} \left[ \text{house} \right] - \mathbb{E} \left[ \text{triangle with a pendant edge} \right] \\ &+ \mathbb{E} \left[ \text{complex graph 1} \right] + \mathbb{E} \left[ \text{complex graph 2} \right] + \mathbb{E} \left[ \text{complex graph 3} \right] + \mathbb{E} \left[ \text{complex graph 4} \right] + \mathbb{E} \left[ \text{complex graph 5} \right] + \mathbb{E} \left[ \text{triangle} \right] \\ &- \tilde{\mathbb{E}} \left[ \text{triangle with red edges} \right] - \tilde{\mathbb{E}} \left[ \text{triangle with red edges} \right] + \tilde{\mathbb{E}} \left[ \text{triangle with red edges} \right] + \dots \end{aligned}$$

# $\mathbb{P}(\text{no triangles in } \mathcal{G}(n, p))$

$$\begin{aligned} \log \mathbb{P}(X=0) &\stackrel{\text{formally}}{=} -\mathbb{E} \left[ \text{graph}_1 \right] + \mathbb{E} \left[ \text{graph}_2 \right] - \mathbb{E} \left[ \text{graph}_3 \right] - \mathbb{E} \left[ \text{graph}_4 \right] - \mathbb{E} \left[ \text{graph}_5 \right] \\ &+ \mathbb{E} \left[ \text{graph}_6 \right] + \mathbb{E} \left[ \text{graph}_7 \right] + \mathbb{E} \left[ \text{graph}_8 \right] + \mathbb{E} \left[ \text{graph}_9 \right] + \mathbb{E} \left[ \text{graph}_{10} \right] + \mathbb{E} \left[ \text{graph}_{11} \right] \\ &- \tilde{\mathbb{E}} \left[ \text{graph}_{12} \right] - \tilde{\mathbb{E}} \left[ \text{graph}_{13} \right] + \tilde{\mathbb{E}} \left[ \text{graph}_{14} \right] + \dots \end{aligned}$$

**Stark and Wormald, 2018; Mousset, Noever, Panagiotou, Samotij, 2020**

If  $p = o(n^{-7/11})$ , then

$$\begin{aligned} \mathbb{P}(X=0) &= \exp \left( -\mathbb{E} \left[ \text{graph}_1 \right] + \mathbb{E} \left[ \text{graph}_2 \right] - \mathbb{E} \left[ \text{graph}_3 \right] - \mathbb{E} \left[ \text{graph}_4 \right] - \mathbb{E} \left[ \text{graph}_5 \right] \right. \\ &\left. + \mathbb{E} \left[ \text{graph}_6 \right] + \mathbb{E} \left[ \text{graph}_7 \right] + \mathbb{E} \left[ \text{graph}_8 \right] + \mathbb{E} \left[ \text{graph}_9 \right] + \mathbb{E} \left[ \text{graph}_{10} \right] + \mathbb{E} \left[ \text{graph}_{11} \right] + o(1) \right). \end{aligned}$$

- Mousset, Noever, Panagiotou, Samotij used [an alternating cumulant series](#), whose terms after simplifications are responding to partitions in cluster expansion.



- Mousset, Noever, Panagiotou, Samotij used [an alternating cumulant series](#), whose terms after simplifications are responding to partitions in cluster expansion.

## Joint cumulant (1929)

For any multiset of random variables  $\{X_i\}_{i \in S}$ ,

$$\kappa(S) = \sum_{\pi \in \Pi(S)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{P \in \pi} \mu(P).$$

Also called *semi-invariant* (1889) in probability, *Ursell function* (1927) in physics.

- Mousset, Noever, Panagiotou, Samotij used [an alternating cumulant series](#), whose terms after simplifications are responding to partitions in cluster expansion.

## Joint cumulant (1929)

For any multiset of random variables  $\{X_i\}_{i \in S}$ ,

$$\kappa(S) = \sum_{\pi \in \Pi(S)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{P \in \pi} \mu(P).$$

Also called *semi-invariant* (1889) in probability, *Ursell function* (1927) in physics.  
For  $G$ -dependent variables and  $S \subseteq V(G)$ ,  
if  $S$  [does not induce a connected subgraph](#) in  $G$ , then  $\kappa(S) = 0$ .

- Mousset, Noever, Panagiotou, Samotij used [an alternating cumulant series](#), whose terms after simplifications are responding to partitions in cluster expansion.

## Joint cumulant (1929)

For any multiset of random variables  $\{X_i\}_{i \in S}$ ,

$$\kappa(S) = \sum_{\pi \in \Pi(S)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{P \in \pi} \mu(P).$$

Also called *semi-invariant* (1889) in probability, *Ursell function* (1927) in physics.  
 For  $G$ -dependent variables and  $S \subseteq V(G)$ ,  
 if  $S$  [does not induce a connected subgraph](#) in  $G$ , then  $\kappa(S) = 0$ .

- Stark and Wormald (2018) also have results for  $\mathcal{G}_{n,m}$ :  
 if  $d = m \binom{n}{2}^{-1} = o(n^{-7/11})$ , then

$$\mathbb{P}(X = 0) = \exp\left(-\frac{1}{6}n^3 d^3 - \frac{1}{8}n^4 d^6 - \frac{1}{2}n^2 d^3 + o(1)\right).$$

## Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex.

## Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$\mathbb{P}(\text{a random hypergraph is linear})$

$= \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).$

## Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$\mathbb{P}(\text{a random hypergraph is linear})$

$= \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).$

McKay and Tian, 2020

If  $p = o(n^{-3/2})$ , then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 p^2 + \frac{2}{3}n^5 p^3 + o(1)\right).$$

## Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$\mathbb{P}(\text{a random hypergraph is linear})$

$= \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).$

McKay and Tian, 2020

If  $p = o(n^{-3/2})$ , then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 p^2 + \frac{2}{3}n^5 p^3 + o(1)\right).$$

Z. 2022; Wormald and Z., 2022+

If  $p = o(n^{-7/5})$ , then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 p^2 + \frac{2}{3}n^5 p^3 - \frac{55}{24}n^6 p^4 + \frac{3}{2}n^3 p^2 + o(1)\right).$$

## Asymptotic linearity of hypergraphs

A hypergraph is linear if each pair of hyperedges intersects in at most one vertex. So

$\mathbb{P}(\text{a random hypergraph is linear})$

$= \mathbb{P}(\text{no hyperedge-pairs intersect in more than one vertex}).$

### McKay and Tian, 2020

If  $p = o(n^{-3/2})$ , then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 p^2 + \frac{2}{3}n^5 p^3 + o(1)\right).$$

### Z. 2022; Wormald and Z., 2022+

If  $p = o(n^{-7/5})$ , then

$$\mathbb{P}(H_3(n, p) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 p^2 + \frac{2}{3}n^5 p^3 - \frac{55}{24}n^6 p^4 + \frac{3}{2}n^3 p^2 + o(1)\right).$$

- In Z. 2022, the truncation error is by Mousset, Noever, Panagiotou, Samotij's results.
- Coefficients match McKay's conjecture based upon numerical simulation.



A recent finding:

### Corollary

$$\begin{aligned}\kappa(S) &= \sum_{\{C_1, \dots, C_n\} \in \Pi(S)} (-1)^{n-1} |\mathcal{T}| \prod_{i \in [n]} \mu(C_i) 1_{\{C_i \text{ is connected}\}} \\ &\leq \sum_{\{C_1, \dots, C_n\} \in \Pi(S)} |\mathcal{T}| \prod_{i \in [n]} \mu(C_i) 1_{\{C_i \text{ is connected}\}}\end{aligned}$$

where  $\mathcal{T} = \mathcal{T}(C_1, \dots, C_n)$  denotes a special set of spanning trees by Penrose (1967), who used it to obtain the first convergence criterion of the cluster expansion.

A recent finding:

### Corollary

$$\begin{aligned}\kappa(S) &= \sum_{\{C_1, \dots, C_n\} \in \Pi(S)} (-1)^{n-1} |\mathcal{T}| \prod_{i \in [n]} \mu(C_i) 1_{\{C_i \text{ is connected}\}} \\ &\leq \sum_{\{C_1, \dots, C_n\} \in \Pi(S)} |\mathcal{T}| \prod_{i \in [n]} \mu(C_i) 1_{\{C_i \text{ is connected}\}}\end{aligned}$$

where  $\mathcal{T} = \mathcal{T}(C_1, \dots, C_n)$  denotes a special set of spanning trees by Penrose (1967), who used it to obtain the first convergence criterion of the cluster expansion.

- This implies Janson's bound (1988) on cumulants, thus gives asymptotic normality.

A recent finding:

### Corollary

$$\begin{aligned}\kappa(S) &= \sum_{\{C_1, \dots, C_n\} \in \Pi(S)} (-1)^{n-1} |\mathcal{T}| \prod_{i \in [n]} \mu(C_i) 1_{\{C_i \text{ is connected}\}} \\ &\leq \sum_{\{C_1, \dots, C_n\} \in \Pi(S)} |\mathcal{T}| \prod_{i \in [n]} \mu(C_i) 1_{\{C_i \text{ is connected}\}}\end{aligned}$$

where  $\mathcal{T} = \mathcal{T}(C_1, \dots, C_n)$  denotes a special set of spanning trees by Penrose (1967), who used it to obtain the first convergence criterion of the cluster expansion.

- This implies Janson's bound (1988) on cumulants, thus gives asymptotic normality.
- This relates to 'weighted dependency graph' by Feray (2018), who also bounds cumulants via spanning trees and obtains asymptotic normality under weak dependence (e.g., small subgraphs in  $\mathcal{G}(n, m)$ ).

# Asymptotic linearity of hypergraphs

## McKay and Tian, 2020

If  $d = \binom{n}{3}^{-1} m = o\left(n^{-3/2}\right)$ , then via switching method,

$$\mathbb{P}(H_3(n, m) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 d^2 - \frac{1}{12}n^5 d^3 + o(1)\right).$$

## Wormald and Z. 2022+

If  $d = \binom{n}{3}^{-1} m = o\left(n^{-7/5}\right)$ , then via perturbation method,

$$\mathbb{P}(H_3(n, m) \text{ is linear}) = \exp\left(-\frac{1}{4}n^4 d^2 - \frac{1}{12}n^5 d^3 - \frac{1}{24}n^6 d^4 + \frac{3}{2}n^3 d^2 + o(1)\right).$$

## Dependency digraphs

### Dependency digraphs

A digraph  $D = ([d], E)$  is a dependency digraph for the events  $\{A_i\}_{i \in [d]}$  if  $A_i$  is mutually independent of all the non-adjacent  $\{A_j : (i, j) \notin E\}$  for every  $i \in [d]$ .


# Dependency digraphs

## Dependency digraphs

A digraph  $D = ([d], E)$  is a dependency digraph for the events  $\{A_i\}_{i \in [d]}$  if  $A_i$  is mutually independent of all the non-adjacent  $\{A_j : (i, j) \notin E\}$  for every  $i \in [d]$ .

## Example

Let  $\{A_i\}_{i \in [5]}$  be events with the following dependency digraph.




$A_1$  is independent of  $A_4$ .

$A_2$  is independent of  $\{A_1, A_3, A_5\}$ .

A natural refinement is

- adjacent vertices  $\rightarrow$  the pairs of 'strongly dependent' events,  
 $D_i \subseteq [d]$  denotes the 'strongly dependent neighbours' of event  $A_i$ .
- non-adjacent vertices  $\rightarrow$  the pairs of 'weakly dependent' events.




$$\mathbb{P}(A_4 | A_1) \approx \mathbb{P}(A_4)$$

$$\mathbb{P}(A_1 \cup A_3 \cup A_5 | A_2) \approx \mathbb{P}(A_1 \cup A_3 \cup A_5)$$

A natural refinement is

- adjacent vertices  $\rightarrow$  the pairs of 'strongly dependent' events,  
 $D_i \subseteq [d]$  denotes the 'strongly dependent neighbours' of event  $A_i$ .
- non-adjacent vertices  $\rightarrow$  the pairs of 'weakly dependent' events.



$$\mathbb{P}(A_4 | A_1) \approx \mathbb{P}(A_4)$$

$$\mathbb{P}(A_1 \cup A_3 \cup A_5 | A_2) \approx \mathbb{P}(A_1 \cup A_3 \cup A_5)$$

If we change ' $\approx$ ' to ' $\geq$ ', then it is the 'negative dependency graph' by Erdős and Spencer (1991) to obtain lopsided Lovász local lemma.



## New dependency digraphs

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  on  $[d]$ :

- $\varphi$ -mixing coefficient is used to measure 'weak dependencies':

$$\varphi = \max_{i \in [d]} \left| \mathbb{P} \left( \bigcup_{j \in [i-1] \setminus D_i} A_j \mid A_i \right) - \mathbb{P} \left( \bigcup_{j \in [i-1] \setminus D_i} A_j \right) \right|.$$

- The influence of 'strongly dependent' events is

$$\Delta_1 = \sum_{i \in [d]} \mathbb{P} \left( A_i \cap \bigcup_{j \in [i-1] \cap D_i} A_j \right) \prod_{k \in [d] \setminus [i]} \mathbb{P}(\overline{A_k})$$
$$\Delta_2 = \sum_{i \in [d]} \mathbb{P}(A_i) \mathbb{P} \left( \bigcup_{j \in [i-1] \cap D_i} A_j \right) \prod_{k \in [d] \setminus [i]} \mathbb{P}(\overline{A_k}).$$

## New dependency digraphs

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  on  $[d]$ :

- $\varphi$ -mixing coefficient is used to measure 'weak dependencies'

$$\varphi = \max_{i \in [d]} \left| \mathbb{P} \left( \bigcup_{j \in [i-1] \setminus D_i} A_j \mid A_i \right) - \mathbb{P} \left( \bigcup_{j \in [i-1] \setminus D_i} A_j \right) \right|.$$

- The influence of 'strongly dependent' events is

$$\Delta_1 \leq \Delta'_1 = \sum_{i \in [d]} \sum_{j \in [i-1] \cap D_i} \mathbb{P}(A_i \cap A_j),$$
$$\Delta_2 \leq \Delta'_2 = \sum_{i \in [d]} \sum_{j \in [i-1] \cap D_i} \mathbb{P}(A_i) \mathbb{P}(A_j).$$

- Chen–Stein Poisson approximation involves similar terms as  $\varphi$ ,  $\Delta'_1$ , and  $\Delta'_2$ .  
Chen (1975) also considered the  $\varphi$ -mixing condition.
- Suen's inequality (1990) also involves  $\Delta'_1$  and  $\Delta'_2$ .

### Theorem (Isaev, Rodionov, Z., Zhukovskii, 2020, 2022+)

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  with vertex set  $[d]$ , we have

$$\left| \mathbb{P} \left( \bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left( 1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.

### Theorem (Isaev, Rodionov, Z., Zhukovskii, 2020, 2022+)

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  with vertex set  $[d]$ , we have

$$\left| \mathbb{P} \left( \bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left( 1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If  $\varphi = o(1)$ ,  $\Delta_1 = o(1)$ , and  $\Delta_2 = o(1)$ , then we have the asymptotic independence.

### Theorem (Isaev, Rodionov, Z., Zhukovskii, 2020, 2022+)

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  with vertex set  $[d]$ , we have

$$\left| \mathbb{P} \left( \bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left( 1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If  $\varphi = o(1)$ ,  $\Delta_1 = o(1)$ , and  $\Delta_2 = o(1)$ , then we have the asymptotic independence.
  - ▶ 'Two Moments Suffice for Poisson Approximations: The Chen–Stein Method' by Arratia, Goldstein and Gordon (1989); also Chen (1975).

### Theorem (Isaev, Rodionov, Z., Zhukovskii, 2020, 2022+)

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  with vertex set  $[d]$ , we have

$$\left| \mathbb{P} \left( \bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left( 1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If  $\varphi = o(1)$ ,  $\Delta_1 = o(1)$ , and  $\Delta_2 = o(1)$ , then we have the asymptotic independence.
  - ▶ 'Two Moments Suffice for Poisson Approximations: The Chen–Stein Method' by Arratia, Goldstein and Gordon (1989); also Chen (1975).
- Max codegrees, max clique-extension counts of random (hyper)graphs, etc. are asymptotically Gumbel.

### Theorem (Isaev, Rodionov, Z., Zhukovskii, 2020, 2022+)

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  with vertex set  $[d]$ , we have

$$\left| \mathbb{P} \left( \bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left( 1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$

- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If  $\varphi = o(1)$ ,  $\Delta_1 = o(1)$ , and  $\Delta_2 = o(1)$ , then we have the asymptotic independence.
  - ▶ '*Two Moments Suffice for Poisson Approximations: The Chen–Stein Method*' by Arratia, Goldstein and Gordon (1989); also Chen (1975).
- Max codegrees, max clique-extension counts of random (hyper)graphs, etc. are asymptotically Gumbel.
- Potential application: asymptotic enumeration of rainbow matchings (Latin transversals).

### Theorem (Isaev, Rodionov, Z., Zhukovskii, 2020, 2022+)

For any events  $\{A_i\}_{i \in [d]}$  and digraph  $D$  with vertex set  $[d]$ , we have

$$\left| \mathbb{P} \left( \bigcap_{i \in [d]} \overline{A_i} \right) - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right| \leq \left( 1 - \prod_{i \in [d]} \mathbb{P}(\overline{A_i}) \right) \varphi + \max(\Delta_1, \Delta_2).$$



- The lower bound generalises Dubickas' bound (2008), and the upper bound is via a similar inductive proof.
- If  $\varphi = o(1)$ ,  $\Delta_1 = o(1)$ , and  $\Delta_2 = o(1)$ , then we have the asymptotic independence.
  - ▶ *'Two Moments Suffice for Poisson Approximations: The Chen–Stein Method'* by Arratia, Goldstein and Gordon (1989); also Chen (1975).
- Max codegrees, max clique-extension counts of random (hyper)graphs, etc. are asymptotically Gumbel.
- Potential application: asymptotic enumeration of rainbow matchings (Latin transversals).
- Lu and Székely (2009+?) also introduced a notion of 'positive' dependency graph to obtain upper bounds for Poisson approximation for random matchings.



## Regular tournaments

A tournament is a digraph obtained by assigning a direction for each edge in an undirected complete graph.



### Example



## Regular tournaments

A tournament is a digraph obtained by assigning a direction for each edge in an undirected complete graph.

### Example




A tournament is regular if the in-degree is equal to the out-degree for each vertex.

## Regular tournaments

A tournament is a digraph obtained by assigning a direction for each edge in an undirected complete graph.

### Example



A tournament is regular if the in-degree is equal to the out-degree for each vertex. Let  $RT(n)$  denote the number of regular tournament on  $n$  vertices.

### Theorem (McKay, 1990)

For odd  $n \rightarrow \infty$  and  $\varepsilon > 0$ ,

$$RT(n) = \left(1 + O\left(n^{-1/2+\varepsilon}\right)\right) \left(\frac{n}{e}\right)^{1/2} \left(\frac{2^{n+1}}{\pi n}\right)^{(n-1)/2}.$$

# Asymptotic enumeration of regular tournaments

The method was developed by McKay and Wormald (1990)  
based on generating function + Cauchy's integral theorem + the saddle point method.

$$\begin{aligned} \text{RT}(n) &= \frac{1}{(2\pi i)^n} \oint \cdots \oint \frac{\prod_{1 \leq j < k \leq n} (x_j/x_k + x_k/x_j)}{x_1 \cdots x_n} dx_1 \cdots dx_n, \\ &= \frac{2^{n(n-1)/2}}{(2\pi)^n} \int_{-\pi}^{\pi} \cdots \int_{-\pi}^{\pi} \prod_{1 \leq j < k \leq n} \cos(\theta_j - \theta_k) d\theta. \end{aligned}$$

The remaining is to estimate the  $n$ -dimensional integral.

### Theorem (McKay, 1990)

For odd  $n \rightarrow \infty$  and  $\varepsilon > 0$ ,

$$\text{RT}(n) = \left(1 + O\left(n^{-1/2+\varepsilon}\right)\right) \left(\frac{n}{e}\right)^{1/2} \left(\frac{2^{n+1}}{\pi n}\right)^{(n-1)/2}.$$

Via the complex martingale method by Isaev and McKay (2018), and an estimate of the integral via cumulants of multivariate Gaussians, we have:

### Theorem (Isaev, McKay, Z., 2022+)



For odd  $n \rightarrow \infty$ ,

$$\begin{aligned} \text{RT}(n) = n^{1/2} \left(\frac{2^{n+1}}{\pi n}\right)^{(n-1)/2} \exp \left( -\frac{1}{2} + \frac{1}{4n} + \frac{1}{4n^2} + \frac{7}{24n^3} + \frac{37}{120n^4} + \frac{31}{60n^5} + \frac{81}{28n^6} \right. \\ \left. + \frac{5981}{336n^7} + \frac{22937}{240n^8} + \frac{90031}{180n^9} + \frac{1825009}{660n^{10}} + \frac{4344847}{264n^{11}} + O\left(n^{-12}\right) \right). \end{aligned}$$

## Eulerian digraphs

A simple directed graph is a directed graph having no multiple edges or loops.


### Example



## Eulerian digraphs

A simple directed graph is a directed graph having no multiple edges or loops.

### Example



An Eulerian digraph is a digraph s.t. the in-degree equals the out-degree for each vertex.

### Theorem (McKay, 1990)

For  $n \rightarrow \infty$  and  $\varepsilon > 0$ ,

$$\text{ED}(n) = \left(1 + O\left(n^{-1/2+\varepsilon}\right)\right) \frac{n^{1/2}}{e^{1/4}} \left(\frac{4^n}{\pi n}\right)^{(n-1)/2}.$$

### Theorem (McKay, 1990)

For  $n \rightarrow \infty$  and  $\varepsilon > 0$ ,

$$\text{ED}(n) = \left(1 + O\left(n^{-1/2+\varepsilon}\right)\right) \frac{n^{1/2}}{e^{1/4}} \left(\frac{4^n}{\pi n}\right)^{(n-1)/2}.$$

### Theorem (Isaev, McKay, Z., 2022+)

For  $n \rightarrow \infty$ ,


$$\begin{aligned} \text{ED}(n) = n^{1/2} \left(\frac{4^n}{\pi n}\right)^{(n-1)/2} \exp \left( -\frac{1}{4} + \frac{3}{16n} + \frac{1}{8n^2} + \frac{47}{384n^3} + \frac{371}{1920n^4} + \frac{1807}{3840n^5} + \frac{655}{448n^6} \right. \\ \left. + \frac{435581}{86016n^7} + \frac{1145941}{61440n^8} + \frac{13318871}{184320n^9} + \frac{99074137}{337920n^{10}} + \frac{1339710847}{1081344n^{11}} + O\left(n^{-12}\right) \right). \end{aligned}$$



## Eulerian oriented graphs

A directed graph having no symmetric pair of directed edges is an oriented graph (a complete oriented graph is a tournament).

### Example



Eulerian oriented graphs are Eulerian digraphs with no symmetric pair of directed edges.

### Theorem (McKay, 1990)

For  $n \rightarrow \infty$  and  $\varepsilon > 0$ ,

$$\text{EOG}(n) = \left(1 + O\left(n^{-1/2+\varepsilon}\right)\right) \frac{n^{1/2}}{e^{3/8}} \left(\frac{3^{n+1}}{4\pi n}\right)^{(n-1)/2}.$$

### Theorem (McKay, 1990)

For  $n \rightarrow \infty$  and  $\varepsilon > 0$ ,

$$\text{EOG}(n) = \left(1 + O\left(n^{-1/2+\varepsilon}\right)\right) \frac{n^{1/2}}{e^{3/8}} \left(\frac{3^{n+1}}{4\pi n}\right)^{(n-1)/2}.$$

### Theorem (Isaev, McKay, Z., 2022+)

For  $n \rightarrow \infty$ ,

$$\begin{aligned} \text{EOG}(n) = & n^{1/2} \left(\frac{3^{n+1}}{4\pi n}\right)^{(n-1)/2} \exp\left(-\frac{3}{8} + \frac{11}{64n} + \frac{7}{64n^2} + \frac{233}{2048n^3} + \frac{497}{2560n^4} + \frac{27583}{61440n^5} \right. \\ & + \frac{55463}{43008n^6} + \frac{33678923}{7340032n^7} + \frac{101414573}{5242880n^8} + \frac{1882520759}{20971520n^9} + \frac{101145677531}{230686720n^{10}} \\ & \left. + \frac{2469157786549}{1107296256n^{11}} + O\left(n^{-12}\right)\right). \end{aligned}$$

Results that are not covered but will be in thesis include

- asymptotic extremes for  $m$ -dependent random variables via cluster expansion, (a not-so-trivial convergent application).

Results that are not covered but will be in thesis include

- asymptotic extremes for  $m$ -dependent random variables via cluster expansion, (a not-so-trivial convergent application).
- asymptotic probability of non-existence of small subgraphs in random graphs with high accuracy (with Wormald).

Results that are not covered but will be in thesis include

- asymptotic extremes for  $m$ -dependent random variables via cluster expansion, (a not-so-trivial convergent application).
- asymptotic probability of non-existence of small subgraphs in random graphs with high accuracy (with Wormald).
- asymptotic enumeration of Eulerian orientations via cumulants (with Isaev, McKay).

Results that are not covered but will be in thesis include

- asymptotic extremes for  $m$ -dependent random variables via cluster expansion, (a not-so-trivial convergent application).
- asymptotic probability of non-existence of small subgraphs in random graphs with high accuracy (with Wormald).
- asymptotic enumeration of Eulerian orientations via cumulants (with Isaev, McKay).

Results that are not covered but will be in thesis include

- asymptotic extremes for  $m$ -dependent random variables via cluster expansion, (a not-so-trivial convergent application).
- asymptotic probability of non-existence of small subgraphs in random graphs with high accuracy (with Wormald).
- asymptotic enumeration of Eulerian orientations via cumulants (with Isaev, McKay).

Some other results include

- concentration under graph-dependence (2022),
- learning under graph-dependence (with Amini),
- ....

## A plan for the structure of thesis

### 1 Cumulants and cluster expansion.

- ▶ Asymptotic probability of non-existence and linearity of binomial random hypergraphs.
- ▶ Asymptotic normality.
- ▶ Asymptotic extremes for  $m$ -dependent random variables.



## A plan for the structure of thesis

### 1 Cumulants and cluster expansion.

- ▶ Asymptotic probability of non-existence and linearity of binomial random hypergraphs.
- ▶ Asymptotic normality.
- ▶ Asymptotic extremes for  $m$ -dependent random variables.

### 2 Asymptotic probability of non-existence via perturbation method (with Wormald).

- ▶ Asymptotic linearity of random hypergraphs.
- ▶ Non-existence of small subgraphs in random graphs with high accuracy.

## A plan for the structure of thesis

- 1 Cumulants and cluster expansion.
  - ▶ Asymptotic probability of non-existence and linearity of binomial random hypergraphs.
  - ▶ Asymptotic normality.
  - ▶ Asymptotic extremes for  $m$ -dependent random variables.
- 2 Asymptotic probability of non-existence via perturbation method (with Wormald).
  - ▶ Asymptotic linearity of random hypergraphs.
  - ▶ Non-existence of small subgraphs in random graphs with high accuracy.
- 3 Asymptotic independence under mixing via dependency digraphs (with Isaev, Rodionov, Zhukovskii).

## A plan for the structure of thesis

- 1 Cumulants and cluster expansion.
  - ▶ Asymptotic probability of non-existence and linearity of binomial random hypergraphs.
  - ▶ Asymptotic normality.
  - ▶ Asymptotic extremes for  $m$ -dependent random variables.
- 2 Asymptotic probability of non-existence via perturbation method (with Wormald).
  - ▶ Asymptotic linearity of random hypergraphs.
  - ▶ Non-existence of small subgraphs in random graphs with high accuracy.
- 3 Asymptotic independence under mixing via dependency digraphs (with Isaev, Rodionov, Zhukovskii).
- 4 Cumulants and asymptotic enumeration of RT, ED, and EOG (with Isaev, McKay).

## A plan for the structure of thesis

- 1 Cumulants and cluster expansion.
  - ▶ Asymptotic probability of non-existence and linearity of binomial random hypergraphs.
  - ▶ Asymptotic normality.
  - ▶ Asymptotic extremes for  $m$ -dependent random variables.
- 2 Asymptotic probability of non-existence via perturbation method (with Wormald).
  - ▶ Asymptotic linearity of random hypergraphs.
  - ▶ Non-existence of small subgraphs in random graphs with high accuracy.
- 3 Asymptotic independence under mixing via dependency digraphs (with Isaev, Rodionov, Zhukovskii).
- 4 Cumulants and asymptotic enumeration of RT, ED, and EOG (with Isaev, McKay).

Thanks!