



Living under surveillance: Gender, psychological distress, and stop-question-and-frisk policing in New York City



Abigail A. Sewell ^{a,*}, Kevin A. Jefferson ^b, Hedwig Lee ^c

^a Emory University, Department of Sociology, USA

^b Emory University, Rollins School of Public Health, USA

^c University of Washington, Department of Sociology, USA

ARTICLE INFO

Article history:

Received 26 October 2015

Received in revised form

18 April 2016

Accepted 20 April 2016

Available online 22 April 2016

Keywords:

Police

Neighborhoods

New York City

Gender

Psychological distress

Mental health

Mass incarceration

Men's health

ABSTRACT

A growing body of research highlights the collateral consequences of mass incarceration, including stop-and-frisk policing tactics. Living in a neighborhood with aggressive policing may affect one's mental health, especially for men who are the primary targets of police stops. We examine whether there is an association between psychological distress and neighborhood-level aggressive policing (i.e., frisking and use of force by police) and whether that association varies by gender. The 2009–2011 New York City (NYC) Stop, Question, and Frisk Database is aggregated to the neighborhood-level ($N = 34$) and merged with individual data from the 2012 NYC Community Health Survey ($N = 8066$) via the United Hospital Fund neighborhood of respondents' residence. Weighted multilevel generalized linear models are used to assess main and gendered associations of neighborhood exposures to aggressive police stops on psychological distress (Kessler-6 items). While the neighborhood stop rate exhibits inconsistent associations with psychological distress, neighborhood-level frisk and use of force proportions are linked to higher levels of non-specific psychological distress among men, but not women. Specifically, men exhibit more non-specific psychological distress and more severe feelings of nervousness, effort, and worthlessness in aggressively surveilled neighborhoods than do women. Male residents are affected by the escalation of stop-and-frisk policing in a neighborhood. Living in a context of aggressive policing is an important risk factor for men's mental health.

© 2016 Elsevier Ltd. All rights reserved.

Media (Balko, 2015; Goodman, 2015; Leovy, 2015; Lewis, 2015; The Associated Press, 2015) and scholarly (Cooper et al., 2004; Lee et al., 2014; Massoglia, 2008; Massoglia and Pridemore, 2015; Sewell and Jefferson, 2016) attention to the over-policing of neighborhoods has grown. For residents of such neighborhoods, police stops may be a chronic stressor occurring repeatedly over days, weeks, months, and years (Brunson and Miller, 2006; Brunson and Weitzer, 2008; Cooper et al., 2004; Engel and Calnon, 2004; Lerman and Weaver, 2014). The health associations of such a stressor are only beginning to be examined. This study uses a multilevel research design to examine the mental health of individuals living in highly surveilled neighborhoods, particularly neighborhoods that are surveilled with aggressive policing tactics of frisking and use of force. Special

attention is paid to gender differences in the strength of the association of living in neighborhoods where pedestrian stops are more likely to incur frisking and use of force because men are the primary targets of stop-and-frisk practices. Specifically, individual-level health data from the New York City Community Health Survey (NYC-CHS) is matched to geocoded administrative data from the New York City Stop, Question, and Frisk Database (NYC-SQF), and a cross-level interaction between male gender and living in an (aggressively) surveilled neighborhood is evaluated.

The policing of pedestrians is gendered. For instance, an analysis of administrative data from the NYC-SQF indicates that men comprise about 88 percent of pedestrian stop suspects in NYC (Ridgeway, 2007). As such, men's mental health may be especially affected by direct contact with the police. Indeed, a recent study of young men in NYC found higher levels of trauma and anxiety symptoms among those men who reported frequent contact with the police, especially if such contact was conceived as "intrusive" or

* Corresponding author. Department of Sociology, Emory University, 1555 Dickey Drive, 204 Tarbutton Hall, Atlanta, GA 30322, USA.

E-mail address: abigail.a.sewell@emory.edu (A.A. Sewell).

“unfair” (Geller et al., 2014). Men living in highly policed neighborhoods, especially men of color, indicate high levels of worry and anticipation caused by the possibility of being stopped by the police at any time, as well as anger, frustration, and resentment caused by the perception that police unfairly target them (Anderson, 1990, 1998, 2003; Dottolo and Stewart, 2008; Goffman, 2009; Jones, 2014; Young, 2006).

Yet, a burgeoning body of research suggests that *indirect* contact with the criminal justice system is also associated with negative health consequences (Hatzenbuehler et al., 2014; Lee et al., 2014; Sewell and Jefferson, 2016; Wildeman et al., 2012). For instance, living in a neighborhood with high incarceration rates is linked to higher risks of depression and anxiety (Hatzenbuehler et al., 2014) and asthma (Frank et al., 2013) for men and women. On one hand, living in neighborhoods where pedestrian stops are more likely to incur frisking and use of force may represent a unique stressor for men, who are more likely to be stopped and, therefore, anticipate being stopped. On the other hand, both men and women living in such neighborhoods may experience policing similarly, suggesting that neighborhood policing invokes the same types of stress and coping processes across the gender continuum.

Research suggests a palpable relationship between aggressive policing and the risks of psychiatric illness for men (Geller et al., 2014). Living in highly policed areas may be harmful to one's mental health, via the negative effects of hypervigilance and perceived unfairness (Brunson and Miller, 2006; Brunson and Weitzer, 2008; Cooper et al., 2004; Geller et al., 2014; Kessler et al., 1999; McEwen, 2004; Shedd, 2012; Unnever and Gabbidon, 2011; Williams and Mohammed, 2009; Williams et al., 2003; Williams et al., 1997). When police stops within a neighborhood frequently escalate to frisking and use of force, residents are more likely to perceive stops as discriminatory or unfair (Brunson and Miller, 2006; Brunson and Weitzer, 2008; Cooper et al., 2004; Unnever and Gabbidon, 2011). Perceived unfairness not only is linked to poor mental health (Kessler et al., 1999; McEwen, 2004; Williams et al., 1997, 2003) but also creates a “climate of fear” in which residents live with knowledge that they could be criminalized at any moment and in turn feel more vigilant (Shedd, 2012). Moreover, hypervigilance, through the psychological expenses of chronically activating coping mechanisms (Cohen et al., 1986; Meyer, 1995), can produce changes in the hippocampus, prefrontal cortex, and amygdala that precipitate depression and/or anxiety (Kessler et al., 1999; Lerman and Weaver, 2014; McEwen, 2004; Williams et al., 1997, 2003). The climate of fear produced by aggressive policing practices may be especially impactful for male residents who are more likely to feel it is unfair that the police target them on a day-to-day basis. Perceptions of injustice in policing, in turn, may be directly linked to pedestrian stops turning aggressive, such that ill effects are associated with frisking and use of force, but not the rate of pedestrian stops in the neighborhood.

To our knowledge, this is the first study to examine the mental health consequences of stop-and-frisk policing at the community-level. We examine (1) the association between psychological distress and the escalation of neighborhood police stops, holding constant key individual- and neighborhood-level correlates of health and (2) gender variation in the association between mental health and community-level escalated police encounters. We propose two hypotheses that are examined using data on pedestrian stops in NYC:

1. Living in (aggressively) surveilled neighborhoods is associated with a greater risk of reporting psychological distress among neighborhood residents.

2. The association between living in such (aggressively) surveilled neighborhoods and psychological distress is stronger for men than for women.

1. Methods

1.1. Data

This multilevel study merges individual-level data from one data source with neighborhood-level data from multiple data sources.

1.1.1. Individual level

The analysis is based on a sample of adults ($N = 8797$) participating in the 2012 NYC-CHS collected by the NYC Department of Health and Mental Hygiene (New York City Department of Health and Mental Hygiene, 2012). NYC-CHS is an annual random-digital health survey of non-institutionalized adult (18+) New Yorkers. This survey evaluates the health of New York residents city-wide, by neighborhood, and across demographic subpopulations. The survey gathers a broad range of health measures and is based on the national Behavioral Risk Factor Surveillance System coordinated by the U.S. Centers for Disease Control and Prevention. Each year, the NYC-CHS interviews approximately 10,000 people. All data, collected by telephone or cellphone, are self-reported, publicly-available, and include community identifiers for individuals surveyed. Respondents live in 1 of 34 United Hospital Fund (UHF) neighborhoods (NYC Department of Health and Mental Hygiene, 2006); only 303 respondents (0.3%) were missing neighborhood identifiers.

1.1.2. Neighborhood level

Administrative data from the 2009–2011 NYC-SQF provide pedestrian stop-level data (New York City Police Department, 2009–2011), which are aggregated to UHF neighborhoods by geocoding stop locations for 1,816,871 of the 1,818,465 pedestrian stops (99.9%) occurring during the study period. These data are used to create neighborhood-level densities of post-stop outcomes by counting the number of stops occurring in each UHF neighborhood and the outcome of the stop. Data from the 2010 decennial census and the New York City Police Department (NYPD) provide additional neighborhood-level variables. The publicly available data are exempt from IRB review by the authors' institutions.

1.2. Measures

1.2.1. Outcome measures

Our outcomes of interest are the six items comprising the Kessler-6 Psychological Distress Scale (K6) (Kessler et al., 2003), as well as a dichotomous measure of psychological distress based on this scale (Table 1). All NYC-CHS respondents are asked the following questions: “During the past 30 days, how often did you feel: (1) So sad that nothing could cheer you up?; (2) Nervous?; (3) Restless or fidgety?; (4) Hopeless?; (5) That everything was an effort?; and (6) Worthless? The response categories for the items are: (1) All of the time; (2) Most of the time; (3) Some of the time; (4) A little of the time; (5) None of the time. Responses are reverse coded, such that higher values indicate more frequent and severe psychological distress. Individuals who responded “don't know”, “not sure”, or refused to provide a response are classified as missing data. As is convention (Kessler et al., 2003), summary scale scores (Range: 6–30) were converted to a dichotomous measure of non-specific psychological distress (NSPD) (1: $K6 = 12+$; 0: $K6 \leq 12$). We treat NSPD as a binary response outcome, while we

Table 1Descriptive statistics for individual-level covariates: 2012 NYC community health survey ($N = 8066$) and 2009–2011 NYC stop, question, and frisk ($N = 34$).

	Mean	Median	SD	Min	Max
Psychological Distress ($N = 8066$)					
Non-Specific Psychological Distress (NSPD) ^a	0.23			0	1
Sadness	1.56	1	0.94	1	5
Nervousness	1.78	1	0.96	1	5
Restlessness	1.73	1	1.01	1	5
Hopelessness	1.36	1	0.79	1	5
Effort	1.89	1	1.19	1	5
Worthlessness	1.30	1	0.75	1	5
Individual-Level Covariates ($N = 8066$)					
Black Non-Latino	0.23			0	1
Latino	0.26			0	1
Asian/Pacific Islander	0.09			0	1
White (Reference Category)	0.42			0	1
Male (0 = Female)	0.41			0	1
Age (in Years)	52.57	52	17.44	18	98
Less than High School (Reference Category)	0.15			0	1
High School	0.21			0	1
Some College	0.20			0	1
College Degree	0.43			0	1
Employed (0 = Unemployed)	0.55			0	1
US Born (0 = Not US Born)	0.60			0	1
Insured (0 = Not Insured)	0.88			0	1
Unmet Medical Care Last Year (0 = Met)	0.10			0	1
Household Size	2.50	2	1.49	1	6
Currently Married/Cohabiting (Reference Category)	0.44			0	1
Formerly Married	0.29			0	1
Never Married	0.27			0	1
Primarily English at Home (Reference Category)	0.73			0	1
Primarily Spanish at Home	0.16			0	1
Primarily Other Language at Home	0.11			0	1

Note: SD = Standard Deviation.

^a NSPD is indicated by a K6 Score of 12 or higher (Min: 6; Max: 30), where the K6 Score is a summation of K6 items (sadness, nervousness, restlessness, hopelessness, required effort, worthlessness).

treat the six constituent items of the K6 as five-level ordinal outcomes.

Prior research using the 2002–2003 NYC-CHS found that K6 items are distributed differentially across the urban environment, with feelings of effort being most prevalent among New Yorkers and feeling of hopelessness being least prevalent (McVeigh et al., 2006). As such, this study examines the dichotomous indicator of NSPD separately from the ordinal measures of its constituent items. Fig. 1 provides a visualization of the spread of the dichotomous NSPD indicator in NYC.

1.2.2. Neighborhood policing measures

We examine two neighborhood-level post-stop policing characteristics using NYC-SQF data: frisking and use of force. All neighborhood-level variables are centered at their means and standardized (Table 2). Fig. 2 provides a map of the distribution of the neighborhood post-stop policing measures.

The post-stop policing characteristics portray the culture of aggressive policing tactics in a neighborhood. Frisking is reported per stop by officers and includes quickly passing the hands over a pedestrian's clothes or through a pedestrian's pockets. Nine types of physical force by the officer are also reported per stop by officers and includes the use of the officer's hands, placing a suspect on the ground, placing a suspect against the wall, drawing the officer's weapon, pointing the weapon at the suspect, and use of baton, handcuffs, pepper spray, or other physical object. If a stop involves any one of the nine types of physical force by the officer, then that stop is considered a stop involving use of force. Two measures are created based off of all stops in a neighborhood: (1) the proportion of stops that involve frisking

and (2) the proportion of stops that involve any use of physical force. As there is a strong relationship between frisking and use of force at the neighborhood-level ($r = 0.64$; $p < 0.001$), the health associations of these two post-stop outcomes are evaluated independently.

1.2.3. Neighborhood-level covariates

A measure of stop productivity (the proportion of stops that result in an arrest being made) per UHF from NYC-SQF data is included, as is the NYPD reported average count of robbery complaints between 2010 and 2012 in a neighborhood and the 2010 decennial Census reported total number of persons living in a neighborhood that are currently incarcerated (i.e., living in correctional facilities). Several measures of ethnoracial and economic segregation from the 2010 decennial Census are included as well: (1) the proportion of the population that identifies as Black or Latino; (2) the proportion of households with incomes below the federal poverty line; and (3) a measure of affluence concentration. Affluence concentration is measured using a z-score that captures the relatedness of the proportion of owner-occupied homes in a neighborhood and the proportion of households with incomes above \$50,000 a year in a neighborhood (Browning and Cagney, 2003). These two indicators of neighborhood affluence are highly correlated ($r = 0.73$). Table 2 provides descriptive statistics for neighborhood-level covariates in their native form.

1.2.4. Individual-level covariates

At the individual level, we consider key sociodemographic (gender, age, race/ethnicity, nativity, marital status, household size,




Fig. 1. Distribution of non-specific psychological distress by United Hospital Fund neighborhoods.

Source: 2012 New York City Community Health Survey; Weighted distributions.

Table 2

Descriptive statistics for neighborhood-level covariates: 2009–2011 NYC stop, question, and frisk ($N = 34$).

	Mean	Median	SD	Min	Max
Neighborhood-Level Covariates ($N = 34$)					
Stop Rate Per 100 Non-Institutionalized Persons, 2009–2011	22.79	17.76	14.95	6.32	65.11
Proportion of Stops Resulting in Frisking, 2009–2011	0.54	0.54	0.09	0.35	0.73
Proportion of Stops Resulting in Use of Force, 2009–2011	0.22	0.20	0.08	0.10	0.36
Proportion of Stops Resulting in Arrest, 2009–2011	0.07	0.06	0.02	0.04	0.12
Average Robbery Complaint Rate, 2010–2012	49.19	0.00	115.32	0.00	477.00
Total Persons Incarcerated, 2010	553.74	24.00	1924.40	0.00	11,101.00
Proportion Black or Latino, 2010	0.51	0.52	0.29	0.11	0.97
Proportion Below Federal Poverty Line, 2010	0.19	0.16	0.09	0.06	0.39
Affluence Concentration, 2010	7.73	7.00	5.48	0.70	23.09
Proportion of Households More than \$50K, 2010	0.50	0.50	0.13	0.22	0.76
Proportion of Housing Units Owner-Occupied, 2010	0.32	0.31	0.17	0.07	0.73

Note: SD = Standard Deviation.

language spoken at home), socioeconomic (educational attainment, income, work status), and healthcare access (health insurance status, unmet medical care needs) variables of importance to health. Except for age and household size, all individual-level

variables are categorical as described in Table 1. Age is mean-centered at 52.6 years. Household size is median-centered at 2 people; household size is top-coded at 6 (the 97th percentile) to lessen the impact of outliers.




Fig. 2. Distribution of neighborhood post-stop characteristics by United Hospital Fund neighborhoods. Source: 2009–2011 New York City Stop, Question, and Frisk Database; Raw distributions.

1.3. Statistical analysis

The statistical modeling framework employed in this study anticipates that individual reports of illness are partly a function of the UHF to which an individual belongs (Bryk and Raudenbush, 1992). UHF-level variables are appended to each individual observation in the NYC-CHS database, and post-stratification survey weights are applied. Individuals are seen as a representative sample of NYC residents. The UHF, meanwhile, is seen as the population of neighborhoods in NYC. As such, inferences are made about the population of individuals living in NYC neighborhoods.

Stata 14.0 is used for all analyses. Weighted multilevel models for binary and ordinal response outcomes are employed that allow for a random-intercept and assume an unstructured covariance structure. A logit-link is used for the NSPD indicator, whereas a cumulative logit-link is used for the constituent items of the K6 scale, which each has 5 levels. After excluding the missing data on the outcome and predictor variables, we evaluate the relationship between mental health and policing with and without considering individual- and neighborhood-level characteristics. We conduct and report on three sets of regression analyses on 8066 individuals (91.7 percent of sample) nested within 34 NYC UHFs for each outcome-policing pair. The first set of regressions (Overall)

evaluates the association between mental health and policing patterns while controlling for gender to answer Hypothesis 1. The next two sets of regressions pertain to Hypothesis 2. The second set of regressions (Stratified) enables estimates of the health associations of policing to be stratified by gender while assuming that the impacts of control variables are not gender-specific; and a third set of regressions (Interaction) evaluates moderated relationships – that is, whether the mental health associations of policing with psychological distress are different for men and women.

For each set of regressions, four main models are computed. The first model adjusts for race/ethnicity. The second model adjusts for compositional effects by including controls for race/ethnicity and individual-level covariates (Table 1). The third model adjusts for other types of contextual effects by including controls for race/ethnicity and neighborhood-level covariates (Table 2). The fourth model adjusts for race/ethnicity, individual- and neighborhood-level covariates. Statistical significance is assessed for all models at $p < 0.05$.

Supplemental analysis, shown in Appendix A, repeats these sets of regressions and models to assess the mental health associations of the neighborhood stop rate. This analysis shows different relationships than observed for post-stop outcomes. First, there is inconsistency in the main effect of the neighborhood stop rate on mental health. Second, the neighborhood stop rate demonstrates no gendered associations with mental health, as do the neighborhood frisk and use of force proportions.

2. Results

2.1. Descriptive statistics

Tables 1 and 2 provide an unweighted summary of the final data considered for analysis. On average, there are approximately 22 pedestrian stops per 100 non-institutionalized residents of a neighborhood (Table 2). Well over half of pedestrian stops in NYC involve frisking by a police officer, and 22.2 percent involve the use of some kind of force. While a substantial portion of pedestrian stops can be considered “aggressive”, the proportion of stops that are productive is much lower. Only 6.8 percent of pedestrian stops produce an arrest.

Fig. 3 indicates that gender differences in the distribution of NSPD are minute. Women are more likely than men to report feeling sad “all of the time” or “most of the time” within the past 30 days. However, on other dimensions of NSPD, men and women exhibit similar probabilities of reporting distress.

2.2. The neighborhood frisking proportion

Table 3 examines the association between psychological distress and the neighborhood frisking proportion. Weighted odds ratios for the association of frisking with the dichotomous NSPD indicator and the six constituent ordinal psychological distress items are shown under different model specifications described above. Each odds ratio includes, in brackets, the 95% confidence interval for the odds ratio. The first column shows the average association of the stop-question-and-frisk measure using an overall sample of males and females. The second column shows the association of the stop-question-and-frisk measure for males. The third column shows the association of the stop-question-and-frisk measure for females. The fourth column evaluates the statistical significance of the male gender \times neighborhood frisking proportion interaction term and shows the gender difference in


the mental health association of the stop-question-and-frisk measure; this column shows differences in size or direction of effect for men compared to women. For each column and each outcome, the four models are shown vertically, with the naïve model controlling for race/ethnicity (Model 1) shown first and the fully-adjusted model (Model 4) shown last. Models 2 and 3 indicate the mental health associations of neighborhood policing patterns, holding constant individual-level and neighborhood-level covariates, respectively.

In the overall analysis, the neighborhood proportion of pedestrian stops that involve frisking is associated with the dichotomous indicator of NSPD for all models computed ($p < 0.05$). Holding constant both individual- and neighborhood-level characteristics (Model 4), living in a neighborhood where pedestrians are more likely to be frisked increases the odds of reporting NSPD in the past month (AOR = 1.08; 95% CI:1.01–1.15). A standard deviation increase in living in a neighborhood characterized by stops that involve frisking increases the odds of NSPD by 8 percent.

Naïve estimates (Model 1) indicate that the mental health associations of the neighborhood frisking proportion appear to be isolated to select items of the K6 – sadness (OR = 1.12; 95% CI:1.07–1.17), nervousness (OR = 1.04; 95% CI:1.00–1.08), and worthlessness (OR = 1.16; 95% CI:1.08–1.24). Yet, these particular associations do not tend to hold when considering individual-level correlates of mental health and policing patterns (Model 2). The association between worthlessness and the neighborhood frisking proportion is independent of neighborhood-level correlates of mental health and policing patterns (Model 3), but the relationship between neighborhood frisking proportion and sadness and nervousness is attenuated when comparing people in similarly-situated neighborhoods. Holding constant both individual- and neighborhood-level characteristics (Model 4), there is no relationship between the neighborhood frisking proportion and constituent psychological distress items outcomes.

The mental health profiles of men and women (Second and Third Column) are not similarly associated with the neighborhood frisking proportion. For instance, men report more NSPD and more sadness, nervousness, hopelessness, and worthlessness if they live in a neighborhood where pedestrians are more likely to be frisked by police. Naïve estimates (Model 1) for men indicate that a standard deviation increase in the neighborhood frisk proportion (SD = 0.093) is associated with a 20 percent increase in the odds of having NSPD (OR = 1.20; 95% CI:1.07–1.35), a 17 percent increase in the odds of reporting severe feelings of sadness (OR = 1.17; 95% CI:1.07–1.29), a 13 percent increase in the odds of reporting severe feelings of nervousness (OR = 1.13; 95% CI:1.07–1.21), a 19 percent increase in the odds of reporting severe feelings of hopelessness (OR = 1.19; 95% CI:1.10–1.29), and a 25 percent increase in the odds of reporting severe feelings of worthlessness (OR = 1.25; 95% CI:1.15–1.36). The relationships between the neighborhood frisking proportion and sadness and hopelessness are not independent of either compositional or other contextual factors. However, when adjusting for individual- and neighborhood-level characteristics (Model 4), men remain 11 percent more likely to report more severe feelings of nervousness ($p < 0.001$) and 17 percent more likely to report more severe feelings of worthlessness ($p < 0.05$) in neighborhoods with a 9 percentage point increase in the risk of frisking among pedestrians. The psychological profiles of women generally are not shaped by the neighborhood frisking proportion.

The association between neighborhood frisking proportions and



Note: * $p < 0.05$ (two-tailed test for statistically significant gender difference); For constituent items (Columns 1–6) of the non-specific psychological distress indicator (Column 7), the percentage of respondents reporting feeling a specific type of distress “all the time” or “most of the time” in the past 30 days is reported.

Fig. 3. Gender Differences in the Distribution of Psychological Distress. Note: * $p < 0.05$ (two-tailed test for statistically significant gender difference); For constituent items (Columns 1–6) of the non-specific psychological distress indicator (Column 7), the percentage of respondents reporting feeling a specific type of distress “all the time” or “most of the time” in the past 30 days is reported.

Source: 2009–2011 New York City Stop, Question, and Frisk Database; Weighted distributions.

psychological distress is differentiated by gender (Fourth Column). Male gender moderates the psychological associations of the neighborhood frisking proportion for nervousness ($p < 0.001$) and worthlessness ($p < 0.05$). Holding all else constant (Model 4), a standard deviation increase in the neighborhood frisk proportion has a stronger association with the likelihood of men, compared to women, reporting more severe feelings of nervousness (AOR = 1.16; 95% CI: 1.07–1.26) and worthlessness (AOR = 1.16; 95% CI: 1.00–1.34).

2.3. The neighborhood use of force proportion

Table 4 examines the association between psychological distress and the neighborhood proportion of stops that involve use of force, following the same sequence of models as Table 3.

In the overall analysis, the neighborhood proportion of pedestrian stops that involve use of force is not associated with the dichotomous indicator of NSPD for all models computed. Naïve estimates (Model 1) indicate that the mental health associations of the neighborhood frisking proportion appear to be

isolated to select items of the K6 Psychological Distress Scale – nervousness (OR = 1.06; 95% CI: 1.02–1.10), effort (OR = 0.95; 95% CI: 0.91–0.99), and worthlessness (OR = 1.11; 95% CI: 1.04–1.19). Yet, these particular associations do not hold when considering individual-level correlates (Model 2). Model 3 indicates the association between worthlessness and the neighborhood use of force proportion is independent of neighborhood-level correlates, but the relationship between neighborhood use of force proportion and nervousness and effort is attenuated when comparing people in similarly-situated neighborhoods. For sadness and restlessness, comparing people in similarly-situated neighborhoods (Model 3) reveals a protective association between the neighborhood use of force proportion and psychological distress ($p < 0.05$). Holding constant both individual- and neighborhood-level characteristics (Model 4), there is a protective association between the neighborhood use of force proportion and both sadness and effort.

Still, the mental health profiles of men and women are not similarly associated with the neighborhood use of force proportion (Second and Third Columns). Naïve estimates indicate that men

Table 3
Weighted generalized linear multilevel regression of neighborhood frisking proportion on psychological distress for overall sample and for men and women.

	Overall	Stratified		Interaction
		Men	Women	Men × policing
NSPD				
Model 1	1.13 [1.08,1.18]***	1.20 [1.07,1.35]**	1.07 [0.96,1.18]	1.12 [0.93,1.36]
Model 2	1.10 [1.03,1.16]**	1.19 [1.03,1.38]*	1.05 [0.93,1.19]	1.12 [0.92,1.36]
Model 3	1.14 [1.09,1.20]***	1.21 [1.08,1.36]***	1.08 [0.97,1.20]	1.12 [0.93,1.35]
Model 4	1.08 [1.01,1.15]*	1.15 [1.00,1.33]+	1.02 [0.90,1.16]	1.11 [0.92,1.35]
Sadness				
Model 1	1.12 [1.07,1.17]***	1.17 [1.07,1.29]***	1.07 [1.00,1.15]+	1.11 [0.96,1.27]
Model 2	1.02 [0.96,1.08]	1.06 [1.00,1.13]+	0.98 [0.89,1.09]	1.08 [0.95,1.24]
Model 3	1.00 [0.95,1.06]	1.07 [0.98,1.17]	0.98 [0.90,1.06]	1.10 [0.96,1.26]
Model 4	1.00 [0.95,1.06]	1.05 [0.96,1.16]	0.97 [0.89,1.05]	1.09 [0.95,1.25]
Nervousness				
Model 1	1.04 [1.00,1.08]*	1.13 [1.07,1.21]***	0.97 [0.93,1.01]	1.17 [1.07,1.28]***
Model 2	1.03 [0.98,1.07]	1.11 [1.05,1.18]***	0.95 [0.90,1.01]+	1.17 [1.07,1.27]***
Model 3	0.99 [0.96,1.03]	1.10 [1.05,1.15]***	0.94 [0.89,0.99]*	1.17 [1.07,1.27]***
Model 4	1.02 [0.98,1.05]	1.11 [1.06,1.16]***	0.95 [0.90,1.01]+	1.16 [1.07,1.26]***
Restlessness				
Model 1	1.04 [1.00,1.08]+	1.05 [0.97,1.13]	1.03 [0.96,1.12]	1.01 [0.90,1.14]
Model 2	1.02 [0.98,1.06]	1.03 [0.94,1.12]	1.01 [0.93,1.09]	1.02 [0.91,1.14]
Model 3	0.94 [0.89,0.98]**	0.94 [0.86,1.03]	0.93 [0.86,1.01]+	1.01 [0.90,1.14]
Model 4	0.96 [0.92,1.01]	0.97 [0.89,1.06]	0.95 [0.88,1.03]	1.02 [0.91,1.14]
Hopelessness				
Model 1	1.02 [0.97,1.08]	1.19 [1.10,1.29]***	1.02 [0.89,1.18]	1.17 [0.96,1.43]
Model 2	1.02 [0.96,1.08]	1.12 [1.00,1.25]+	0.94 [0.81,1.09]	1.17 [0.97,1.41]+
Model 3	1.04 [0.99,1.10]	1.14 [1.01,1.28]*	0.97 [0.86,1.09]	1.17 [0.96,1.43]
Model 4	1.02 [0.96,1.08]	1.12 [0.96,1.30]	0.94 [0.83,1.07]	1.17 [0.96,1.42]
Effort				
Model 1	1.01 [0.95,1.06]	0.99 [0.92,1.06]	0.93 [0.86,1.00]+	1.07 [0.97,1.19]
Model 2	0.94 [0.90,0.97]***	0.97 [0.89,1.05]	0.91 [0.84,0.99]*	1.07 [0.96,1.19]
Model 3	0.97 [0.92,1.02]	1.00 [0.92,1.10]	0.94 [0.87,1.01]+	1.07 [0.97,1.18]
Model 4	0.97 [0.92,1.01]	1.00 [0.90,1.10]	0.94 [0.87,1.01]+	1.06 [0.96,1.18]
Worthlessness				
Model 1	1.16 [1.08,1.24]***	1.25 [1.15,1.36]***	1.07 [0.97,1.18]	1.17 [1.03,1.34]*
Model 2	1.07 [0.99,1.15]+	1.13 [0.99,1.30]+	0.95 [0.87,1.04]	1.17 [1.01,1.35]*
Model 3	1.12 [1.05,1.19]***	1.23 [1.12,1.36]***	1.05 [0.97,1.15]	1.17 [1.03,1.33]*
Model 4	1.06 [0.97,1.15]	1.17 [1.03,1.31]*	0.98 [0.87,1.11]	1.16 [1.00,1.34]*

+ $p < 0.10$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (two-tailed test; 95% confidence interval shown in brackets).

Source: NYC Community Health Survey ($N_i = 8066$ Residents); NYC Stop, Question, and Frisk Database ($N_j = 34$ United Health Fund Neighborhoods). Model 1 includes for race/ethnicity and, where appropriate (Overall and Interaction), gender. Model 2 adds individual-level controls to Model 1, including age, education, employment, nativity, insurance, unmet medical need, household size, marital status, and language. Model 3 adds neighborhood-level controls to Model 1, including the proportion of stops resulting in arrest, average robbery complaint rate, total persons incarcerated, proportion Black or Latino, proportion below federal poverty line, and affluence concentration. Model 4 adds individual-level and neighborhood-level controls to Model 1.

report more NSPD and more sadness, nervousness, hopelessness, and worthlessness if they live in a neighborhood where pedestrians are more likely to have force used against them by police. Model 1 indicates that for men a standard deviation increase in the neighborhood use of force proportion ($SD = 0.077$) is associated with a 23 percent increase in the odds of having NSPD ($OR = 1.23$; 95% $CI: 1.11–1.37$), a 10 percent increase in the odds of reporting severe feelings of sadness ($OR = 1.10$; 95% $CI: 1.01–1.19$), a 13 percent increase in the odds of reporting severe feelings of nervousness ($OR = 1.13$; 95% $CI: 1.06–1.22$), a 15 percent increase in the odds of reporting severe feelings of hopelessness ($OR = 1.15$; 95% $CI: 1.02–1.28$), and a 23 percent increase in the odds of reporting severe feelings of worthlessness ($OR = 1.23$; 95% $CI: 1.08–1.40$). The relationship between the neighborhood use of force proportion and worthlessness is independent of neighborhood-level covariates but attenuates when comparing people with similar demographic characteristics. When adjusting for individual- and neighborhood-level characteristics (Model 4), men remain 7 percent more likely to report more severe feelings of nervousness in neighborhoods with a 7.7 percentage point increase in the risk of use of force among pedestrians.

Meanwhile, the psychological profiles of women are generally not shaped by the neighborhood use of force proportion. If anything, women in similarly-situated neighborhoods report less severe feelings of sadness and effortfulness if they live in neighborhoods where pedestrians face a higher risk of having force used against them. Holding all else constant (Model 4), a standard deviation increase in the neighborhood use of force proportion decreases severe feelings of sadness by 13 percent ($AOR = 0.87$; 95% $CI: 0.80–0.94$) and severe feelings of effort by 14 percent ($AOR = 0.80$; 95% $CI: 0.80–0.92$).

The association between neighborhood use of force proportions and psychological distress is differentiated by gender (Fourth Column). In fact, gender differences in the relationship between the neighborhood use of force proportion and psychological distress are more pronounced than gender differences in the relationship between the neighborhood frisking proportion and psychological distress. Holding all else constant (Model 4), male gender moderates the psychological associations of the neighborhood use of force proportion for the dichotomous NSPD indicator, nervousness, effort, and worthlessness ($p < 0.05$). Compared to women, a standard deviation increase in the neighborhood use of force

Table 4

Weighted generalized linear multilevel regression of neighborhood use of force proportion on psychological distress for overall sample and for men and women.

	Overall	Stratified		Interaction
		Men	Women	Men × policing
NSPD				
Model 1	1.02 [0.95,1.08]	1.23 [1.11,1.37]***	1.01 [0.92,1.11]	1.22 [1.04,1.43]*
Model 2	0.99 [0.92,1.06]	1.10 [0.99,1.22]+	0.90 [0.79,1.03]	1.20 [1.02,1.41]*
Model 3	1.01 [0.94,1.09]	1.12 [1.02,1.24]*	0.92 [0.82,1.04]	1.21 [1.03,1.42]*
Model 4	1.05 [0.98,1.12]	1.17 [1.05,1.31]**	0.95 [0.84,1.08]	1.21 [1.02,1.43]*
Sadness				
Model 1	1.06 [1.00,1.12]+	1.10 [1.01,1.19]*	0.99 [0.89,1.09]	1.13 [0.99,1.30]+
Model 2	1.03 [0.97,1.09]	1.07 [0.99,1.15]+	0.96 [0.86,1.07]	1.12 [0.96,1.29]
Model 3	0.92 [0.88,0.97]**	0.99 [0.90,1.08]	0.88 [0.80,0.96]**	1.14 [0.99,1.31]+
Model 4	0.91 [0.87,0.96]***	0.97 [0.88,1.07]	0.87 [0.80,0.94]***	1.12 [0.97,1.30]
Nervousness				
Model 1	1.06 [1.02,1.10]**	1.13 [1.06,1.22]***	1.00 [0.94,1.06]	1.14 [1.02,1.27]*
Model 2	0.98 [0.93,1.02]	1.10 [1.03,1.18]**	0.97 [0.90,1.04]	1.14 [1.03,1.27]*
Model 3	1.00 [0.97,1.03]	1.08 [1.01,1.15]*	0.95 [0.89,1.00]+	1.14 [1.02,1.27]*
Model 4	1.00 [0.97,1.04]	1.07 [1.02,1.13]**	0.94 [0.88,1.01]+	1.14 [1.03,1.26]*
Restlessness				
Model 1	1.02 [0.97,1.06]	1.05 [0.96,1.15]	0.99 [0.92,1.06]	1.06 [0.93,1.20]
Model 2	1.01 [0.96,1.05]	1.04 [0.95,1.15]	0.98 [0.91,1.05]	1.06 [0.94,1.20]
Model 3	0.96 [0.93,1.00]*	0.99 [0.91,1.08]	0.94 [0.88,1.00]+	1.05 [0.93,1.19]
Model 4	0.96 [0.93,1.00]+	1.02 [0.93,1.11]	0.95 [0.89,1.03]	1.06 [0.94,1.20]
Hopelessness				
Model 1	1.05 [1.00,1.11]+	1.15 [1.02,1.28]*	0.94 [0.83,1.07]	1.22 [0.99,1.50]+
Model 2	1.00 [0.94,1.06]	1.12 [0.97,1.28]	0.91 [0.78,1.05]	1.21 [0.97,1.49]+
Model 3	1.00 [0.95,1.05]	1.11 [0.97,1.27]	0.91 [0.81,1.01]+	1.22 [0.99,1.51]+
Model 4	0.99 [0.94,1.05]	1.11 [0.95,1.30]	0.90 [0.80,1.02]	1.21 [0.97,1.50]+
Effort				
Model 1	0.95 [0.91,0.99]*	0.99 [0.92,1.06]	0.87 [0.81,0.93]***	1.13 [1.03,1.24]**
Model 2	1.00 [0.95,1.05]	1.08 [0.97,1.19]	0.94 [0.88,1.01]+	1.13 [1.02,1.25]*
Model 3	0.99 [0.95,1.04]	1.07 [0.99,1.16]+	0.94 [0.88,1.00]+	1.13 [1.03,1.24]**
Model 4	0.91 [0.87,0.95]***	0.97 [0.89,1.06]	0.86 [0.80,0.92]***	1.12 [1.02,1.24]*
Worthlessness				
Model 1	1.11 [1.04,1.19]**	1.23 [1.08,1.40]**	1.02 [0.96,1.08]	1.21 [1.05,1.41]*
Model 2	1.02 [0.95,1.09]	1.14 [0.99,1.30]+	0.93 [0.85,1.02]	1.20 [1.02,1.40]*
Model 3	1.10 [1.03,1.17]**	1.22 [1.10,1.36]***	1.02 [0.94,1.11]	1.20 [1.04,1.39]*
Model 4	0.99 [0.91,1.08]	1.10 [0.98,1.23]+	0.91 [0.80,1.04]	1.18 [1.01,1.38]*

+ $p < 0.10$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (two-tailed test; 95% confidence interval shown in brackets).

Source: NYC Community Health Survey ($N_i = 8,066$ Residents); NYC Stop, Question, and Frisk Database ($N_j = 34$ United Health Fund Neighborhoods). Model 1 includes for race/ethnicity and, where appropriate (Overall and Interaction), gender. Model 2 adds individual-level controls to Model 1, including age, education, employment, nativity, insurance, unmet medical need, household size, marital status, and language. Model 3 adds neighborhood-level controls to Model 1, including the proportion of stops resulting in arrest, average robbery complaint rate, total persons incarcerated, proportion Black or Latino, proportion below federal poverty line, and affluence concentration. Model 4 adds individual-level and neighborhood-level controls to Model 1.

proportion has a stronger association with men reporting high levels of NSPD (AOR = 1.21; 95% CI:1.02–1.43) and more severe feelings of nervousness (AOR = 1.14; 95% CI:1.03–1.26), effort (AOR = 1.12; 95% CI:1.02–1.24), and worthlessness (AOR = 1.18; 95% CI:1.01–1.38). Living in a community where pedestrian stops are likely to involve use of force by police is associated with NSPD and more severe reports of nervousness, effort, and worthlessness more so for male than for female respondents.

3. Discussion

This paper provides a starting point to evaluate the relationship between psychological distress and the escalation of pedestrian stops across the 34 UHF neighborhoods of NYC. Underreporting of frisking and use of force by police (Spitzer, 1999) may attenuate the mental health associations of neighborhood-level policing patterns. Yet, we identify an association between escalated police encounters, measured at the neighborhood-level, and higher psychological distress for men, measured at the individual-level, using multilevel models that adjust for complaints of criminal activity, racial/ethnic/class composition, and known sociodemographic correlates of mental health. This study indicates that po-

lice surveillance matters not only for men who have contact with police (Geller et al., 2014), but also for men living in highly policed areas. Living in a neighborhood with a higher density of frisking is associated with experiencing more severe psychological distress for all residents, while living in a neighborhood with a higher density of use of force is associated with experiencing fewer feelings of sadness and effort.

The psychological associations of living under surveillance, however, are gendered. Men who live in neighborhoods where pedestrians are more likely to be frisked by police are more likely to report feelings of nervousness and worthlessness and more severe psychological distress. Men who live in neighborhoods where pedestrians are more likely to have force used against them by police are more likely to report feelings of nervousness, effort, and worthlessness and more severe psychological distress. Women, however, are not affected psychologically by living in a neighborhood with a high density of frisking, and, in some cases, living in a high use of force neighborhood is protective of women's mental health.

The reasons why some patterns of police surveillance may be protective for women is unclear and cannot be discerned from this study design. While women may also be frustrated about police

neglect and abuse (Cooper et al., 2004; Lee and Wildeman, 2013), Black mothers, in particular, report wanting a stronger police presence in their neighborhood, so as to mitigate against concerns about neighborhood violence and safety (Dias and Whitaker, 2013). Ideally, we would have been able to include individual-level measures of perceived neighborhood safety and crime in this analysis, but such measures were not available. Nonetheless, women may feel safer in neighborhoods where police are more likely to aggressively treat pedestrians. Possibly, when police have a strong presence in a neighborhood, men may be less likely to exhibit aggressive behavior towards women, such as cat-calling. However, such feelings of safety among women, if present, do not transfer to men living in such neighborhoods. Moreover, these analyses suggest that the protective association of stop-and-frisk policing is specific to certain dimensions of mental health, such as feelings of restlessness and worthlessness.

Another possibility for these counterintuitive findings is that women may be more vulnerable to clinical indicators of mental health than men. For instance, a recent study of the K6 found men endorsed items less often than women with similar underlying levels of distress (Drapeau et al., 2010). Future research should explore additional measures of mental health and well-being to ascertain the dimensionality of stop-and-frisk policing on women's mental health.

Combining different racial and ethnic status groups obscures points of vulnerability among women of color, who are not immune to police stops. Specifically, previous research indicates that the health effects of stop-question-and-frisk policing may vary by race (Sewell and Jefferson, 2016). Moreover, incarceration is associated with a loss to the family, so it will have repercussive effects on all family members, particularly women who take on a larger care-taking burden (Lee and Wildeman, 2013; Lee et al., 2014). Given the disproportionate representation of minorities in the criminal justice system and the tendency of police to stop minority men on the street, future research should examine whether race or ethnicity buffers or amplifies the gendered mental health associations of neighborhood stop-question-and-frisk policing. Still, minority men may be more vulnerable to the mental health associations of neighborhood frisking and use of force proportions, thus leading to strengthened associations for Black, Latino, and Asian men. Yet, because police contact is so frequent for minority men, they may normalize this experience, leading to weakened associations. Such findings would align with research showing attenuated relationships between perceived discrimination and mental health for minorities (Kessler et al., 1999).

While this study demonstrates an association between contextual patterns of policing and emotional well-being for the community at-large, this study has limitations. First, the NYC-CHS does not include measures of police contact for respondents. It is possible that police are more likely to stop the mentally ill (Engel and Silver, 2001; Fisher et al., 2006) or that displays of emotionality and erratic behavior may precipitate police aggression. However, time ordering the policing measures to occur within the three years prior to the collection of mental health data situates our results as useful for understanding the association between current health status and living in neighborhoods that have been highly surveilled.

Second, this analysis relies on police-reported data. It is possible that police may underreport their use of force, particularly against Black and Latino individuals. Should this be the case, this would likely bias our results towards a null hypothesis and, thus, would not threaten the validity of our observation that policing associates

with indicators of distress.

Third, community factors not considered in this study, such as the density of parks or the quality of available social services, may affect the relationship between police stop-and-frisk actions and psychological distress. While we do not directly control for specific measures of neighborhood quality, we do attempt to control for neighborhood socioeconomic status to help lessen this concern. However, the absence of individual-level perceptions of neighborhood crime and safety, as well as individual-level perceptions of collective efficacy and social cohesion, represent important subjective measures missing from this study. Although these measures were not available for inclusion in our analyses, we do include an objective measure of neighborhood crime (robbery complaints) that may help explain how neighborhood safety and social disorder could affect relationships between policing and psychological distress.

Fourth, this study explores a limited range of neighborhood policing patterns. Adjusted and unadjusted analyses showed that the stop rate itself is not associated with indicators of NSPD. However, other patterns to consider are the relative risk of being searched, receiving a summons, and undergoing an arrest. Moreover, exposure in this study is considered in relation to the number of stops executed in a community. Exposure could also be calculated relative to the population size or land area. Relatedly, a wider range of localities should be considered in future research. Each UHF neighborhood represents a large swath of people and land-mass. In all, NYC has over two thousand census tracts covering nearly one hundred acres. There may be substantial variation in the association of neighborhood policing patterns within UHF neighborhoods across smaller locales or egocentric neighborhoods.

Nonetheless, this study uses a multilevel design with distinct data sources corresponding to separate levels and draws attention to the relationship between neighborhood policing patterns and psychological distress. Just as mass incarceration has broader consequences beyond incarcerated individuals (Frank et al., 2013; Hatzenbuehler et al., 2014), police actions within a neighborhood also affect the health of the community at-large. Our findings are consistent with recent research that identifies ill health associations of stop-and-frisk policing (Geller et al., 2014; Sewell and Jefferson, 2016). The actions and inactions of police spillover to shape the health and well-being of communities. Our findings show that living in aggressively policed communities is of detriment to the health of male residents in the neighborhood. As such, this study highlights an important tension between violence prevention, which is a public health issue, and over-policing, which we argue is a health risk factor.

Acknowledgments

This work was completed while the first author was a Vice Provost's Postdoctoral Fellow at the University of Pennsylvania.

Appendix A

Mental health, gender, and the neighborhood stop rate

As stopping pedestrians is the most basic action necessary to implement NYC's Stop, Question, and Frisk policy, we also consider the mental health associations of the neighborhood stop rate. The neighborhood stop rate is calculated as the number of pedestrian stops per 100 non-institutionalized residents of a neighborhood. Fig. A-1 provides a map of the distribution of the neighborhood stop rate.




Fig. A-1. Distribution of Neighborhood Stop Rate by United Hospital Fund Neighborhoods.

Table A-1 examines the association between psychological distress and the neighborhood rate of pedestrian stops per the non-institutionalized population, following the same sequence of models as Tables 3 and 4.

Table A-1

. Weighted Generalized Linear Multilevel Regression of Neighborhood Stop Rate on Psychological Distress for Overall Sample and for Men and Women.

	Overall	Stratified		Interaction Men × policing
		Men	Women	
NSPD				
Model 1	1.04 [0.99,1.10]	1.07 [0.94,1.21]	1.02 [0.92,1.14]	1.04 [0.85,1.28]
Model 2	0.94 [0.88,1.01]+	0.96 [0.81,1.14]	0.93 [0.85,1.03]	1.04 [0.83,1.30]
Model 3	0.97 [0.90,1.05]	1.00 [0.87,1.15]	0.96 [0.86,1.09]	1.04 [0.84,1.28]
Model 4	0.97 [0.92,1.04]	1.00 [0.86,1.16]	0.96 [0.86,1.08]	1.04 [0.83,1.30]
Sadness				
Model 1	1.06 [1.01,1.12]*	1.11 [1.02,1.21]*	1.05 [0.96,1.14]	1.05 [0.95,1.17]
Model 2	0.98 [0.93,1.02]	1.01 [0.93,1.09]	0.95 [0.90,1.01]+	1.06 [0.96,1.17]
Model 3	1.02 [0.96,1.08]	1.05 [0.96,1.15]	0.99 [0.91,1.08]	1.06 [0.95,1.17]
Model 4	1.00 [0.94,1.07]	0.98 [0.88,1.09]	0.92 [0.85,1.01]+	1.06 [0.96,1.17]
Nervousness				
Model 1	0.93 [0.89,0.97]***	0.97 [0.90,1.04]	0.89 [0.83,0.96]**	1.07 [0.97,1.19]

(continued on next page)

Table A-1 (continued)

	Overall	Stratified		Interaction Men × policing
		Men	Women	
Model 2	0.91 [0.86,0.95]***	0.94 [0.87,1.01]	0.86 [0.80,0.92]***	1.09 [0.99,1.21]+
Model 3	1.05 [1.00,1.10]*	1.09 [1.01,1.19]*	1.01 [0.95,1.07]	1.08 [0.97,1.19]
Model 4	0.97 [0.92,1.03]	1.08 [1.00,1.17]+	0.99 [0.93,1.05]	1.09 [0.98,1.21]
Restlessness				
Model 1	1.06 [1.01,1.11]*	1.07 [0.98,1.17]	1.02 [0.96,1.09]	1.05 [0.93,1.20]
Model 2	0.98 [0.93,1.02]	1.02 [0.92,1.13]	0.97 [0.91,1.03]	1.06 [0.93,1.21]
Model 3	0.94 [0.91,0.98]***	0.97 [0.89,1.05]	0.92 [0.86,0.99]*	1.06 [0.93,1.20]
Model 4	0.89 [0.84,0.94]***	0.91 [0.82,1.02]	0.87 [0.81,0.93]***	1.06 [0.93,1.22]
Hopelessness				
Model 1	1.13 [1.06,1.20]***	1.14 [1.03,1.26]**	1.12 [1.03,1.21]**	1.02 [0.90,1.16]
Model 2	1.01 [0.96,1.06]	1.03 [0.92,1.15]	1.00 [0.93,1.07]	1.03 [0.91,1.17]
Model 3	1.06 [0.99,1.13]+	1.07 [0.96,1.19]	1.05 [0.97,1.14]	1.02 [0.90,1.16]
Model 4	0.99 [0.93,1.06]	0.97 [0.86,1.08]	0.94 [0.87,1.02]	1.03 [0.91,1.17]
Effort				
Model 1	1.06 [1.01,1.11]*	1.09 [0.99,1.21]+	1.08 [1.01,1.15]*	1.02 [0.89,1.17]
Model 2	1.02 [0.97,1.08]	1.04 [0.93,1.16]	1.02 [0.95,1.09]	1.04 [0.90,1.19]
Model 3	0.97 [0.93,1.03]	0.94 [0.84,1.05]	0.93 [0.87,1.00]*	1.02 [0.89,1.17]
Model 4	0.98 [0.94,1.03]	0.99 [0.90,1.09]	0.97 [0.90,1.06]	1.03 [0.90,1.19]
Worthlessness				
Model 1	1.07 [0.99,1.15]	1.14 [1.02,1.28]	1.03 [0.93,1.13]	1.11 [0.96,1.29]
Model 2	1.00 [0.93,1.07]	1.05 [0.91,1.21]	0.96 [0.90,1.03]	1.09 [0.92,1.30]
Model 3	0.88 [0.83,0.93]***	0.90 [0.79,1.02]+	0.81 [0.73,0.89]***	1.11 [0.96,1.29]
Model 4	0.93 [0.85,1.02]	0.98 [0.85,1.13]	0.90 [0.82,0.98]*	1.09 [0.92,1.29]

Source: NYC Community Health Survey ($N_i = 8066$ Residents); NYC Stop, Question, and Frisk Database ($N_j = 34$ United Health Fund Neighborhoods). Model 1 includes for race/ethnicity and, where appropriate (Overall and Interaction), gender. Model 2 adds individual-level controls to Model 1, including age, education, employment, nativity, insurance, unmet medical need, household size, marital status, and language. Model 3 adds neighborhood-level controls to Model 1, including the proportion of stops resulting in arrest, average robbery complaint rate, total persons incarcerated, proportion Black or Latino, proportion below federal poverty line, and affluence concentration. Model 4 adds individual-level and neighborhood-level controls to Model 1.

+ $p < 0.10$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (two-tailed test; 95% confidence interval shown in brackets).

In the overall analysis, the neighborhood rate of stopping pedestrian is not associated with the dichotomous indicator of NSPD for any of the models computed. However, the neighborhood stop rate is associated with five of the six constituent items, but not always in the expected direction. Controlling for race/ethnicity and gender (Model 1), a standard deviation increase in the neighborhood stop rate ($SD = 14.95$) increases the odds of reporting more severe feelings of sadness, restlessness, hopelessness, and effortfulness by at least 6 percent and decreases the odds of reporting more severe feelings of nervousness by 7 percent. Living in a neighborhood with a higher stop rate is associated with a higher odds of reporting sadness ($OR = 1.06$; 95% $CI: 1.01–1.12$), restlessness ($OR = 1.06$; 95% $CI: 1.01–1.11$), hopelessness ($OR = 1.13$; 95% $CI: 1.06–1.20$), and effort ($OR = 1.06$; 95% $CI: 1.01–1.11$) but a lower odds of reporting feelings of nervousness ($OR = 0.93$; 95% $CI: 0.89–0.97$) in the past month.

These relationships do not hold when considering compositional attributes of highly surveilled neighborhoods or other contextual characteristics. Specifically, when individual level covariates (Model 2) or neighborhood level covariates (Model 3) are added the associations between the neighborhood stop rate and sadness, restlessness, hopelessness, and effort dissipate. The aforementioned protective association between the neighborhood stop rate and nervousness also dissipates once both individual- and neighborhood-level correlates are considered (Model 4). While a protective association between the neighborhood stop rate and restlessness in the past month appears ($OR = 0.89$; 95% $CI: 0.84–0.94$), this association may be an artifact of comparing people in similarly-situated neighborhoods (Model 3) and holds when further controlling for the demographic composition of people living in neighborhoods with higher stop rates (Model 4).

Moreover, contrary to expectation, the psychological associations of the neighborhood stop rate are not differentiated by gender. In models that adjust for race/ethnicity only (Model 1), there are gender-specific psychological distress associations with

the neighborhood stop rate. For instance, men (Second Column) do report more sadness ($OR = 1.11$; 95% $CI: 1.02–1.21$), hopelessness ($OR = 1.14$; 95% $CI: 1.03–1.26$), and worthlessness ($OR = 1.14$; 95% $CI: 1.02–1.28$) in neighborhoods with higher stop rates. Furthermore, women (Third Column) do report less severe feelings of nervousness ($OR = 0.89$; 95% $CI: 0.83–0.96$). Yet, women also report more severe feelings of hopelessness ($OR = 1.12$; 95% $CI: 1.03–1.21$) and effort ($OR = 1.08$; 95% $CI: 1.01–1.15$) in neighborhoods with a higher stop rate. Nevertheless, these associations are attenuated once individual- and neighborhood-level covariates are included in Models 2 and 3, respectively.

Holding all else constant, men are not affected by the neighborhood stop rate (Model 4). Meanwhile, women report less severe feelings of restlessness ($OR = 0.87$; 95% $CI: 0.81–0.93$) and worthlessness ($OR = 0.90$; 95% $CI: 0.82–0.98$) in neighborhoods with higher stop rates. Yet, the absence of a statistically significant multilevel association for the male gender × neighborhood stop rate interaction term (Men-Women: Fourth Column) indicates there is not enough evidence to suggest a difference between men and women.

References

- Anderson, E., 1990. *Streetwise: Race, Class, and Change in an Urban Community*. University of Chicago Press.
- Anderson, E., 1998. The police and the black male. In: Anderson, M.L., Collins, P.H. (Eds.), *Race, Class, and Gender: an Anthology*, pp. 476–480.
- Anderson, E., 2003. *A Place on the Corner*. University of Chicago Press.
- Balko, R., 2015. The Black Lives Matter policy agenda is practical, thoughtful - and urgent. *Wash. Post*.
- Browning, C.R., Cagney, K.A., 2003. Moving beyond poverty: neighborhood structure, social processes, and health. *J. Health Soc. Behav.* 44 (4), 552–571. Retrieved from: <http://www.jstor.org/stable/1519799>.
- Brunson, R.K., Miller, J., 2006. Gender, race, and urban policing: the experience of African American youths. *Gend. Soc.* 20 (4), 531–552. <http://dx.doi.org/10.1177/0891243206287727>.
- Brunson, R.K., Weitzer, R., 2008. Police relations with black and white youths in different urban neighborhoods. *Urban Aff. Rev.* <http://dx.doi.org/10.1177/>

- 1078087408326973.
- Bryk, A.S., Raudenbush, S.W., 1992. *Hierarchical Linear Models: Applications and Data Analysis Methods*. Sage Publications, Inc.
- Cohen, S., Evans, G., Stokols, D., Krantz, D., 1986. Stress processes and the costs of coping. In: *Behavior, Health, and Environmental Stress*. Springer US, pp. 1–23.
- Cooper, H., Moore, L., Gruskin, S., Krieger, N., 2004. Characterizing perceived police violence: implications for public health. *Am. J. Public Health* 94 (7), 1109–1118. <http://dx.doi.org/10.2105/AJPH.94.7.1109>.
- Dias, J.J., Whitaker, R.C., 2013. Black mothers' perceptions about urban neighborhood safety and outdoor play for their preadolescent daughters. *J. Health Care Poor Underserved* 24 (1), 206–219.
- Dottolo, A.L., Stewart, A.J., 2008. "Don't ever forget now, You're a black man in America": intersections of race, class and gender in encounters with the police. *Sex Roles* 59 (5–6), 350–364. <http://dx.doi.org/10.1007/s11199-007-9387-x>.
- Drapeau, A., Beaulieu-Prévost, D., Marchand, A., Boyer, R., Prévile, M., Kairouz, S., 2010. A life-course and time perspective on the construct validity of psychological distress in women and men. Measurement invariance of the K6 across gender. *BMC Med. Res. Methodol.* 10 (1), 1–16. <http://dx.doi.org/10.1186/1471-2288-10-68>.
- Engel, R.S., Calnon, J.M., 2004. Examining the influence of drivers' characteristics during traffic stops with police: results from a national survey. *Justice Q.* 21 (1), 49–90. <http://dx.doi.org/10.1080/07418820400095741>.
- Engel, R.S., Silver, E., 2001. Policing mentally disordered subjects: a reexamination of the criminalization hypothesis. *Criminology* 39 (2), 225–252. <http://dx.doi.org/10.1111/j.1745-9125.2001.tb00922.x>.
- Fisher, W.H., Silver, E., Wolff, N., 2006. Beyond criminalization: toward a criminologically informed framework for mental health policy and services research. *Adm. Policy Ment. Health Ment. Health Serv. Res.* 33 (5), 544–557. <http://dx.doi.org/10.1007/s10488-006-0072-0>.
- Frank, J.W., Hong, C.S., Subramanian, S.V., Wang, E.A., 2013. Neighborhood incarceration rate and asthma prevalence in New York City: a multilevel approach. *Am. J. Public Health* 103 (5), e38–e44. <http://dx.doi.org/10.2105/AJPH.2013.301255>.
- Geller, A., Fagan, J., Tyler, T., Link, B.G., 2014. Aggressive policing and the mental health of young urban men. *Am. J. Public Health* 104 (12), 2321–2327. <http://dx.doi.org/10.2105/AJPH.2014.302046>.
- Goffman, A., 2009. On the run: wanted men in a Philadelphia ghetto. *Am. Sociol. Rev.* 74 (3), 339–357. <http://dx.doi.org/10.1177/000312240907400301>.
- Goodman, J.D., 2015. As shootings rise in New York, police focus on a small number of young men. *N. Y. Times*.
- Hatzenbuehler, M.L., Keyes, K., Hamilton, A., Uddin, M., Galea, S., 2014. The collateral damage of mass incarceration: risk of psychiatric morbidity among non-incarcerated residents of high-incarceration neighborhoods. *Am. J. Public Health* 105 (1), 138–143. <http://dx.doi.org/10.2105/AJPH.2014.302184>.
- Jones, N., 2014. "The regular routine": proactive policing and adolescent development among young, poor black men. *New Dir. Child Adolesc. Dev.* 143, 33–54. <http://dx.doi.org/10.1002/cad.20053>.
- Kessler, R.C., Barker, P.R., Colpe, L.J., Epstein, J.F., Gfroerer, J.C., Hiripi, E., et al., 2003. Screening for serious mental illness in the general population. *Arch. General Psychiatry* 60 (2), 184–189. <http://dx.doi.org/10.1001/archpsyc.60.2.184>.
- Kessler, R.C., Mickelson, K.D., Williams, D.R., 1999. The prevalence, distribution, and mental health correlates of perceived discrimination in the United States. *J. Health Soc. Behav.* 40 (3), 208–230. <http://dx.doi.org/10.2307/2676349>.
- Lee, H., Wildeman, C., 2013. Things fall apart: health consequences of mass imprisonment for African American women. *Rev. Black Polit. Econ.* 40 (1), 39–52.
- Lee, H., Wildeman, C., Wang, E.A., Matusko, N., Jackson, J.S., 2014. A heavy burden: the cardiovascular health consequences of having a family member incarcerated. *Am. J. Publ. Health* 104 (3), 421–427.
- Leovy, J., January 24, 2015 Saturday. The underpolicing of black America. *Wall Str. J.* Abstr. C; Pg. 3, 21 words <http://www.wsj.com/articles/the-underpolicing-of-black-america-1422049080>.
- Lerman, A.E., Weaver, V., 2014. Staying out of sight? Concentrated policing and local political action. *Ann. Am. Acad. Polit. Soc. Sci.* 651 (1), 202–219. <http://dx.doi.org/10.1177/0002716213503085>.
- Lewis, T., 2015. ACLU finds Chicago police use stop-and-frisk four times more often than NYPD. *Essence*. <http://www.essence.com/2015/03/25/aclu-chicago-police-stop-and-frisk-four-times-nypd>.
- Massoglia, M., 2008. Incarceration as exposure: the prison, infectious disease, and other stress-related illnesses. *J. Health Soc. Behav.* 49 (1), 56–71. Retrieved from: <http://hsb.sagepub.com/content/49/1/56.full.pdf>.
- Massoglia, M., Pridemore, W.A., 2015. Incarceration and health. *Annu. Rev. Sociol.* 41 (1).
- McEwen, B.S., 2004. Protection and damage from acute and chronic stress: allostasis and allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Ann. N. Y. Acad. Sci.* 1032 (1), 1–7. <http://dx.doi.org/10.1196/annals.1314.001>.
- McVeigh, K.H., Galea, S., Thorpe, L.E., Malsby, C., Henning, K., Sederer, L.I., 2006. The epidemiology of nonspecific psychological distress in New York City, 2002 and 2003. *J. Urban Health* 83 (3), 394–405. <http://dx.doi.org/10.1007/s11524-006-9049-2>.
- Meyer, I.H., 1995. Minority stress and mental health in gay men. *J. Health Soc. Behav.* 36 (1), 38–56. <http://dx.doi.org/10.2307/2137286>.
- New York City Department of Health and Mental Hygiene, 2012. *Community Health Survey* 2012.
- New York City Police Department, 2009–2011. 2009, 2010, 2011 Stop, Question, and Frisk Data.
- NYC Department of Health and Mental Hygiene, 2006. *New York City United Hospital Fund (UHF) Neighborhoods and NYC ZIP Code Areas*.
- Ridgeway, G., 2007. *Analysis of Racial Disparities in the New York Police Department's Stop, Question, and Frisk Practices*. Rand Corporation.
- Sewell, A.A., Jefferson, K.A., 2016. Collateral damage: the health effects of invasive police encounters in New York City. *J. Urban Health* 93 (Supplement 1), 42–67.
- Shedd, C., 2012. What about the other 99%? The broader impact of street stops on minority communities. In: La Vigne, N., Lachman, P., Matthews, A., Neusteter, S.R. (Eds.), *Key Issues in the Police Use of Pedestrian Stops and Searches: Discussion Papers from an Urban Institute Roundtable*. The Urban Institute, New York, pp. 24–29.
- Spitzer, E., 1999. *The New York City Police Department's Stop & Frisk Practices: a Report to the People of the State of New York from the Office of the Attorney General*. DIANE Publishing.
- The Associated Press, 2015. Chicago stop-and-frisk to be monitored. *N. Y. Times*.
- Unnever, J.D., Gabbidon, S.L., 2011. *A Theory of African American Offending: Race, Racism, and Crime*. Taylor & Francis.
- Wildeman, C., Schnittker, J., Turney, K., 2012. Despair by association? The mental health of mothers with children by recently incarcerated fathers. *Am. Sociol. Rev.* 77 (2), 216–243. Retrieved from: <http://asr.sagepub.com/content/77/2/216.abstract>.
- Williams, D.R., Mohammed, S.A., 2009. Discrimination and racial disparities in health: evidence and needed research. *J. Behav. Med.* 32 (1), 20–47. <http://dx.doi.org/10.1007/s10865-008-9185-0>.
- Williams, D.R., Neighbors, H.W., Jackson, J.S., 2003. Racial/ethnic discrimination and health: findings from community studies. *Am. J. Public Health* 93 (2), 200–208. <http://dx.doi.org/10.2105/AJPH.93.2.200>.
- Williams, D.R., Yan, Y., Jackson, J.S., Anderson, N.B., 1997. Racial differences in physical and mental health: socio-economic status, stress and discrimination. *J. Health Psychol.* 2 (3), 335–351. Retrieved from: <http://hpq.sagepub.com/content/2/3/335.abstract>.
- Young, A.A., 2006. *The Minds of Marginalized Black Men: Making Sense of Mobility, Opportunity, and Future Life Chances*. Princeton University Press.