

Nested ANOVA and MANOVA

Rui Zhou

目录

1	Nested ANOVA	1
1.1	0	1
1.2	visualization	2
1.3	nested ANOVA	2
2	MANOVA	5
2.1	0	5
2.2	visualization	6
2.3	assumptions	8
2.4	MANOVA	11
2.5	effect size	12
2.6	univariate ANOVA	12
2.7	post-hoc tests	13

1 Nested ANOVA


1.1 0

```
library(tidyverse)
library(car)
```

```
nestedData = readxl::read_excel("nested anova.xlsx")
```

1.2 visualization

```
# we should eyeball the data first to make sense of the test.  
library(ggplot2)  
# create boxplots to visualize the data  
ggplot(nestedData, aes(x=factor(location), y=DV)) +  
  geom_boxplot()
```



1.3 nested ANOVA

Theoretically, the order of computing the variance sum of squares matters, thus appointing the `summation = 'I'`.

However, here the summation type does not matter.

Note that the result of the main effect of the group is wrong. We can manually make it straight from the SS of the group and the subgroup.

```
# note that nested anova can use error type I and III.
# aov() has a default type I to process factors in an sequential order
# here type I and III don't differ, since effectively we have no orders
aov_model = aov(DV ~ factor(method)/factor(location),
               data = nestedData,summation="I")
```

```
## Warning: In lm.fit(x, y, offset = offset, singular.ok = singular.ok, ...) :
## extra argument 'summation' will be disregarded
```

```
summary(aov_model)
```

```
##                                Df Sum Sq Mean Sq F value    Pr(>F)
## factor(method)                  2  665.7    332.8    255.7 1.45e-10 ***
## factor(method):factor(location)  9 1720.7    191.2    146.9 6.98e-11 ***
## Residuals                       12   15.6      1.3
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
aov_model = aov(DV ~ factor(method)/factor(location),
               data = nestedData,summation="III")
```

```
## Warning: In lm.fit(x, y, offset = offset, singular.ok = singular.ok, ...) :
## extra argument 'summation' will be disregarded
```

```
summary(aov_model)
```

```
##                                Df Sum Sq Mean Sq F value    Pr(>F)
## factor(method)                  2  665.7    332.8    255.7 1.45e-10 ***
## factor(method):factor(location)  9 1720.7    191.2    146.9 6.98e-11 ***
## Residuals                       12   15.6      1.3
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

1.3.1 error adjustment

An alternative refinement is to adjust the error term.

```
aov.result2 = aov(DV ~ factor(method) +
                  Error(factor(method):factor(location)),
                  data = nestedData)
```

```
## Warning in aov(DV ~ factor(method) + Error(factor(method):factor(location)), :
## Error() model is singular
```

```
summary(aov.result2)
```

```
##
## Error: factor(method):factor(location)
##           Df Sum Sq Mean Sq F value Pr(>F)
## factor(method)  2  665.7   332.8   1.741   0.23
## Residuals      9 1720.7   191.2
##
## Error: Within
##           Df Sum Sq Mean Sq F value Pr(>F)
## Residuals 12  15.62    1.302
```

1.3.2 comparison with one-way

If we wrongly ignore the nested factor and use one-way ANOVA, the results would be different

Note that there's only one group factor, therefore, the type of summation does not matter.

```
aov_model = aov(DV ~ factor(method),
                data = nestedData, summation="III")
```

```
## Warning: In lm.fit(x, y, offset = offset, singular.ok = singular.ok, ...) :
## extra argument 'summation' will be disregarded
```

```
summary(aov_model)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## factor(method)  2  665.7   332.8    4.026 0.0331 *
## Residuals      21 1736.3    82.7
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
aov_model = aov(DV ~ factor(method),
                data = nestedData)
summary(aov_model)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## factor(method)  2  665.7   332.8    4.026 0.0331 *
## Residuals      21 1736.3    82.7
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

2 MANOVA

2.1 0

```
library("readxl")
library("rstatix")
```

```
##
## Attaching package: 'rstatix'

## The following object is masked from 'package:stats':
##
##   filter
```

```
library("MVTests")
```

```
##
## Attaching package: 'MVTests'

## The following object is masked from 'package:datasets':
##
##      iris
```

```
library("heplots")
```

```
## Loading required package: broom
```

```
# salary data example
salaryData = readxl::read_xlsx("example.xlsx")
DV = cbind(salaryData$salary, salaryData$salbegin)
jobcat = as.factor(salaryData$jobcat)
gender = as.factor(salaryData$gender)
```

2.2 visualization

Note that in order to generate the boxplot according to the “categorical” data, “factor” type should be designated in the mapping parameter.

If not “factored”, the scatter plot works fine with grouped effect, but the boxplot fails.

```
# eyeball the data
library(gridExtra)
```

```
##
## Attaching package: 'gridExtra'


## The following object is masked from 'package:dplyr':
##
##      combine

p1 = ggplot(salaryData, aes(x = factor(jobcat), y = salary)) +
  geom_boxplot(outlier.shape = NA) + geom_jitter(width = 0.2) +
```

```

theme(legend.position="top")
p2 = ggplot(salaryData, aes(x = factor(jobcat), y = salbegin)) +
  geom_boxplot(outlier.shape = NA) + geom_jitter(width = 0.2) +
  theme(legend.position="top")
grid.arrange(p1, p2, ncol=2)

```




We can also eyeball the DVs.

```

library(car)
scatterplot(salary ~ salbegin, data=salaryData, ellipse=TRUE,
  smooth=list(style="lines"))

```



2.3 assumptions

2.3.1 normality

2.3.1.1 pipe-friendly (Preferred) Preferred!

Note that `mshapiro_test()` is from the package `rstatix`, one step further from `tidyverse`.

```
salaryData |> select(salary,salbegin) |> mshapiro_test()
```

```
## # A tibble: 1 x 2
##   statistic p.value
##   <dbl>     <dbl>
## 1      0.707 6.13e-28
```

```
salaryData |> group_by(jobcat) |>
  shapiro_test(salary,salbegin)
```

```
## # A tibble: 6 x 4
##   jobcat variable statistic      p
```

```
##      <dbl> <chr>          <dbl>    <dbl>
## 1      1 salary          0.882 4.57e-16
## 2      1 salbegin       0.931 6.10e-12
## 3      2 salary          0.818 2.86e- 4
## 4      2 salbegin       0.499 1.64e- 8
## 5      3 salary          0.929 1.72e- 4
## 6      3 salbegin       0.860 2.14e- 7
```

2.3.1.2 classical (Not Preferred) Not preferred. It's not efficient to do this way.

The specific function `mshapiro.test()` requires to library the package `mvnrmtest`, which surprisingly has a single function! Moreover, it requires the data to be transposed before being processed.

The most annoying part is that the data has to be packed purposefully.

```
mvnrmtest::mshapiro.test(t(DV))
```

```
##
##  Shapiro-Wilk normality test
##
## data:  Z
## W = 0.70689, p-value < 2.2e-16
```

2.3.2 homogeneity of variance & covariance

Using the Box's M-test and Levene's Test.

2.3.2.1 pipe-friendly (Preferred) Using `box_m()` from the package `rstatix` to test the homogeneity of covariance matrices, in which the first parameter is the DVs needed to be analyzed (No group variable! This can be achieved through `select()`), and the second is the `group` parameter specifying the group variable.

Moreover, the function `heplots::boxM()` is helpful. Instead of taking the formula to specify the parameters, using the “traditional” way to write `group = ...` works pretty well combined with pipe-friendly environment!

The drawback of the functions is that they could not search column variables automatically, and that should be specified manually. And more annoying one is that, when encountering with NAs, an error will be raised by `rstatix::box_m()`, while that’s fine with `heplots::boxM()`.

```
salaryData |> select(salary,salbegin) |>
  box_m(group = jobcat)
```

```
## # A tibble: 1 x 4
##   statistic    p.value parameter method
##   <dbl>      <dbl>      <dbl> <chr>
## 1      509. 1.14e-106          6 Box's M-test for Homogeneity of Covariance Matr~
```

```
salaryData |> select(salary,salbegin) |>
  heplots::boxM(group = jobcat)
```

```
##
## Box's M-test for Homogeneity of Covariance Matrices
##
## data:  select(salaryData, salary, salbegin)
## Chi-Sq (approx.) = 508.68, df = 6, p-value < 2.2e-16
```

Do not forget that the assumption of homogeneity of variance for each DV is needed either! However, the traditional `levene_test()` (or its variants) can deal with one DV at a time, which is tedious when we have many DVs to check.

Hopefully, we have `heplots::leveneTests(data,group)` to deal with that issue. Combined with pipe-friendly procedures, that’s fabulous!

```
salaryData |> select(salbegin,salary) |>
  heplots::leveneTests(group = jobcat)
```

```
## Levene's Tests for Homogeneity of Variance (center = median)
```

```
##
##           df1 df2 F value    Pr(>F)
## salbegin   2 471  68.862 < 2.2e-16 ***
## salary     2 471  51.189 < 2.2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

2.3.2.2 classical The most annoying part is that the data has to be packed purposefully.

An alternative way to use `heplots::boxM()` has been mentioned above. We can renovate the efficiency to carry this test by replacing the “formula” specifying-pattern with the “group” one.

```
heplots::boxM(cbind(salary, salbegin) ~ factor(jobcat),
              data = salaryData)
```

```
##
## Box's M-test for Homogeneity of Covariance Matrices
##
## data: Y
## Chi-Sq (approx.) = 508.68, df = 6, p-value < 2.2e-16
```

2.4 MANOVA

Use the function `manova()`, which is from the `baseR`, no special need to library packages.

Similarly, the most annoying part is that the DVs and IVs has to be packed previously.

```
fit = manova(DV ~ jobcat)
```

As before, simply use `summary()` of the fitted model to see the results.

```
summary(fit) #for manova

##              Df  Pillai approx F num Df den Df    Pr(>F)
## jobcat        2 0.67298   119.43      4    942 < 2.2e-16 ***
## Residuals 471
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

2.5 effect size

Checking the effect size is also indispensable.

Use `eta_squared()` from the package `effectsize`, which contains plenty of useful functions computing effect size under various scenarios.

Note that there's also an `eta_squared()` from the package `rstatix`, but that can only deal with univariate ANOVA, it would fail when encountering MANOVA models.

Also note that the package `effectsize` is incorporated in `bruceR`!

```
effectsize::eta_squared(fit)

## # Effect Size for ANOVA (Type I)
##
## Parameter | Eta2 (partial) |          95% CI
## -----
## jobcat    |          0.34 | [0.30, 1.00]
##
## - One-sided CIs: upper bound fixed at [1.00].
```

2.6 univariate ANOVA

Needed if MANOVA reports significant results.

`summary.aov()` will report the univariate ANOVA respectively.

```
summary.aov(fit) #for indDVi dual anovas

## Response 1 :
##           Df      Sum Sq    Mean Sq F value    Pr(>F)
## jobcat      2 8.9438e+10 4.4719e+10  434.48 < 2.2e-16 ***
## Residuals  471 4.8478e+10 1.0293e+08
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Response 2 :
##           Df      Sum Sq    Mean Sq F value    Pr(>F)
## jobcat      2 1.7926e+10 8962772266  371.11 < 2.2e-16 ***
## Residuals  471 1.1375e+10  24151508
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

2.7 post-hoc tests

Note that the post-hoc tests here are essentially for the univariate ANOVA.

Note that the `games_howell_test()` is different from `tukey_hsd()`.

Note that for many DVs to take the post-hoc tests simultaneously, first `gather()` the targeted columns, and then take the **grouped** data for the test.

```
salaryData %>% gather(key = "var",
                      value = "value",
                      salary, salbegin) |>
  group_by(var) |>
  games_howell_test(value~jobcat)
```

```
# A tibble: 6 x 9
```

```
var      .y.   group1 group2 estimate conf.low conf.high    p.adj p.adj.signif
```

* <chr>	<chr>	<chr>	<chr>	<dbl>	<dbl>	<dbl>	<dbl>	<chr>
1 salary	value 1		2	3100.	1746.	4455.	1.22e- 6	****
2 salary	value 1		3	36139.	31302.	40977.	4.25e-10	****
3 salary	value 2		3	33039.	28196.	37881.	4.21e-10	****
4 salbegin	value 1		2	982.	256.	1707.	6 e- 3	**
5 salbegin	value 1		3	16162.	13539.	18784.	2.74e-10	****
6 salbegin	value 2		3	15180.	12514.	17846.	4.77e-10	****