

# Introduction to Optimization

## Lecture 13: Constrained problems and optimality conditions.



# From last class...

## Linear programming and optimality conditions

We have

$$(LP) \quad \min_{x \in \mathbb{R}^N} \{ c \cdot x : Ax + s = b, \text{ and } s \geq 0 \}.$$

$$(DLP) \quad \min_{y \in \mathbb{R}^M} \{ b \cdot y : A^T y + c = 0, \text{ and } y \geq 0 \}.$$

### Primal-dual optimality conditions

**Primal feasibility:**  $Ax + s = b$  and  $s \geq 0$ .

**Dual feasibility:**  $A^T y + c = 0$  and  $y \geq 0$ .

**Complementarity:**  $y_i s_i = 0$  for  $i = 1, \dots, M$ .

# Idea of the proof

Write  $K = N + M$  and  $z = (x, s)^T$ . Define  $F : \mathbb{R}^K \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $G : \mathbb{R}^M \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $P \in \mathbb{R}^{K \times M}$  by

$$F(z) = c \cdot x + \iota_{\mathbb{R}_+^M}(s), \quad G(y) = \iota_{\{b\}}(y) \quad \text{and} \quad Pz = Ax + s,$$

respectively, so that  $(LP)$  consists in minimizing  $F(z) + G(Pz)$ .

# Idea of the proof

Write  $K = N + M$  and  $z = (x, s)^T$ . Define  $F : \mathbb{R}^K \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $G : \mathbb{R}^M \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $P \in \mathbb{R}^{K \times M}$  by

$$F(z) = c \cdot x + \iota_{\mathbb{R}_+^M}(s), \quad G(y) = \iota_{\{b\}}(y) \quad \text{and} \quad Pz = Ax + s,$$

respectively, so that  $(LP)$  consists in minimizing  $F(z) + G(Pz)$ . We compute

$$\partial F(z) = \begin{bmatrix} c \\ \partial \iota_{\mathbb{R}_+^M}(s) \end{bmatrix}$$

# Idea of the proof

Write  $K = N + M$  and  $z = (x, s)^T$ . Define  $F : \mathbb{R}^K \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $G : \mathbb{R}^M \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $P \in \mathbb{R}^{K \times M}$  by

$$F(z) = c \cdot x + \iota_{\mathbb{R}_+^M}(s), \quad G(y) = \iota_{\{b\}}(y) \quad \text{and} \quad Pz = Ax + s,$$

respectively, so that  $(LP)$  consists in minimizing  $F(z) + G(Pz)$ . We compute

$$\partial F(z) = \begin{bmatrix} c \\ \partial \iota_{\mathbb{R}_+^M}(s) \end{bmatrix}, \quad \partial G(y) = \begin{cases} \emptyset & \text{if } y \neq b \\ \mathbb{R}^M & \text{if } y = b \end{cases}$$

# Idea of the proof

Write  $K = N + M$  and  $z = (x, s)^T$ . Define  $F : \mathbb{R}^K \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $G : \mathbb{R}^M \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $P \in \mathbb{R}^{K \times M}$  by

$$F(z) = c \cdot x + \iota_{\mathbb{R}_+^M}(s), \quad G(y) = \iota_{\{b\}}(y) \quad \text{and} \quad Pz = Ax + s,$$

respectively, so that  $(LP)$  consists in minimizing  $F(z) + G(Pz)$ . We compute

$$\partial F(z) = \begin{bmatrix} c \\ \partial \iota_{\mathbb{R}_+^M}(s) \end{bmatrix}, \quad \partial G(y) = \begin{cases} \emptyset & \text{if } y \neq b \\ \mathbb{R}^M & \text{if } y = b \end{cases}$$

and

$$P^T y = \begin{bmatrix} A^T y \\ y \end{bmatrix}.$$

## Idea of the proof, continued

The primal-dual optimality conditions are

$$-P^T \hat{y} \in \partial F(\hat{z}) \quad \text{and} \quad \hat{y} \in \partial G(P\hat{x}),$$

which we translate as

$$\begin{aligned} -P^T \hat{y} \in \partial F(\hat{x}) &\Leftrightarrow -A^T \hat{y} = c, \quad \hat{s} \geq 0 \quad \text{and} \quad -\hat{y} \in \partial \iota_{\mathbb{R}_+^M}(\hat{s}). \\ \hat{y} \in \partial G(P\hat{x}) &\Leftrightarrow A\hat{x} + \hat{s} = b, \quad \hat{y} \in \mathbb{R}^M. \end{aligned}$$

## Idea of the proof, continued

The primal-dual optimality conditions are

$$-P^T \hat{y} \in \partial F(\hat{z}) \quad \text{and} \quad \hat{y} \in \partial G(P\hat{x}),$$

which we translate as


$$\begin{aligned} -P^T \hat{y} \in \partial F(\hat{x}) &\Leftrightarrow -A^T \hat{y} = c, \quad \hat{s} \geq 0 \quad \text{and} \quad -\hat{y} \in \partial \iota_{\mathbb{R}_+^M}(\hat{s}). \\ \hat{y} \in \partial G(P\hat{x}) &\Leftrightarrow A\hat{x} + \hat{s} = b, \quad \hat{y} \in \mathbb{R}^M. \end{aligned}$$

On the other hand,  $\hat{y} \in \partial \iota_{\mathbb{R}_+^M}(\hat{s})$  means that, for  $i = 1, \dots, M$ ,  $\hat{y}_i = 0$  if  $\hat{s}_i > 0$ , and  $\hat{y}_i \geq 0$  if  $\hat{s}_i = 0$ .




# Example I

Distance from a plane to the origin



# Example I

Distance from a plane to the origin




- Minimize distance to the origin

$$\sqrt{x^2 + y^2 + z^2}.$$

# Example I

Distance from a plane to the origin



- Minimize distance to the origin


$$\sqrt{x^2 + y^2 + z^2}.$$

Equivalently, minimize

$$x^2 + y^2 + z^2.$$

# Example I

Distance from a plane to the origin



- Minimize distance to the origin

$$\sqrt{x^2 + y^2 + z^2}.$$

Equivalently, minimize

$$x^2 + y^2 + z^2.$$

- Subject to the constraint

$$ax + by + cz - d = 0.$$

# Optimization problems with equality constraints

Let  $f, h_1, h_2, \dots, h_M \in \mathcal{C}^1(\mathbb{R}^N; \mathbb{R})$ , and consider the **constrained problem**

$$(\mathcal{P}) \quad \min_{x \in C} f(x),$$

where the **feasible set**  $C$  is given by

$$C = \{x \in \mathbb{R}^N : h_m(x) = 0, \quad m = 1, \dots, M\}.$$

# Optimization problems with equality constraints

Let  $f, h_1, h_2, \dots, h_M \in \mathcal{C}^1(\mathbb{R}^N; \mathbb{R})$ , and consider the **constrained problem**

$$(\mathcal{P}) \quad \min_{x \in C} f(x),$$

where the **feasible set**  $C$  is given by

$$C = \{x \in \mathbb{R}^N : h_m(x) = 0, \quad m = 1, \dots, M\}.$$

A local solution (constrained minimizer)  $\hat{x}$  is **regular** if the set

$$V = \{\nabla h_1(\hat{x}), \nabla h_2(\hat{x}), \dots, \nabla h_M(\hat{x})\}$$

is linearly independent.

# Necessary conditions for optimality

## Theorem (Lagrange Multiplier Theorem)

Let  $\hat{x}$  be a regular local minimizer of  $(\mathcal{P})$ . There is a unique  $\hat{\lambda} \in \mathbb{R}^M$ , called *Lagrange multiplier vector*, such that

$$\nabla f(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla h_m(\hat{x}) = 0.$$

# Necessary conditions for optimality

## Theorem (Lagrange Multiplier Theorem)

Let  $\hat{x}$  be a regular local minimizer of  $(\mathcal{P})$ . There is a unique  $\hat{\lambda} \in \mathbb{R}^M$ , called **Lagrange multiplier vector**, such that

$$\nabla f(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla h_m(\hat{x}) = 0.$$

If, in addition,  $f, h_1, h_2, \dots, h_M \in \mathcal{C}^2(\mathbb{R}^N; \mathbb{R})$ , then

$$y \cdot \left( \nabla^2 f(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla^2 h_m(\hat{x}) \right) y \geq 0$$

for all  $y \in V^\perp$ .



# Sketch of the proof

- Pick  $x_k$  that minimizes

$$f_k(x) = f(x) + \frac{\varepsilon}{2} \|x - \hat{x}\|^2 + \frac{k}{2} \sum_{m=1}^M \|h_m(x)\|^2$$

over a small ball around  $\hat{x}$ .

# Sketch of the proof

- Pick  $x_k$  that minimizes

$$f_k(x) = f(x) + \frac{\varepsilon}{2} \|x - \hat{x}\|^2 + \frac{k}{2} \sum_{m=1}^M \|h_m(x)\|^2$$

over a small ball around  $\hat{x}$ .

- Let  $k \rightarrow \infty$  and use the optimality condition  $\nabla f_k(x_k) = 0$  to deduce that  $x_k \rightarrow \hat{x}$  and  $kh_m(x_k) \rightarrow s_m$ .

# Sketch of the proof

- Pick  $x_k$  that minimizes

$$f_k(x) = f(x) + \frac{\varepsilon}{2} \|x - \hat{x}\|^2 + \frac{k}{2} \sum_{m=1}^M \|h_m(x)\|^2$$

over a small ball around  $\hat{x}$ .

- Let  $k \rightarrow \infty$  and use the optimality condition  $\nabla f_k(x_k) = 0$  to deduce that  $x_k \rightarrow \hat{x}$  and  $kh_m(x_k) \rightarrow s_m$ .
- Show that  $\hat{\lambda}_m = s_m$  has the desired property.

# Sketch of the proof

- Pick  $x_k$  that minimizes


$$f_k(x) = f(x) + \frac{\varepsilon}{2} \|x - \hat{x}\|^2 + \frac{k}{2} \sum_{m=1}^M \|h_m(x)\|^2$$

over a small ball around  $\hat{x}$ .

- Let  $k \rightarrow \infty$  and use the optimality condition  $\nabla f_k(x_k) = 0$  to deduce that  $x_k \rightarrow \hat{x}$  and  $kh_m(x_k) \rightarrow s_m$ .
- Show that  $\hat{\lambda}_m = s_m$  has the desired property.
- Proceed analogously to obtain the second order condition.


# Back to Example I

Find the point in the plane  $ax + by + cz = d$  that is closest to the origin



# Back to Example I

Find the point in the plane  $ax + by + cz = d$  that is closest to the origin




- Minimize distance to the origin

$$\sqrt{x^2 + y^2 + z^2}.$$

# Back to Example I

Find the point in the plane  $ax + by + cz = d$  that is closest to the origin



- Minimize distance to the origin


$$\sqrt{x^2 + y^2 + z^2}.$$

Equivalently, minimize

$$x^2 + y^2 + z^2.$$

# Back to Example I

Find the point in the plane  $ax + by + cz = d$  that is closest to the origin



- Minimize distance to the origin

$$\sqrt{x^2 + y^2 + z^2}.$$

Equivalently, minimize

$$x^2 + y^2 + z^2.$$

- Subject to the constraint

$$ax + by + cz - d = 0.$$



# Second order sufficient conditions

## Theorem

Let  $f, h_1, h_2, \dots, h_M : \mathbb{R}^N \rightarrow \mathbb{R}$  be twice continuously differentiable, and let  $\hat{x} \in \mathcal{C}$  and  $\hat{\lambda} \in \mathbb{R}^M$  satisfy


$$\nabla f(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla h_m(\hat{x}) = 0$$

and

$$y \cdot \left( \nabla^2 f(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla^2 h_m(\hat{x}) \right) y > 0$$

for all  $y \in V^\perp \setminus \{0\}$ . Then,  $\hat{x}$  is a strict local minimizer of  $(\mathcal{P})$ .

# Break




# Example II

## Hydraulic Cylinder Design



# Example II


## Hydraulic Cylinder Design – Manufacturing Constraints



- Minimal force required
  - $f = \pi r^2 p \geq f_{min}$ .

# Example II

## Hydraulic Cylinder Design – Manufacturing Constraints



- Minimal force required


- $f = \pi r^2 p \geq f_{min}$ .

- Maximal hoop stress

- $s = \frac{pr}{t} \leq s_{max}$

# Example II


## Hydraulic Cylinder Design – Manufacturing Constraints



- Minimal force required
  - $f = \pi r^2 p \geq f_{min}$ .
- Maximal hoop stress
  - $s = \frac{pr}{t} \leq s_{max}$
- Minimal thickness
  - $t \geq t_{min}$

## Example II


### Hydraulic Cylinder Design – Manufacturing Constraints



- Minimal force required
  - $f = \pi r^2 p \geq f_{min}$ .
- Maximal hoop stress
  - $s = \frac{pr}{t} \leq s_{max}$
- Minimal thickness
  - $t \geq t_{min}$
- Maximal pressure
  - $p \leq p_{max}$ .

# Example II

## Hydraulic Cylinder Design – Optimization Problem



- Minimize the total width  $2(r + t)$
- Subject to the constraints
  - $pr - ts_{max} \leq 0$
  - $f_{min} - \pi r^2 p \leq 0$
  - $t_{min} - t \leq 0$
  - $p - p_{max} \leq 0$ .



# Inequality constraints

Let  $f, g_j, h_m \in \mathcal{C}^1(\mathbb{R}^N; \mathbb{R})$ , and consider the constrained problem

$$(\mathcal{P}) \quad \min_{x \in C} f(x),$$

where the feasible set  $C$  is now given by

$$C = \{x \in \mathbb{R}^N : h_m(x) = 0, g_j(x) \leq 0, \forall m, j\}.$$

# Inequality constraints

Let  $f, g_j, h_m \in \mathcal{C}^1(\mathbb{R}^N; \mathbb{R})$ , and consider the constrained problem

$$(\mathcal{P}) \quad \min_{x \in C} f(x),$$

where the feasible set  $C$  is now given by


$$C = \{x \in \mathbb{R}^N : h_m(x) = 0, g_j(x) \leq 0, \forall m, j\}.$$

The set of **active inequality constraints** at a local solution  $\hat{x}$  is

$$A(\hat{x}) = \{j : g_j(\hat{x}) = 0\}.$$

# Back to Example II

## Hydraulic Cylinder Design – Optimization Problem



- Minimize the total width  
 $2(r + t)$
- Subject to the constraints
  - $pr - ts_{max} \leq 0$
  - $f_{min} - \pi r^2 p \leq 0$
  - $t_{min} - t \leq 0$
  - $p - p_{max} \leq 0$ .

# Necessary conditions for optimality

## Theorem (Karush-Kuhn-Tucker conditions)

Let  $\hat{x}$  be a regular local minimizer of  $(\mathcal{P})$ . There exist unique Lagrange multiplier vectors  $\hat{\lambda} \in \mathbb{R}^M$  and  $\hat{\mu} \in \mathbb{R}_+^J$ , such that

$$\nabla f(\hat{x}) + \sum_{j=1}^J \hat{\mu}_j \nabla g_j(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla h_m(\hat{x}) = 0,$$

and  $\hat{\mu}_j = 0$  for all  $j \notin A(\hat{x})$ , which means that  $\hat{\mu}_j g_j(\hat{x}) = 0$  for all  $j$ .

# Necessary conditions for optimality

## Theorem (Karush-Kuhn-Tucker conditions)

Let  $\hat{x}$  be a regular local minimizer of  $(\mathcal{P})$ . There exist unique Lagrange multiplier vectors  $\hat{\lambda} \in \mathbb{R}^M$  and  $\hat{\mu} \in \mathbb{R}_+^J$ , such that

$$\nabla f(\hat{x}) + \sum_{j=1}^J \hat{\mu}_j \nabla g_j(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla h_m(\hat{x}) = 0,$$

and  $\hat{\mu}_j = 0$  for all  $j \notin A(\hat{x})$ , which means that  $\hat{\mu}_j g_j(\hat{x}) = 0$  for all  $j$ .

If, in addition,  $f, g_j, h_m \in \mathcal{C}^2(\mathbb{R}^N; \mathbb{R})$ , then

$$y \cdot \left( \nabla^2 f(\hat{x}) + \sum_{m=1}^M \hat{\lambda}_m \nabla^2 h_m(\hat{x}) \right) y \geq 0$$

for all  $y \in \mathcal{V}^\perp$ , where  $\mathcal{V} = \{ \nabla g_j(\hat{x}), \nabla h_m(\hat{x}), j \in A(\hat{x}), m = 1, \dots, M \}$ .