

Introduction to Optimization

Lecture 11: Convergence of the proximal-gradient algorithm. Conjugate functions.



university of
 groningen

Proximal-gradient algorithm

Suppose we want to find the minima of $f = g + h$, where $g : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is closed and convex, and $h : \mathbb{R}^N \rightarrow \mathbb{R}$ is L -smooth and convex.

Proximal-gradient algorithm

Suppose we want to find the minima of $f = g + h$, where $g : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is closed and convex, and $h : \mathbb{R}^N \rightarrow \mathbb{R}$ is L -smooth and convex.

Example

A typical example in image and signal processing, statistics, ML, is

$$f(x) = \frac{1}{2} \|Ax - b\|^2 + \rho \|x\|_1$$

for $x \in \mathbb{R}^N$.

Proximal-gradient algorithm

The **proximal-gradient** method consists in applying proximal iterations while linearizing the smooth function:

$$x_{n+1} = \text{prox}_{\alpha g}(x_n - \alpha \nabla h(x_n))$$

Proximal-gradient algorithm

The **proximal-gradient** method consists in applying proximal iterations while linearizing the smooth function:

$$\begin{aligned}x_{n+1} &= \text{prox}_{\alpha g}(x_n - \alpha \nabla h(x_n)) \\ &= \text{Argmin} \left\{ g(x) + \frac{1}{2\alpha} \|x - (x_n - \alpha \nabla h(x_n))\|^2 \right\}.\end{aligned}$$

Proximal-gradient algorithm

The **proximal-gradient** method consists in applying proximal iterations while linearizing the smooth function:

$$\begin{aligned}x_{n+1} &= \text{prox}_{\alpha g}(x_n - \alpha \nabla h(x_n)) \\ &= \text{Argmin} \left\{ g(x) + \frac{1}{2\alpha} \|x - (x_n - \alpha \nabla h(x_n))\|^2 \right\}.\end{aligned}$$

This subproblem has a unique solution characterized by

$$0 \in \partial g(x_{n+1}) + \nabla h(x_n) + \frac{1}{\alpha}(x_{n+1} - x_n).$$

Convergence of proximal-gradient sequences

Theorem

Let $f = g + h$, where $g : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is closed and convex, and $h : \mathbb{R}^N \rightarrow \mathbb{R}$ is L -smooth and convex. Take $\alpha \in (0, 1/L]$ and define (x_n) by

$$x_{n+1} = \text{prox}_{\alpha g}(x_n - \alpha \nabla h(x_n)), \quad n \geq 0.$$

If $S \neq \emptyset$, x_n converges to an $\hat{x} \in S$, and

$$f(x_n) - \min(f) \leq \frac{\text{dist}(x_0, S)^2}{2\alpha n}, \quad n \geq 1.$$

Moreover, $\lim_{n \rightarrow \infty} n(f(x_n) - \min(f)) = 0$.

Sketch of the proof

$f(x_n) + g(x_n)$ is nonincreasing

$$\begin{cases} f(x_{n+1}) &\leq f(x_n) + \nabla f(x_n) \cdot (x_{n+1} - x_n) + \frac{L}{2} \|x_{n+1} - x_n\|^2 \\ g(x_{n+1}) &\leq g(x_n) + \left(\frac{x_n - x_{n+1}}{\alpha} - \nabla f(x_n) \right) \cdot (x_{n+1} - x_n). \end{cases}$$

Sketch of the proof

$f(x_n) + g(x_n)$ is nonincreasing

$$\begin{cases} f(x_{n+1}) &\leq f(x_n) + \nabla f(x_n) \cdot (x_{n+1} - x_n) + \frac{L}{2} \|x_{n+1} - x_n\|^2 \\ g(x_{n+1}) &\leq g(x_n) + \left(\frac{x_n - x_{n+1}}{\alpha} - \nabla f(x_n) \right) \cdot (x_{n+1} - x_n). \end{cases}$$

Compatibility

$$\|x_{n+1} - x_n - \alpha(\nabla f(x_{n+1}) - \nabla f(x_n))\|^2 \leq \|x_{n+1} - x_n\|^2.$$

Sketch of the proof

$f(x_n) + g(x_n)$ is nonincreasing

$$\begin{cases} f(x_{n+1}) &\leq f(x_n) + \nabla f(x_n) \cdot (x_{n+1} - x_n) + \frac{L}{2} \|x_{n+1} - x_n\|^2 \\ g(x_{n+1}) &\leq g(x_n) + \left(\frac{x_n - x_{n+1}}{\alpha} - \nabla f(x_n) \right) \cdot (x_{n+1} - x_n). \end{cases}$$

Compatibility

$$\|x_{n+1} - x_n - \alpha(\nabla f(x_{n+1}) - \nabla f(x_n))\|^2 \leq \|x_{n+1} - x_n\|^2.$$

Convergence rate

$$\begin{cases} f(x_n) &\leq f(p) + \nabla f(x_n) \cdot (x_n - p) \\ g(x_{n+1}) &\leq g(p) + \left(\frac{x_n - x_{n+1}}{\alpha} - \nabla f(x_n) \right) \cdot (x_{n+1} - p). \end{cases}$$

Break

The Fenchel conjugate

The **Fenchel conjugate** of a closed convex function $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is the function $f^* : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$f^*(x^*) = \sup_{x \in \mathbb{R}^N} \{x^* \cdot x - f(x)\}.$$

The Fenchel conjugate

The **Fenchel conjugate** of a closed convex function $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is the function $f^* : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$f^*(x^*) = \sup_{x \in \mathbb{R}^N} \{x^* \cdot x - f(x)\}.$$

Examples

- In \mathbb{R} : (1) $f(x) = x$;

The Fenchel conjugate

The **Fenchel conjugate** of a closed convex function $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is the function $f^* : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$f^*(x^*) = \sup_{x \in \mathbb{R}^N} \{x^* \cdot x - f(x)\}.$$

Examples

- In \mathbb{R} : (1) $f(x) = x$; (2) $f(x) = \frac{1}{2}x^2$;

The Fenchel conjugate

The **Fenchel conjugate** of a closed convex function $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is the function $f^* : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$f^*(x^*) = \sup_{x \in \mathbb{R}^N} \{x^* \cdot x - f(x)\}.$$

Examples

- In \mathbb{R} : (1) $f(x) = x$; (2) $f(x) = \frac{1}{2}x^2$; (3) $f(x) = e^x$.

The Fenchel conjugate

The **Fenchel conjugate** of a closed convex function $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is the function $f^* : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$f^*(x^*) = \sup_{x \in \mathbb{R}^N} \{x^* \cdot x - f(x)\}.$$

Examples

- In \mathbb{R} : (1) $f(x) = x$; (2) $f(x) = \frac{1}{2}x^2$; (3) $f(x) = e^x$.
- In \mathbb{R}^N : (4) $f(x) = c \cdot x + \alpha$;

The Fenchel conjugate

The **Fenchel conjugate** of a closed convex function $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is the function $f^* : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$f^*(x^*) = \sup_{x \in \mathbb{R}^N} \{x^* \cdot x - f(x)\}.$$

Examples

- In \mathbb{R} : (1) $f(x) = x$; (2) $f(x) = \frac{1}{2}x^2$; (3) $f(x) = e^x$.
- In \mathbb{R}^N : (4) $f(x) = c \cdot x + \alpha$; (5) $f(x) = \phi(\|x\|)$.

Basic properties

Proposition

- 1 The function f^* is closed, convex and not identically $+\infty$.

Basic properties

Proposition

- 1 The function f^* is closed, convex and not identically $+\infty$.
- 2 If $f \leq g$, then $f^* \geq g^*$.

Basic properties

Proposition

- 1 The function f^* is closed, convex and not identically $+\infty$.
- 2 If $f \leq g$, then $f^* \geq g^*$.
- 3 *Fenchel-Young Inequality*: $f(x) + f^*(x^*) \geq \langle x^*, x \rangle$.

Basic properties

Proposition

- ① The function f^* is closed, convex and not identically $+\infty$.
- ② If $f \leq g$, then $f^* \geq g^*$.
- ③ **Fenchel-Young Inequality:** $f(x) + f^*(x^*) \geq \langle x^*, x \rangle$.
There is equality if, and only if, $x^* \in \partial f(x)$.

Basic properties

Proposition

- ① The function f^* is closed, convex and not identically $+\infty$.
- ② If $f \leq g$, then $f^* \geq g^*$.
- ③ **Fenchel-Young Inequality:** $f(x) + f^*(x^*) \geq \langle x^*, x \rangle$.
There is equality if, and only if, $x^* \in \partial f(x)$.
- ④ $f^{**} = f$.

Basic properties

Proposition

- ① The function f^* is closed, convex and not identically $+\infty$.
- ② If $f \leq g$, then $f^* \geq g^*$.
- ③ **Fenchel-Young Inequality:** $f(x) + f^*(x^*) \geq \langle x^*, x \rangle$.
There is equality if, and only if, $x^* \in \partial f(x)$.
- ④ $f^{**} = f$.
- ⑤ **Legendre-Fenchel Reciprocity Formula:** $x^* \in \partial f(x)$ if, and only if, $x \in \partial f^*(x^*)$.

Basic properties

Proposition

- ❶ The function f^* is closed, convex and not identically $+\infty$.
- ❷ If $f \leq g$, then $f^* \geq g^*$.
- ❸ **Fenchel-Young Inequality:** $f(x) + f^*(x^*) \geq \langle x^*, x \rangle$.
There is equality if, and only if, $x^* \in \partial f(x)$.
- ❹ $f^{**} = f$.
- ❺ **Legendre-Fenchel Reciprocity Formula:** $x^* \in \partial f(x)$ if, and only if, $x \in \partial f^*(x^*)$.
- ❻ Let $\mu\ell = 1$. Then, f is μ -strongly convex if, and only if, f^* is ℓ -smooth.