



Research Article

Algae 2022, 37(4): 265-279
<https://doi.org/10.4490/algae.2022.37.11.28>

Open Access



Primary production by phytoplankton in the territorial seas of the Republic of Korea

An Suk Lim¹ and Hae Jin Jeong^{2,3,*}

¹Division of Life Science, Gyeongsang National University, Jinju 52828, Korea

²School of Earth and Environmental Sciences, College of Natural Sciences, Seoul National University, Seoul 08826, Korea

³Research Institute of Oceanography, Seoul National University, Seoul 08826, Korea

The primary production (PP) by phytoplankton in marine ecosystems is essential for carbon cycling and fueling food webs. Hence, estimating the PP in the territorial sea of each country is a necessary step to achieving carbon neutrality. To estimate the PP in the territorial sea of the Republic of Korea from 2005 to 2021, we analyzed various physiochemical parameters, such as sea surface temperature (SST), Secchi depth, and concentrations of chlorophyll-*a* and nutrients in the seas of five regions, including the East Sea, West Sea, western South Sea, eastern South Sea, and the waters off Jeju Island. During the 17-year study period, the SST tended to increase, while the nutrient concentrations declined, except in the Jeju area. Overall, the PP did not show a specific temporal trend, but daily PP in the western South Sea was the highest among the five regions. Moreover, the maximum PP in the Korean territorial waters ($76,450 \text{ km}^2$) was estimated at $11,227 \text{ Gg C y}^{-1}$, which accounts for 0.03% of the global PP. The results may give insights into a better understanding of the PP, further resource utilization, and environmental sustainability in the studied region.

Keywords: carbon cycle; light utilization; photosynthetic assimilation; phytoplankton; primary productivity

INTRODUCTION

Oceans are one of the largest storages of carbon dioxide (CO_2) on the planet and play a major role in the global carbon cycle by absorbing approximately 30% of atmospheric CO_2 each year (Raven and Falkowski 1999). Although both pelagic and benthic communities of primary producers are responsible for primary production (PP) in the ocean, approximately half of the global annual net PP is produced by marine phytoplankton (Field et al. 1998, Spector and Edward 2020, Luo et al. 2021). This primary source of energy supports the entire marine biomass through food webs (Turner 2004, Eom et al. 2021, Jeong et al. 2021, Lee et al. 2021, Ok et al. 2021). Thus, it is crucial to accurately assess the geographical and temporal vari-

ability in phytoplankton production to better understand the global carbon dynamics and trophic energy flow.

Coastal waters support a high PP of approximately 10–30% of the global ocean production (Muller-Karger et al. 2005, Salgado-Hernanz et al. 2022). Moreover, physical and biological processes in coastal waters play an important role in carbon transport to deep oceans (Cai 2011). Therefore, estimating and comprehending PP in coastal waters is crucial for improving our understanding of the marine carbon cycle.

However, PP by marine phytoplankton is influenced by various physiochemical factors that are directly related to phytoplankton physiology, such as light, turbidity, water

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received October 10, 2022, Accepted November 28, 2022

*Corresponding Author

E-mail: hjeong@snu.ac.kr

Tel: +82-2-880-6746, Fax: +82-2-874-9695

temperature, salinity, and nutrients (Valiela 1984, Cloern 1987). Due to the difficulty in acquiring all of these variables, the estimation of marine PP is challenging. Several methods, including the measurement of oxygen production using light and dark bottle incubation (Nielsen 1975), have been used to determine the PP. Moreover, photosynthetic carbon assimilation in water samples can be measured using radioactive or stable isotopes (e.g., ^{14}C and ^{13}C) (Yoo and Shin 1995, Park 1996, Kwak et al. 2013a, 2013b). More recently, active fluorescence techniques, such as pulse-amplitude-modulated fluorometry and fast repetition rate fluorometry, have been applied to estimate photosynthetic parameters and / or PP (Falkowski et al. 2004, Smyth et al. 2004, Ko et al. 2019). As an indirect method, some algorithms developed using satellite images have been used to provide a global estimation of PP (Eppley et al. 1985, Yoon et al. 2012).

Over the last 10 years, long-term monitoring programs have been developed, and many scientists have monitored various environmental parameters in a bid to evaluate the quality of coastal waters. Therefore, in the present study, the PP in the Korean coastal waters was estimated using environmental parameters such as light intensity, chlorophyll-*a* (Chl-*a*) concentration, and Secchi depth (SeD) from 2005 to 2021. The photosynthetic assimilation rate and light utilization efficiency were obtained from the literature for each season to calculate the primary productivity of marine phytoplankton. The temporal PP in each area was then estimated by combining environmental and physiological parameters. The results have implications for understanding PP trends and coastal water management.

MATERIALS AND METHODS

The dataset

Data were obtained from the National Fisheries Research and Development Institute (NFRDI, https://www.nifs.go.kr/femo/data_obs.femo) and the Korea Marine Environment Management Cooperation (KOEM, <https://www.meis.go.kr/mei/observe/port.do>). A total of 376 sampling stations (231 from NFRDI and 145 from KOEM) were located near or along the coastline of the Korean Peninsula (Fig. 1). The dataset of each station included sea surface temperature (SST), SeD, dissolved inorganic nitrogen (DIN), dissolved inorganic phosphorus (DIP), and Chl-*a* concentrations in the surface and bottom waters. The concentrations (μM) of DIN and DIP were con-




Fig. 1. A map showing the study area. The bold solid line indicates the terrestrial waters and, the area of the terrestrial waters was divided into five regions where the dotted lines were indicated. The area of each region indicated the numbers under each region's name.

verted from those in $\mu\text{g L}^{-1}$ with conversion factors of 14 for DIN and 31 for DIP. Data from sites with a salinity of less than 30 were excluded because these sampling sites had extremely high nutrient concentrations (e.g., $>200 \mu\text{M}$ DIN) and were highly likely to have been influenced by artificial effects. The total number of data points was 21,540 from 2005 to 2021 (2011–2019 from NFRDI and 2005–2021 from KOEM). The data included samples from every other month (NFRDI) or from February, May, August, and November (KOEM) (Supplementary Table S1). Only data on both Chl-*a* and SeD that were available simultaneously were included in the PP calculation. The area of the territorial waters of Korea was obtained from the Korea Hydrographic and Oceanographic Agency (KHOA, <https://www meis.go.kr/map/oemsBaseMap.do>) (Fig. 1).

Estimation of PP

The daily PP was calculated from the photosynthesis-irradiance parameters, calculated depth profiles of photosynthetic active radiation (PAR), and the Chl-*a* concentration. To estimate the PP of phytoplankton in a water column for a day, the hourly primary productivity at each water depth was integrated from the sea surface (Z_0) to the euphotic depth (Z_{eu}) for the day length. It was assumed that no photoinhibition occurred. Therefore,

$$\text{PP}_{\text{day}} = \int_{Z_0}^{Z_{eu}} B \times P_m^B [1 - \exp(-\alpha^B \times I_z/P_m^B)] \times D_{\text{irr}}$$

, where PP_{day} is depth-integrated PP ($\text{mg C m}^{-2} \text{d}^{-1}$), B is the

Chl-*a* concentration (mg Chl-*a* m⁻³) at each depth, P_m^B is the maximum Chl-*a* normalized photosynthetic rate (mg C mg Chl-*a*⁻¹ h⁻¹), α^B is the photosynthetic efficiency (mg C mg Chl-*a*⁻¹ h⁻¹ [μmol photons m⁻² s⁻¹]⁻¹), I_z is the irradiance (μmol photons m⁻² s⁻¹) at each depth, and D_{irr} is the day length (h).

As a physiological response of phytoplankton to light, the maximum Chl-*a* normalized photosynthetic rate (P_m^B), known as the assimilation number or the activity coefficient of chlorophyll, is the maximum primary productivity under optimal light conditions. Photosynthetic efficiency (α^B) is controlled by the counteracting effects of light limitation and depends on the capability of the cell to gather incident light (Côté and Platt 1983). The photosynthetic parameters P_m^B and α^B were obtained from the literature, which measured each parameter using a short-term ¹⁴C experiment within the Korean coastal waters. The mean values for each parameter were categorized according to season and region (Table 1, Supplementary Table S2). The Chl-*a* concentration at each depth of each sampling station was calculated using the linear equations that were obtained by plotting the concentrations of Chl-*a* at the surface and bottom against the two depths. We considered the bottom depth to be 1 m above the reported water depth of each sampling station. It is possible that Chl-*a* exhibits a non-linear decrease from the surface to the bottom or a subsurface chlorophyll maximum layer exists. However, data on the vertical profiles of the Chl-*a* are scarce. Therefore, it was assumed that the concentration of Chl-*a* decreased linearly from the surface to the bottom waters.

The irradiance (μmol photons m⁻² s⁻¹) was converted from the total solar radiation (MJ m⁻² d⁻¹) obtained from the Korea Meteorological Administration (KMA). The monthly average values were applied for the conversion. In the present study, irradiance is considered at wave-

lengths between 400 and 700 nm (PAR, I), which consists of 44% of solar radiation (Moon 1940, Morel and Smith 1974, Shim 2003) (Supplementary Table S1). The irradiance at a given depth in the water column (I_z) was estimated using the following equation:

$$I_z = I_0 \times e^{-kz}$$

, where I_z is the irradiance at each depth (μmol photons m⁻² s⁻¹), I_0 is the irradiance at the surface (μmol photons m⁻² s⁻¹), k is the attenuation coefficient, and z is the depth (m). To estimate k , SeD was used, and k was converted using the relationship $k = 1.44 / \text{SeD}$ (Holmes 1970). The euphotic depth (Z_{eu}) was calculated as the depth at which PAR is 1% of the surface PAR and is given by $4.6 / k$ (Morel and Berthon 1989). Given the relationship with SeD, Z_{eu} was then given by $Z_{eu} = 3.2 \times \text{SeD}$. When the water column depth at the station was shallower than the calculated value of Z_{eu} , the water depth was considered as the euphotic zone depth. The average day length (h) in the sampling month was obtained from the KMA and applied to the equation to calculate the PP_{day}.

Statistical analyses

The photosynthetic parameters in each season in each group of regions were analyzed using a one-way analysis of variance (ANOVA). Prior to analysis, the normality and homogeneity of the data were tested using Kolmogorov-Smirnov and Levene's median tests, respectively. If the data did not satisfy the assumption of homogeneity, Welch's one-way ANOVA was performed. If the data did not fit the Kolmogorov-Smirnov and Levene's tests, the Kruskal-Wallis H test was conducted. Moreover, the photosynthetic parameters between the two groups of regions were analyzed using the t-test or Mann-Whitney U

Table 1. The photosynthetic parameters used in this study

Region	Season	P_m^B	α^B	Reference
East Sea, eastern South Sea and waters off Jeju island	Winter	1.42 (1.44)	0.020 (0.016)	Park (1996), Yoshikawa and Furuya (2008),
	Spring	3.11 (2.17)	0.051 (0.039)	Ko (2014), Kwon et al. (2015)
	Summer	3.91 (3.13)	0.025 (0.016)	
	Autumn	5.73 (3.94)	0.034 (0.018)	
West Sea and western South Sea	Winter	4.34 (2.50)	0.155 (0.200)	Kang et al. (1992), Yoo and Shin (1995), Chung (1996), Son et al. (2005), Park and Kim (2010), Ko (2014)
	Spring	7.50 (5.49)	0.141 (0.140)	
	Summer	5.84 (2.85)	0.054 (0.035)	
	Autumn	7.34 (6.41)	0.168 (0.247)	

P_m^B , maximum Chl-*a* normalized photosynthetic rate, mg C mg Chl-*a*⁻¹ h⁻¹; α^B , photosynthetic efficiency, mg C mg Chl-*a*⁻¹ h⁻¹ [μmol photons m⁻² s⁻¹]⁻¹. The numbers in parentheses are standard deviations. Winter includes data from January, February, and December. Spring includes data from March, April, and May. Summer includes data from June, July, and August. Autumn includes data from September, October, and November.

test. Linear regression analyses were performed to determine whether there was a long-term trend in the change in SST and nutrient concentration. All statistical analyses were performed using SPSS ver. 27.0 (IBM-SPSS Corp., Armonk, NY, USA). The significance criterion was set at $p < 0.05$.

RESULTS

Photosynthetic parameters

Overall, the maximum photosynthetic rate (P_m^B) and photosynthetic efficiency (α^B) values in the East Sea, eastern South Sea, and waters off Jeju Island were significantly lower than those in the West Sea and western South Sea (t-test, $t_{101} = 2.732$, $p = 0.004$ for P_m^B and Mann-Whitney U = 1,863.5, $p = 0.000$ for α^B). In the East Sea, eastern South Sea, and waters off Jeju Island, the highest mean P_m^B (mg C mg Chl- α^1 h $^{-1}$) was 5.73 in autumn, while the lowest rate was 1.42 in winter (Table 1). The mean P_m^B (mg C mg Chl- α^1 h $^{-1}$) differed significantly depending on the season (ANOVA, $F_{3,41} = 3.736$, $p = 0.018$). Moreover, the mean α^B (mg C mg Chl- α^1 h $^{-1}$ [$\mu\text{mol photons m}^{-2} \text{s}^{-1}$] $^{-1}$) was the highest in spring (0.051), but the lowest in winter (0.020) in these regions (Table 1). Unlike photosynthetic rates, the α^B in one season was not significantly different from that in the other seasons (Welch's ANOVA, $F_{3,15.6} = 2.536$, $p = 0.94$).

In the West and western South Seas, the highest mean photosynthetic rate was 7.50 mg C mg Chl- α^1 h $^{-1}$ in




Fig. 2. Long-term trends in annual mean of sea surface temperature (SST) (A), euphotic depth (B), dissolved inorganic nitrogen (DIN) concentration (C), dissolved inorganic phosphorus (DIP) concentration (D), chlorophyll- α (Chl- α) concentration (E), and primary production (F) in the West Sea. Regression lines were shown with statistical significance.

Table 2. Physicochemical environmental conditions and the primary production (PP) in the West Sea

Year	SST (°C)	Euphotic depth (m)	DIN (µM)	DIP (µM)	Chl- α (µg L⁻¹)	PP (mg C m⁻² d⁻¹)
2005	2.4–25.8 (13.7 ± 7.3)	2.6–19.8 (7.0 ± 3.7)	0.7–31.3 (16.2 ± 7.5)	0.19–1.39 (0.57 ± 0.3)	0.5–9.3 (2.4 ± 1.9)	33–807 (236 ± 214)
2006	2.5–25.7 (12.4 ± 7.3)	1.6–13.1 (6.6 ± 3.1)	1.1–36.9 (16.3 ± 10.4)	0.10–1.32 (0.61 ± 0.3)	0.2–16.7 (1.6 ± 2.7)	10–1,244 (134 ± 203)
2007	4.7–22.9 (13.6 ± 6.4)	0.6–12.5 (5.1 ± 3.2)	1.3–27.9 (15.7 ± 7.0)	0.08–1.06 (0.70 ± 0.2)	0.4–3.1 (1.5 ± 0.6)	19–279 (122 ± 71)
2008	2.4–26.1 (12.6 ± 7.1)	2.6–14.4 (7.1 ± 3.4)	0.4–33.2 (10.2 ± 7.9)	0.13–0.87 (0.53 ± 0.2)	0.2–14.9 (2.8 ± 2.9)	32–1,237 (300 ± 308)
2009	4.2–24.3 (13.6 ± 6.9)	1.6–11.5 (5.0 ± 2.7)	0.3–15.9 (7.7 ± 5.0)	0.10–0.97 (0.48 ± 0.3)	0.0–8.4 (1.7 ± 1.6)	3–477 (134 ± 127)
2010	2.8–26.9 (12.1 ± 7.2)	1.6–20.2 (8.7 ± 5.5)	0.9–19.6 (8.6 ± 5.5)	0.03–0.87 (0.50 ± 0.3)	0.1–2.6 (1.1 ± 0.8)	4–248 (89 ± 69)
2011	1.4–20.7 (11.3 ± 6.9)	1.0–11.2 (5.9 ± 3.3)	3.1–16.5 (11.7 ± 3.5)	0.12–0.87 (0.55 ± 0.2)	0.7–9.2 (2.8 ± 2.3)	39–468 (164 ± 98)
2012	0.7–29.6 (12.3 ± 8.4)	0.6–19.8 (5.4 ± 3.8)	0.2–35.0 (9.6 ± 7.4)	0.00–0.97 (0.39 ± 0.3)	0.0–37.2 (1.8 ± 3.6)	2–1,231 (128 ± 191)
2013	1.2–28.0 (11.7 ± 7.3)	0.3–24.0 (5.2 ± 3.8)	0.2–25.7 (9.6 ± 5.8)	0.00–1.29 (0.44 ± 0.3)	0.0–16.7 (2.1 ± 2.9)	2–1,723 (180 ± 264)
2014	1.4–25.9 (13.5 ± 7.4)	0.3–17.6 (5.5 ± 3.5)	0.7–40.0 (10.7 ± 7.4)	0.00–1.61 (0.57 ± 0.3)	0.2–28.2 (2.0 ± 2.8)	3–1,355 (161 ± 198)
2015	1.8–29.1 (14.0 ± 7.7)	0.3–19.8 (4.9 ± 3.0)	0.7–39.3 (10.7 ± 7.7)	0.00–1.61 (0.53 ± 0.4)	0.1–18.8 (2.2 ± 3.1)	4–2,039 (211 ± 332)
2016	2.4–30.2 (14.0 ± 7.7)	0.3–19.8 (4.8 ± 3.2)	0.7–48.6 (11.3 ± 8.0)	0.00–1.94 (0.62 ± 0.4)	0.1–13.1 (2.2 ± 2.3)	2–1,048 (191 ± 224)
2017	1.9–31.1 (15.3 ± 8.5)	0.3–25.6 (5.5 ± 3.7)	0.1–50.7 (7.9 ± 8.0)	0.00–1.61 (0.45 ± 0.4)	0.4–21.5 (4.0 ± 3.7)	28–2,322 (376 ± 372)
2018	-0.9–30.7 (12.8 ± 8.7)	0.3–26.2 (6.2 ± 3.5)	0.7–50.7 (8.5 ± 8.1)	0.00–2.26 (0.48 ± 0.4)	0.0–25.5 (4.1 ± 4.6)	3–3,118 (407 ± 571)
2019	2.1–28.1 (13.6 ± 7.7)	1.0–24.0 (6.7 ± 3.3)	0.3–37.1 (9.3 ± 7.4)	0.00–14.84 (0.66 ± 0.9)	0.4–20.7 (3.9 ± 3.6)	26–1,777 (358 ± 361)
2020	5.4–22.3 (12.6 ± 5.3)	1.6–22.4 (6.2 ± 4.4)	0.4–15.3 (9.6 ± 3.5)	0.04–0.88 (0.57 ± 0.2)	0.6–3.6 (1.6 ± 0.8)	49–436 (155 ± 117)
2021	3.3–25.4 (13.4 ± 7.2)	3.2–22.4 (6.5 ± 3.7)	0.3–17.3 (9.6 ± 4.8)	0.01–1.13 (0.56 ± 0.3)	0.3–4.0 (1.3 ± 0.6)	28–311 (111 ± 61)

SST, sea surface temperature; DIN, dissolved inorganic nitrogen; DIP, dissolved inorganic phosphorus; Chl- α , chlorophyll- α .

The numbers in parenthesis are mean \pm standard deviation.

spring, whereas the lowest rate was 4.34 mg C mg Chl- α $^{-1}$ h $^{-1}$ in winter (Table 1). The highest mean α^B was 0.168 in autumn, while the lowest α^B was 0.054 in summer (Table 1). The P_m^B and α^B in one season were not significantly different from those in the other seasons (Kruskal-Wallis, $H_3 = 1.858$, $p = 0.602$ for P_m^B and $H_3 = 2.647$, $p = 0.449$ for α^B).

Environmental parameters and calculated PP

From 2005 to 2021, in the West Sea of Korea, the annual mean SST fluctuated slightly from 11.3 to 15.3°C (Fig. 2A). The highest surface temperature was observed in 2017, whereas the lowest was observed in 2018 (Table 2). The calculated annual mean euphotic depth (Z_{eu}) ranged from 4.8 to 8.7 m (Fig. 2B); the deepest Z_{eu} was 26.2 m (Table 2). The annual mean DIN concentration in the surface waters significantly decreased from 2005 to 2021 ($p = 0.007$), whereas there was no distinct trend in the DIP concentration (Fig. 2C & D). The annual mean Chl- α concentration at the surface water fluctuated between 1.1 and 2.8 $\mu\text{g L}^{-1}$ from 2005 to 2016 but maintained in high concentration (3.9–4.1 $\mu\text{g L}^{-1}$) in 2017–2019, and then decreased as low as 1.3 $\mu\text{g L}^{-1}$ in 2021 (Fig. 2E). The calculated mean PP showed a similar pattern to that of the Chl- α concentration, ranging from 89 mg C m $^{-2}$ d $^{-1}$ in 2010 to 407 mg C m $^{-2}$ d $^{-1}$ in 2018 (Fig. 2F).

In the western South Sea of Korea, the annual mean SST from 2005 to 2021 fluctuated slightly from 14.9 to 17.8°C (Fig. 3A). The minimum and maximum SST were -0.2 and 32.0°C in 2018 (Table 3). The annual mean Z_{eu} ranged from 3.2 to 5.6 m (Fig. 3B) and the greatest Z_{eu}




Fig. 3. Long-term trends in annual mean of sea surface temperature (SST) (A), euphotic depth (B), dissolved inorganic nitrogen (DIN) concentration (C), dissolved inorganic phosphorus (DIP) concentration (D), chlorophyll- α (Chl- α) concentration (E), and primary production (F) in the western South Sea. Regression lines were shown with statistical significance.

Table 3. Physicochemical environmental conditions and the primary production (PP) in the western South Sea

Year	SST (°C)	Euphotic depth (m)	DIN (μM)	DIP (μM)	Chl- α ($\mu\text{g L}^{-1}$)	PP (mg C m $^{-2}$ d $^{-1}$)
2005	2.4–28.5 (16.2 ± 6.4)	0.3–14.0 (4.9 ± 3.0)	1.1–31.3 (9.4 ± 5.4)	0.07–2.43 (0.56 ± 0.4)	0.2–8.8 (2.2 ± 1.6)	3–779 (186 ± 157)
2006	2.7–27.7 (14.9 ± 6.4)	0.3–10.2 (4.7 ± 2.4)	1.9–41.1 (13.4 ± 7.2)	0.06–5.84 (0.71 ± 0.6)	0.0–36.2 (2.5 ± 3.9)	1–1,675 (188 ± 206)
2007	6.1–28.5 (16.4 ± 6.2)	0.3–12.8 (3.2 ± 2.4)	1.5–47.8 (10.7 ± 8.5)	0.00–3.87 (0.48 ± 0.6)	0.0–23.7 (4.6 ± 3.6)	0–1,358 (363 ± 315)
2008	2.8–29.0 (15.2 ± 6.7)	0.6–13.6 (4.4 ± 2.9)	1.5–28.0 (8.2 ± 4.6)	0.00–2.16 (0.31 ± 0.4)	1.1–24.0 (5.2 ± 4.1)	42–1,842 (450 ± 388)
2009	4.9–28.6 (16.3 ± 5.9)	0.3–16.0 (4.0 ± 3.0)	0.0–18.4 (7.3 ± 4.3)	0.00–0.81 (0.28 ± 0.2)	0.1–24.9 (3.3 ± 3.6)	1–1,394 (205 ± 193)
2010	2.9–29.3 (15.3 ± 6.7)	0.3–16.0 (4.7 ± 2.8)	0.1–12.9 (4.7 ± 3.8)	0.00–0.68 (0.27 ± 0.2)	0.1–8.0 (2.5 ± 1.6)	14–633 (205 ± 151)
2011	2.0–27.3 (15.0 ± 6.7)	0.6–14.4 (5.3 ± 2.8)	0.0–25.8 (4.4 ± 4.3)	0.00–1.29 (0.36 ± 0.3)	0.2–12.5 (2.1 ± 2.1)	16–864 (156 ± 139)
2012	2.3–30.0 (15.4 ± 8.1)	0.3–24.0 (4.6 ± 2.7)	0.0–22.1 (4.3 ± 4.2)	0.00–0.97 (0.30 ± 0.2)	0.2–14.3 (2.2 ± 1.9)	4–1,032 (152 ± 142)
2013	2.4–31.8 (16.0 ± 7.5)	0.3–29.6 (4.0 ± 2.6)	0.0–22.1 (6.6 ± 4.7)	0.00–1.40 (0.37 ± 0.3)	0.3–17.3 (2.4 ± 2.1)	6–1,180 (195 ± 176)
2014	3.4–26.9 (15.5 ± 6.2)	0.6–13.5 (4.2 ± 2.2)	0.2–35.7 (6.2 ± 4.5)	0.00–0.97 (0.31 ± 0.2)	0.1–88.2 (7.7 ± 11.4)	5–5,410 (566 ± 768)
2015	3.6–31.1 (16.2 ± 7.1)	0.3–17.5 (4.2 ± 2.2)	0.3–38.6 (9.2 ± 6.7)	0.00–0.97 (0.39 ± 0.2)	0.2–167.1 (11.9 ± 17.3)	2–11,805 (940 ± 1,280)
2016	1.6–31.8 (16.3 ± 7.7)	0.3–26.9 (4.3 ± 3.5)	0.5–27.9 (5.4 ± 4.6)	0.00–1.94 (0.30 ± 0.3)	0.1–72.9 (7.1 ± 10.1)	2–5,287 (585 ± 843)
2017	4.2–30.0 (17.8 ± 7.6)	0.3–14.0 (4.4 ± 2.3)	0.0–20.7 (4.3 ± 3.8)	0.00–1.61 (0.28 ± 0.3)	0.0–62.9 (3.4 ± 4.5)	1–5,456 (292 ± 369)
2018	-0.2–32.0 (15.7 ± 8.1)	1.3–14.4 (4.8 ± 2.2)	0.0–23.6 (5.8 ± 4.5)	0.00–1.29 (0.39 ± 0.3)	0.1–23.5 (2.7 ± 3.0)	7–2,320 (239 ± 276)
2019	2.9–30.7 (15.2 ± 6.9)	0.3–15.4 (4.7 ± 2.5)	0.1–27.1 (5.5 ± 4.0)	0.00–1.42 (0.39 ± 0.2)	0.1–15.5 (3.0 ± 2.9)	8–1,519 (257 ± 265)
2020	5.9–25.7 (15.9 ± 5.1)	0.3–15.0 (4.7 ± 3.2)	0.4–12.6 (5.8 ± 2.9)	0.01–0.72 (0.38 ± 0.2)	0.1–17.9 (2.4 ± 3.3)	10–1,178 (169 ± 202)
2021	5.4–29.6 (17.2 ± 6.9)	1.0–16.0 (5.6 ± 3.7)	0.3–16.4 (6.0 ± 3.8)	0.00–1.02 (0.41 ± 0.2)	0.1–7.4 (1.5 ± 1.3)	6–561 (114 ± 98)

SST, sea surface temperature; DIN, dissolved inorganic nitrogen; DIP, dissolved inorganic phosphorus; Chl- α , chlorophyll- α .

The numbers in parenthesis are mean ± standard deviation.

among the sites was 29.6 m in 2013 (Table 3). The annual mean DIN concentration in this region slightly decreased during the 17-year period ($p = 0.012$), whereas the annual mean DIP concentration showed no distinct trend (Fig. 3C & D). The minimum and maximum mean concentrations for DIN and DIP were 4.3 and 13.4 μM and 0.27 and 0.71 μM , respectively. An annual mean Chl- a concentration lower than 5.2 $\mu\text{g L}^{-1}$ was maintained from 2005 to 2013, but largely increased to 11.9 $\mu\text{g L}^{-1}$ in 2015 and then decreased to 1.5 $\mu\text{g L}^{-1}$ in 2021 (Fig. 3E). The mean PP trend in the western South Sea was similar to that of Chl- a . The PP ranged from 114 $\text{mg C m}^{-2} \text{d}^{-1}$ in 2021 to 940 $\text{mg C m}^{-2} \text{d}^{-1}$ in 2015 (Fig. 3F).

In the eastern South Sea of Korea, the annual mean SST increased slightly from 2005 to 2021, ranging from 15.2 to 19.2°C ($p = 0.045$) (Fig. 4A). The minimum SST was 3.3°C in 2018, but the maximum SST was 31.6°C in 2013 (Table 4). The annual mean Z_{eu} was between 10.7 m in 2008 and 15.0 m in 2012 (Fig. 4B) and the maximum Z_{eu} was 64.0 m in 2012 (Table 4). The annual mean concentrations of DIN and DIP decreased slightly during 2005–2021 ($p = 0.001$ for DIN and $p = 0.003$ for DIP) (Fig. 4C & D). The minimum and maximum mean DIN concentrations were 3.5 μM in 2021 and 7.0 μM in 2007, respectively, while those of DIP were 0.28 μM in 2021 and 0.56 μM in 2007, respectively (Fig. 4). The annual mean Chl- a concentration fluctuated from 2005 to 2021, ranging from 2.0 $\mu\text{g L}^{-1}$ in 2021 to 4.3 $\mu\text{g L}^{-1}$ in 2015 (Fig. 4E). Similarly, the PP in the Korea Strait fluctuated between 69 $\text{mg C m}^{-2} \text{d}^{-1}$ in 2021 and 183 $\text{mg C m}^{-2} \text{d}^{-1}$ in 2015 (Fig. 4F).

In the East Sea of Korea, the annual mean SST gradu-




Fig. 4. Long-term trends in annual mean of sea surface temperature (SST) (A), euphotic depth (B), dissolved inorganic nitrogen (DIN) concentration (C), dissolved inorganic phosphorus (DIP) concentration (D), chlorophyll- a (Chl- a) concentration (E), and primary production (F) in the eastern South Sea. Regression lines were shown with statistical significance.

Table 4. Physicochemical environmental conditions and the primary production (PP) in the eastern South Sea

Year	SST (°C)	Euphotic depth (m)	DIN (μM)	DIP (μM)	Chl- a ($\mu\text{g L}^{-1}$)	PP ($\text{mg C m}^{-2} \text{d}^{-1}$)
2005	5.3–29.8 (16.5 ± 5.2)	3.2–38.4 (13.7 ± 6.6)	0.6–19.4 (4.5 ± 3.5)	0.00–1.64 (0.41 ± 0.3)	0.0–14.1 (2.1 ± 1.9)	1–419 (77 ± 79)
2006	5.4–31.3 (16.1 ± 5.4)	3.2–35.2 (11.9 ± 5.7)	0.0–22.9 (6.2 ± 4.7)	0.00–2.29 (0.52 ± 0.5)	0.1–15.6 (2.7 ± 2.9)	4–722 (108 ± 138)
2007	6.5–25.8 (16.8 ± 4.7)	3.2–38.4 (13.4 ± 6.3)	0.7–32.0 (7.0 ± 5.3)	0.01–2.00 (0.56 ± 0.4)	0.1–79.9 (4.0 ± 6.4)	3–2,811 (148 ± 235)
2008	5.0–29.3 (16.5 ± 5.2)	0.0–34.3 (10.7 ± 6.2)	0.5–18.0 (5.8 ± 4.1)	0.00–1.95 (0.52 ± 0.4)	0.2–12.9 (2.6 ± 2.4)	4–477 (88 ± 89)
2009	5.7–27.0 (16.7 ± 5.1)	3.2–30.4 (11.5 ± 5.1)	0.3–15.9 (5.5 ± 3.5)	0.01–1.26 (0.30 ± 0.2)	0.1–13.9 (2.4 ± 2.3)	1–697 (80 ± 107)
2010	5.1–28.1 (16.3 ± 5.3)	1.6–35.2 (12.2 ± 6.0)	0.1–18.9 (4.6 ± 3.6)	0.00–1.12 (0.29 ± 0.2)	0.1–24.4 (3.0 ± 2.9)	1–954 (99 ± 110)
2011	3.5–26.6 (15.2 ± 5.6)	3.2–44.0 (12.7 ± 5.8)	0.0–34.3 (4.0 ± 4.2)	0.00–1.61 (0.34 ± 0.3)	0.0–57.7 (3.9 ± 5.0)	1–1,880 (122 ± 154)
2012	3.8–29.5 (16.4 ± 6.9)	2.2–64.0 (15.0 ± 8.4)	0.0–52.1 (4.4 ± 4.9)	0.00–3.87 (0.34 ± 0.4)	0.0–235.1 (3.3 ± 10.5)	0–3,709 (87 ± 175)
2013	4.9–31.6 (16.7 ± 6.0)	1.5–44.8 (13.4 ± 7.1)	0.0–33.6 (4.9 ± 4.8)	0.00–2.90 (0.35 ± 0.4)	0.0–205.7 (3.8 ± 8.3)	2–7,017 (133 ± 296)
2014	5.5–27.7 (16.3 ± 5.2)	1.5–51.2 (12.9 ± 6.7)	0.0–52.1 (5.2 ± 5.1)	0.00–1.61 (0.36 ± 0.3)	0.0–32.9 (3.6 ± 4.3)	1–1,371 (136 ± 186)
2015	5.4–31.4 (17.0 ± 6.0)	2.6–48.0 (12.1 ± 6.3)	0.0–37.1 (5.5 ± 5.2)	0.00–1.54 (0.37 ± 0.3)	0.1–50.6 (4.3 ± 5.9)	1–3,894 (183 ± 394)
2016	4.1–30.0 (16.6 ± 6.3)	0.6–43.0 (12.8 ± 6.7)	0.3–27.9 (4.2 ± 3.9)	0.00–1.61 (0.31 ± 0.3)	0.1–84.5 (3.7 ± 6.4)	2–2,216 (117 ± 169)
2017	5.6–29.6 (19.2 ± 6.4)	2.0–51.2 (13.6 ± 6.7)	0.0–24.3 (3.6 ± 3.5)	0.00–1.61 (0.31 ± 0.3)	0.1–18.3 (2.6 ± 2.5)	1–835 (119 ± 120)
2018	3.3–30.9 (17.4 ± 7.0)	2.6–54.4 (13.1 ± 7.0)	0.0–35.0 (4.4 ± 4.2)	0.00–1.94 (0.34 ± 0.3)	0.0–52.2 (2.7 ± 3.2)	0–841 (96 ± 104)
2019	6.2–30.3 (17.2 ± 5.8)	2.0–58.0 (14.2 ± 7.0)	0.0–17.9 (3.5 ± 3.3)	0.00–1.94 (0.32 ± 0.3)	0.2–22.3 (2.2 ± 2.0)	6–1,217 (87 ± 106)
2020	8.5–26.7 (16.2 ± 3.7)	3.2–41.0 (12.8 ± 6.1)	0.1–17.2 (3.9 ± 3.2)	0.00–1.24 (0.31 ± 0.2)	0.3–9.7 (2.6 ± 1.8)	4–336 (91 ± 72)
2021	7.7–30.0 (18.1 ± 6.1)	3.0–41.6 (13.9 ± 6.3)	0.2–17.2 (3.5 ± 3.3)	0.00–1.12 (0.28 ± 0.2)	0.1–6.4 (2.0 ± 1.3)	4–255 (69 ± 54)

SST, sea surface temperature; DIN, dissolved inorganic nitrogen; DIP, dissolved inorganic phosphorus; Chl- a , chlorophyll- a .

The numbers in parenthesis are mean ± standard deviation.

ally increased from 14.2 to 17.5°C during the study period ($p < 0.001$) (Fig. 5A). The minimum and maximum SST were 2.3°C in 2011 and 29.3°C in 2018, respectively (Table 5). The annual mean Z_{eu} was between 23.0 and 31.7 m in 2020 and 2015, respectively (Fig. 5B), and the maximum Z_{eu} was 73.6 m in 2016 (Table 5). The annual mean concentrations of DIN and DIP decreased slightly during the study period ($p = 0.002$ for both DIN and DIP) (Fig. 5C & D). The maximum and minimum mean DIN concentrations were 7.4 µM in 2006 and 2.4 µM in 2017, respectively (Fig. 5C). Meanwhile, the maximum and minimum mean DIP concentrations were 0.36 µM in 2005 and 0.17 µM in 2015, respectively (Fig. 5D). The annual mean Chl-*a* concentrations were maintained between 0.5 and 2.2 µg L⁻¹ observed in 2015 and 2020, respectively (Fig. 5E). The PP in the East Sea also changed in the range of 17 and 78 mg C m⁻² d⁻¹, and the minimum and maximum values were obtained in 2015 and 2020, respectively (Fig. 5F).

In the waters off Jeju Island, the annual mean SST showed no significant change during the study period, ranging from 18.2 to 20.1°C (Fig. 6A). The minimum and maximum SST were 8.5°C in 2015 and 31.1°C in 2018, respectively (Table 6). The annual mean Z_{eu} fluctuated until 2015, reaching the shallowest at 17.1 m, but increased in recent years. The deepest annual mean Z_{eu} was 52.0 m in 2017 (Fig. 6B). The annual mean concentrations of DIN and DIP tended to increase, except in recent years. The maximum and minimum annual mean DIN were 12.9 µM in 2015 and 3.1 µM in 2020, respectively (Fig. 6C). Meanwhile, the maximum and minimum DIP concentrations were 0.59 µM in 2012 and 0.16 µM in 2007, respectively.




Fig. 5. Long-term trends in annual mean of sea surface temperature (SST) (A), euphotic depth (B), dissolved inorganic nitrogen (DIN) concentration (C), dissolved inorganic phosphorus (DIP) concentration (D), chlorophyll-*a* (Chl-*a*) concentration (E), and primary production (F) in the East Sea. Regression lines were shown with statistical significance.

Table 5. Physicochemical environmental conditions and the primary production (PP) in the East Sea

Year	SST (°C)	Euphotic depth (m)	DIN (µM)	DIP (µM)	Chl- <i>a</i> (µg L ⁻¹)	PP (mg C m ⁻² d ⁻¹)
2005	4.2–25.6 (14.2 ± 5.8)	8.0–48.0 (27.4 ± 8.4)	0.6–20.8 (4.5 ± 3.1)	0.03–1.77 (0.36 ± 0.3)	0.1–7.5 (1.3 ± 1.2)	1–439 (48 ± 55)
2006	3.5–27.7 (13.9 ± 6.1)	8.0–49.0 (26.5 ± 7.3)	0.4–33.8 (7.4 ± 4.6)	0.01–1.06 (0.28 ± 0.2)	0.1–4.4 (1.2 ± 0.6)	1–201 (43 ± 31)
2007	7.4–22.5 (15.1 ± 3.3)	8.0–48.0 (26.1 ± 6.9)	0.4–15.8 (4.3 ± 2.4)	0.00–0.94 (0.28 ± 0.2)	0.0–7.8 (1.1 ± 1.2)	1–390 (39 ± 53)
2008	8.0–26.2 (15.8 ± 5.1)	8.0–51.2 (26.5 ± 8.7)	0.5–15.4 (4.4 ± 3.1)	0.02–0.74 (0.29 ± 0.2)	0.0–8.6 (1.4 ± 1.4)	2–377 (61 ± 81)
2009	6.1–25.9 (15.8 ± 4.9)	8.0–48.0 (24.1 ± 7.5)	0.2–14.6 (4.4 ± 3.5)	0.01–1.30 (0.31 ± 0.2)	0.0–3.4 (1.2 ± 0.6)	0–155 (37 ± 33)
2010	3.4–24.2 (14.4 ± 5.3)	4.8–44.8 (23.7 ± 8.0)	0.4–10.0 (4.8 ± 2.8)	0.00–0.61 (0.18 ± 0.2)	0.5–18.3 (1.7 ± 1.6)	5–692 (64 ± 68)
2011	2.3–25.1 (14.5 ± 5.9)	3.2–57.6 (24.5 ± 9.0)	0.3–11.9 (3.2 ± 2.3)	0.00–0.62 (0.21 ± 0.2)	0.2–6.9 (1.5 ± 1.4)	5–115 (35 ± 25)
2012	6.9–28.1 (16.3 ± 6.1)	9.6–57.6 (28.7 ± 9.6)	0.2–11.4 (3.8 ± 2.9)	0.00–0.65 (0.19 ± 0.2)	0.0–4.5 (0.9 ± 0.6)	1–146 (28 ± 26)
2013	5.8–28.4 (17.0 ± 5.9)	1.6–54.4 (26.6 ± 7.8)	0.1–15.9 (3.3 ± 2.8)	0.00–0.65 (0.18 ± 0.2)	0.0–32.4 (1.8 ± 4.3)	1–1,391 (70 ± 184)
2014	4.7–24.6 (16.3 ± 4.9)	6.4–54.4 (28.8 ± 8.3)	0.0–22.1 (4.8 ± 5.3)	0.00–0.97 (0.18 ± 0.2)	0.0–18.8 (0.8 ± 1.3)	1–532 (31 ± 51)
2015	5.1–26.8 (15.3 ± 5.9)	9.6–51.2 (31.7 ± 8.7)	0.0–10.6 (3.2 ± 2.9)	0.00–0.65 (0.17 ± 0.2)	0.0–2.7 (0.5 ± 0.5)	1–159 (17 ± 22)
2016	4.7–28.5 (15.7 ± 5.5)	7.0–73.6 (29.6 ± 9.4)	0.0–23.6 (2.9 ± 2.9)	0.00–0.65 (0.18 ± 0.2)	0.0–5.0 (0.9 ± 0.9)	2–266 (33 ± 39)
2017	7.9–28.6 (17.5 ± 5.5)	6.4–56.0 (27.3 ± 9.5)	0.0–10.5 (2.4 ± 2.4)	0.00–0.65 (0.19 ± 0.2)	0.0–7.8 (1.1 ± 1.2)	0–431 (54 ± 71)
2018	2.7–29.3 (16.4 ± 6.2)	6.5–57.6 (27.0 ± 9.3)	0.0–13.6 (3.4 ± 3.4)	0.00–0.97 (0.20 ± 0.3)	0.0–18.1 (1.2 ± 1.3)	0–790 (49 ± 62)
2019	5.2–27.3 (16.9 ± 5.2)	7.0–57.6 (26.8 ± 9.1)	0.0–10.7 (3.0 ± 2.6)	0.00–0.73 (0.18 ± 0.2)	0.0–16.1 (1.3 ± 1.5)	3–1,076 (53 ± 79)
2020	10.8–24.2 (16.3 ± 3.8)	6.0–49.0 (23.0 ± 8.3)	0.1–8.7 (2.9 ± 2.6)	0.00–0.55 (0.17 ± 0.2)	0.4–8.8 (2.2 ± 1.6)	8–336 (78 ± 70)
2021	6.4–27.8 (16.3 ± 5.9)	6.4–57.6 (25.4 ± 8.5)	0.3–11.0 (3.4 ± 2.9)	0.00–0.58 (0.24 ± 0.2)	0.2–5.3 (1.5 ± 1.1)	6–243 (57 ± 54)

SST, sea surface temperature; DIN, dissolved inorganic nitrogen; DIP, dissolved inorganic phosphorus; Chl-*a*, chlorophyll-*a*.

The numbers in parenthesis are mean ± standard deviation.

(Fig. 6D). The annual mean Chl-*a* concentration ranged from 0.6 µg L⁻¹ in 2019 to 1.9 µg L⁻¹ in 2015 (Fig. 6E). Similar to Chl-*a* concentration, the PP fluctuated between 24 and 68 mg C m⁻² d⁻¹ in 2009 and 2015, respectively (Fig. 6F).

The annual total PP in the territorial waters (TPP, 76,450 km²) (Fig. 1) was calculated by multiplying 365 days and the area of each region by the daily depth integrated mean PP and then summing the PP in each region. The TPP fluctuated between 2,981 Gg C y⁻¹ in 2012 and 7,278 Gg C y⁻¹ in 2014, but increased to 11,227 Gg C y⁻¹ in 2015, and then decreased to 2,542 Gg C y⁻¹ in 2021 (Fig. 7A). In 2015, when the TPP was the greatest among the studied years, the annual PP in the western South Sea accounted for the greatest, followed by the West Sea (Fig. 7B). The annual PPs in the western South Sea and West Sea were 343,024 and 76,834 kg C km⁻² y⁻¹, respectively (Fig. 7C). The annual PP in the East Sea in 2015 was the lowest (1% of the TPP and 6,072 kg C km⁻² y⁻¹) among the five regions of the territorial waters (Fig. 7B & C).

DISCUSSION

The photosynthetic rate and efficiency are essential parameters to evaluate the photosynthetic performance of the phytoplankton community and to analyze primary productivity. In our results, the photosynthetic rates in a season in the East Sea, the eastern South Sea, and the waters off Jeju Island were significantly different from those in the other seasons ($p = 0.018$), in agreement with




Fig. 6. Long-term trends in annual mean of sea surface temperature (SST) (A), euphotic depth (B), dissolved inorganic nitrogen (DIN) concentration (C), dissolved inorganic phosphorus (DIP) concentration (D), chlorophyll-*a* (Chl-*a*) concentration (E), and primary production (F) in the waters off Jeju Island. Regression lines were shown with statistical significance.

Table 6. Physicochemical environmental conditions and the primary production (PP) in the waters off Jeju Island

Year	SST (°C)	Euphotic depth (m)	DIN (µM)	DIP (µM)	Chl- <i>a</i> (µg L ⁻¹)	PP (mg C m ⁻² d ⁻¹)
2005	14.1–26.7 (19.5 ± 4.3)	14.0–44.8 (24.3 ± 6.8)	1.3–12.1 (5.3 ± 1.9)	0.00–0.85 (0.28 ± 0.2)	0.2–4.0 (1.2 ± 0.8)	4–232 (48 ± 53)
2006	13.2–28.7 (18.4 ± 4.6)	14.0–46.0 (25.5 ± 7.6)	1.7–25.1 (6.3 ± 3.2)	0.00–1.35 (0.41 ± 0.4)	0.0–3.4 (0.9 ± 0.6)	2–165 (28 ± 27)
2007	14.6–28.2 (19.4 ± 4.4)	14.0–46.0 (24.9 ± 7.7)	0.9–16.1 (5.9 ± 3.3)	0.00–0.40 (0.16 ± 0.1)	0.3–5.0 (1.1 ± 0.7)	9–172 (35 ± 26)
2008	13.8–27.4 (19.0 ± 4.4)	14.0–45.7 (26.0 ± 7.8)	0.6–30.1 (7.8 ± 5.3)	0.03–0.39 (0.16 ± 0.1)	0.3–5.5 (1.3 ± 1.1)	12–245 (63 ± 66)
2009	14.2–26.9 (19.1 ± 4.1)	14.0–44.8 (24.6 ± 7.0)	0.8–12.6 (6.7 ± 3.5)	0.02–0.46 (0.18 ± 0.1)	0.3–2.8 (0.9 ± 0.6)	5–81 (24 ± 16)
2010	13.8–28.6 (19.6 ± 5.0)	14.0–43.0 (23.9 ± 7.2)	1.9–14.4 (6.2 ± 2.9)	0.05–0.44 (0.22 ± 0.1)	0.6–3.4 (1.3 ± 0.6)	8–147 (50 ± 34)
2011	13.1–25.8 (19.0 ± 4.2)	1.6–44.8 (17.2 ± 8.5)	0.7–30.0 (7.8 ± 6.2)	0.00–0.97 (0.24 ± 0.2)	0.0–6.5 (1.5 ± 1.0)	7–315 (45 ± 41)
2012	12.7–27.8 (18.6 ± 3.7)	9.0–40.0 (23.6 ± 6.4)	0.0–9.8 (3.7 ± 2.2)	0.00–9.68 (0.59 ± 1.4)	0.4–4.7 (1.3 ± 1.0)	7–142 (41 ± 35)
2013	12.6–29.9 (19.3 ± 4.7)	5.0–48.0 (19.0 ± 7.8)	0.3–121.4 (9.3 ± 15.0)	0.00–4.19 (0.50 ± 0.7)	0.2–11.6 (1.7 ± 1.5)	8–581 (67 ± 77)
2014	11.8–26.8 (18.2 ± 3.3)	4.4–50.0 (19.4 ± 7.7)	0.7–122.9 (8.0 ± 12.8)	0.00–5.16 (0.34 ± 0.5)	0.0–5.1 (1.7 ± 1.1)	2–235 (62 ± 53)
2015	8.5–27.3 (18.6 ± 4.2)	0.3–35.2 (17.1 ± 6.4)	0.4–113.6 (12.9 ± 17.9)	0.00–5.81 (0.40 ± 0.7)	0.2–5.1 (1.9 ± 1.1)	1–223 (68 ± 57)
2016	13.2–28.4 (18.5 ± 3.7)	7.0–46.0 (18.6 ± 7.1)	1.2–85.7 (11.8 ± 13.9)	0.00–3.23 (0.36 ± 0.5)	0.0–3.7 (1.6 ± 0.8)	2–136 (53 ± 40)
2017	13.7–29.3 (20.1 ± 4.7)	8.4–52.0 (21.4 ± 8.2)	0.1–121.4 (12.9 ± 20.6)	0.00–3.55 (0.46 ± 0.6)	0.0–3.1 (0.7 ± 0.7)	0–183 (33 ± 34)
2018	12.9–31.1 (18.9 ± 4.2)	7.0–39.4 (19.7 ± 6.2)	0.1–81.4 (10.2 ± 14.8)	0.00–6.13 (0.58 ± 0.9)	0.0–5.7 (0.8 ± 0.8)	1–256 (35 ± 41)
2019	11.3–28.5 (19.5 ± 4.5)	7.0–46.0 (21.5 ± 6.9)	0.1–65.0 (8.2 ± 12.5)	0.00–3.87 (0.42 ± 0.7)	0.0–2.4 (0.6 ± 0.5)	1–104 (25 ± 25)
2020	14.5–29.4 (19.8 ± 4.6)	9.6–42.0 (22.6 ± 6.9)	0.3–14.0 (3.1 ± 2.5)	0.00–0.48 (0.17 ± 0.2)	0.1–7.7 (1.4 ± 1.3)	2–369 (56 ± 65)
2021	14.8–26.8 (19.4 ± 3.7)	10.0–41.6 (24.0 ± 7.1)	0.3–19.5 (4.1 ± 2.8)	0.00–0.45 (0.17 ± 0.1)	0.1–9.2 (1.3 ± 1.4)	2–211 (43 ± 43)

SST, sea surface temperature; DIN, dissolved inorganic nitrogen; DIP, dissolved inorganic phosphorus; Chl-*a*, chlorophyll-*a*.

The numbers in parenthesis are mean ± standard deviation.




Fig. 7. Trend of annual primary production (PP) in the terrestrial waters (A), the percentage (B) and the amount (C) of PP of each region in 2015. The black bar in (A) indicates the year showing the highest PP during the study period. WS, West Sea; WSS, western South Sea; ESS, eastern South Sea; ES, East Sea; JI, waters off Jeju Island.

the fact that the photosynthetic rate is affected by water temperature and / or nutrient concentration (Park and Kim 2010). Moreover, the photosynthetic rates in the East Sea, eastern South Sea, and waters off Jeju Island were significantly lower than those in the West Sea and western South Sea ($p = 0.004$). Therefore, the results showed that different photosynthetic rates according to environmental conditions should be applied to the models or algorithms developed for estimating PP. Estimation of photosynthetic parameters based on the ecological environment should also be continued to accurately measure PP. The seasonal mean values of the photosynthetic rates ranged from 1.42 to 7.50 mg C mg Chl- α^{-1} h $^{-1}$ in the Korean coastal waters, showing similar ranges to the values in the various marine environments ranging from 0.4 to 10 mg C mg Chl- α^{-1} h $^{-1}$ (Eppley 1972). In particular, the values in the present study considerably overlapped the photosynthetic rates of phytoplankton in upwelling regions when water temperatures were between 17–25°C and ranged from 5 to 10 mg C Chl- α^{-1} h $^{-1}$ (Eppley 1972).

The photosynthetic efficiencies of the phytoplankton communities from the West Sea and western South Seas were significantly higher than those of the East Sea, eastern South Sea, and waters off Jeju Island ($p = 0.000$). This means that the phytoplankton communities in the West and western South Seas likely assimilated carbon more

efficiently through photosynthesis under the same light conditions. According to Côté and Platt (1983), photosynthetic efficiency is correlated with community structure, such as species composition, cell size, and species diversity. The disparate phytoplankton communities derived from the different environmental conditions between regions may result in differences in photosynthetic efficiency.

The annual mean SSTs in the eastern South Sea and East Sea significantly increased during the 17-year study period (Figs 4 & 5). Although the increase in the annual mean temperatures in other regions was not significant in the present study, there was a tendency to increase the SST in the Korean coastal waters, which is in agreement with previous studies (Han and Lee 2020, Lee and Park 2020).

The calculated euphotic depths did not exhibit any particular temporal pattern. However, it was clearly shown that the euphotic depths in the West and western South Seas were shallower than those in the eastern South Sea, East Sea, and waters off Jeju Island. This is not only because the average water depth in the West Sea and western South Sea of Korea (mostly less than 30 m, <http://www.mels.go.kr>) is shallower than that in the eastern South Sea and / or East Sea, but also because there are several large rivers in the western part of Korea. Moreover, the coastlines of the West Sea and western South Sea of Korea are characterized by a complicated coastline consisting of numerous islands, extensive tidal flats, and relatively strong tides (Kang et al. 2009). Strong tidal currents and mixing over shallow bathymetry may increase turbidity (Park et al. 1998) and cause shallow euphotic depths in the West and western South Seas of Korea.

The concentrations of DIN in the Korean coastal waters were significantly decreased, except in the Jeju area, indicating a reduction in nutrient concentrations in the coastal waters of the Korean Peninsula (Figs 2–6). Moreover, DIP concentrations in the eastern South Sea and East Sea decreased significantly. These decreases in nutrients in the Korean coastal waters have been reported in many studies (Kim et al. 2013, Ministry of Oceans and Fisheries and Korea Marine Environment Management Corporation 2022). Several processes, such as input from rivers, submarine groundwater discharge, and topographic environmental changes, may contribute to the dynamics of nutrient concentrations in Korean coastal waters (Hwang et al. 2005, Kim et al. 2013). For example, the reduction of nutrients in river discharge is partially responsible for decreasing nutrients in Korean coastal waters (Kim et al. 2013). Massive civil constructions, such as tidal power

plants or sea dikes, are also suggested as possible reasons for the reduction of nutrients by decreasing nutrient input from sediments due to the decrease in tidal mixing (Kim et al. 2013).

Unlike other regions, the nutrient concentrations in the waters off Jeju Island tended to increase, except in recent years (Fig. 6). Several researchers have studied the quality and eutrophication of coastal waters off Jeju Island, and some nutrient sources have been suggested as possible reasons for the high concentration of nutrients. For example, there are concentrated inland aquacultures

on Jeju Island, accounting for more than 50% of Korea in 2021 (KOSIS 2022), and thus the discharge of aquaculture effluent water has effects on the nutrient concentration in the coastal area of Jeju (Koh et al. 2013, Oh et al. 2021). Nutrient concentrations in the coastal area of Jeju are influenced by the submarine groundwater discharge (Hwang et al. 2005). Moreover, the excrement of migrating birds may be partially responsible for the high concentration of organic matter in the local coastal waters in Jeju (Kim and Kim 2017).

The trend in the Chl- α concentration differed across

Table 7. Comparison of the primary production (PP) in each region

Region	Year	Month	PP ($\text{mg C m}^{-2} \text{d}^{-1}$)	Method	Reference
West Sea	1989	3, 5, 6, 8, 10	37-708	^{14}C	Kang et al. (1992)
	1990	2, 3, 5, 7, 8, 10	37-1,104	^{14}C	Kang et al. (1992)
	1992	1, 4, 8, 10	45-2,523	^{14}C	Lee and Choi (2000)
	1994	7	1,324	^{14}C	Yoo and Shin (1995)
	1995	1, 10	28-2,698	^{14}C	Yoo and Shin (1995), Choi et al. (1997)
	1996	3, 8	1,382-1,888	^{14}C	Choi et al. (1997)
	2004	4	94	^{13}C	Lee et al. (2006)
	2006	4	100-700	^{14}C	Yoo et al. (2018)
	2008	10	226-521	^{14}C	Park and Kim (2010)
	2016	8	67-456	^{13}C	Jang et al. (2018)
	2018	2, 4, 8, 10	68-487	^{13}C	Jang et al. (2021)
	2005-2021	All	2-3,118	Calculated	This study
Western South Sea	1992	9	620	^{14}C	Choi et al. (1995)
	2004	3	51-67	^{13}C	Kim et al. (2016)
	2009	2, 5, 7, 8, 11	9-8,927	^{13}C	Lee et al. (2011), Min et al. (2011, 2012)
	1992-1998	5, 9	554-734	^{14}C	Son et al. (2005)
	2011-2012	7-12	17-1,052	Numerical model	Jeong and Cho (2018)
	2005-2021	All	0.4-11,805	Calculated	This study
Eastern South Sea	1985	10	700-2,740	^{14}C	Shim and Park (1986)
	1992	4, 7, 10, 12	317-3,539	^{14}C	Yeo et al. (1996)
	1994	10-12	72-656	Calculated	Jeong et al. (2009)
	1995	All	32-1,527	^{14}C , calculated	Han et al. (1998), Jeong et al. (2009)
	1996	1-4	151-917	Calculated	Jeong et al. (2009)
	2013-2014	All	160-2,880	^{14}C	Kim et al. (2019)
	2005-2021	All	0.4-3,894	Calculated	This study
East Sea	1984	4, 5, 10	204-637	Calculated	Shim et al. (1985)
	1986	10	750-2,040	^{14}C	Chung et al. (1989)
	1988	5	284-4,574	^{14}C	Shim et al. (1992)
	1989	7	444-3,605	^{14}C	Shim et al. (1992)
	1990	4, 8	310-3,909	^{14}C	Park et al. (1991), Shim et al. (1992)
	2006	4	159-777	^{14}C	Hyun et al. (2009)
	2008	6, 8, 10	280-2,410	^{13}C	Kwak et al. (2013a)
	2009	5	490-940	^{13}C	Kwak et al. (2013a)
	2010	5, 11	550-1,074	^{13}C	Kwak et al. (2013b)
	2011	6	1,026	^{13}C	Kwak et al. (2013b)
	2012	10	181	^{13}C	Lee et al. (2017)
	2015	5	442	^{13}C	Lee et al. (2017)
	2016	4	378-986	^{13}C	Kang et al. (2020)
Waters off Jeju Island	2003-2012	All	447-1,263	MODIS-derived	Joo et al. (2014)
	2005-2021	All	0.03-1,391	Calculated	This study
	1989	4, 11	517-1,727	^{14}C	Chung and Yang (1991)
	1990	3	314	^{14}C	Chung and Yang (1991)
	1991	9, 11	25-145	^{14}C	Lee et al. (1993)
	1992	1, 3, 5, 6	4-805	^{14}C	Lee et al. (1993)
	2005-2021	All	0.2-581	Calculated	This study

regions. However, abnormally high Chl-*a* concentrations were observed from 2014 to 2016 in the western South Sea of Korea (Fig. 3). Heavy phytoplankton blooms frequently occur in the western South Sea of Korea. During 2014–2015, there also were massive algal bloom events caused by *Margalefidinium* (previously known as *Cochlodinium*) (Jeong et al. 2017, National Institute of Fisheries Science 2020). Even if the data for 2014–2016 were not included, the overall concentrations of Chl-*a* (1.5–5.2 µg L⁻¹) (Fig. 3) were greater than those in the East Sea and waters of Jeju Island, of which the maximum values were 2.2 and 1.9 µg L⁻¹, respectively (Figs 5 & 6). The relatively higher nutrient concentrations and photosynthetic abilities, as described above, in the western South Sea than in the East Sea may be partially responsible for the high Chl-*a* concentration range.

The trends and magnitudes of the PP differed in each region. Among the five regions, the PP in the western South Sea was the highest, reaching a maximum value of 940 mg C m⁻² d⁻¹ (Fig. 3, Supplementary Fig. S1). Even if the exceptionally high PP in 2014–2016 was excluded, the PP in the western South Sea exceeded 150 mg C m⁻² d⁻¹ on average. The high range of Chl-*a* concentrations may be a reason for the high PP in this region. The second largest region in the PP was the West Sea, although its Chl-*a* concentration range was similar to that of the eastern South Sea. Overall, the nutrient concentrations in the West Sea were higher than those in the eastern South Sea, and these eutrophic environments may have led to the high photosynthetic rate and efficiency values in the region. Thus, the physiological characteristics of the plankton communities, such as high photosynthetic ability, may be responsible for the high PP in the West Sea and are fundamental to evaluating PP.

The ranges of PP in the West Sea and western and eastern South Seas in the present study were similar to previously reported PP values in the same areas (Table 7). However, the PP derived from this study in the East Sea (1,391 mg C m⁻² d⁻¹) was approximately one-third of the previously reported highest PP values in 1988 (4,574 mg C m⁻² d⁻¹) but was similar to the values reported in the same years as the study period. In the waters off Jeju Island, the maximum PP in this study was one-third of that in 1989 (Table 7). However, recent values of PP in the waters off Jeju Island were not available; it could not be said that the PP was reduced compared to the past.

Based on the daily production from our results, the maximum annual PP was 343 g C m⁻² y⁻¹ in the western South Sea in 2015 (Table 3, Fig. 7C), indicating that it is one of the most productive marine environments com-

pared to other regions. Previously, PP in various environments, including upwelling waters, continental shelves, and oceanic waters, was reported to be between 55 and 350 g C m⁻² y⁻¹, except for a few abnormal values of 521 and 858 g C m⁻² y⁻¹ in the Mediterranean Sea (Eppley et al. 1985, Kwak et al. 2013b, Tiselius et al. 2016, Salgado-Hernanz et al. 2022). Therefore, the western South Sea of Korea may have a considerable effect on the carbon cycle in the Korean coastal waters and further studies should focus more on the western South Sea of Korea. Moreover, as a global aspect, the maximum annual TPP in the Korean territorial waters (11,227 Gg C y⁻¹ = 0.011 Pg C y⁻¹) (Fig. 7A) accounted for approximately 0.03% of the global PP (42.5 Pg C y⁻¹) (Gregg et al. 2003), suggesting that they play an essential role in the global carbon cycle.

In the present study, PP was calculated with the assumption that the concentration of Chl-*a* linearly decreased from the surface to the bottom waters owing to a lack of data on the vertical profile of Chl-*a*. If the Chl-*a* concentrations exponentially decreased with increasing water depths, the PP calculated in the present study might have been overestimated. On the other hand, if a subsurface chlorophyll maximum layer existed, the calculated PP might have been underestimated. Acquisition of data on Chl-*a* at several depths may increase the accuracy of PP estimation.

In the Korean coastal regions, red tide events seem to have considerable effects on PP. Moreover, coastal marine environments, such as nutrients and water temperature, have changed in the study area, and thus potentially change PP in the future. Recent studies have revealed that increased SST and decreased nitrate concentrations decrease global PP (Gregg et al. 2003, Gregg and Rousseaux 2019). Therefore, the PP should be monitored to better understand how ongoing climate change affects the same and for further resource utilization and environmental sustainability.

ACKNOWLEDGEMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation (NRF) grants funded by the Ministry of Education (NRF-2021R1I1A3048458), by the Korean government (MSIT) (2022R1A5A1031361), and by the Ministry of Science and ICT (NRF- 2020M3F6A1110582; NRF-2021M3I6A1091272; NRF-2021R1A2C1093379) award to HJJ.

CONFLICTS OF INTEREST

The authors declare that they have no potential conflicts of interest.

SUPPLEMENTARY MATERIALS

Supplementary Table S1. Dataset used for the analysis in the present study (<https://www.e-algae.org>).

Supplementary Table S2. Photosynthetic parameters used in the present study (<https://www.e-algae.org>).

Supplementary Fig. S1. Long-term trend of primary production in each region (<https://www.e-algae.org>).

REFERENCES

- Cai, W. -J. 2011. Estuarine and coastal ocean carbon paradox: CO₂ sinks or sites of terrestrial carbon incineration? Annu. Rev. Mar. Sci. 3:123–145.
- Choi, J. K., Lee, E. H., Noh, J. H. & Huh, S. H. 1997. The study on the phytoplankton bloom and primary productivity in lake Shihwa and adjacent coastal areas. The Sea 2:78–86.
- Choi, J. K., Noh, J. H., Shin, K. S. & Hong, K. H. 1995. The early autumn distribution of chlorophyll-a and primary productivity in the Yellow Sea, 1992. The Yellow Sea 1:68–80.
- Chung, C. S., Shim, J. H., Park, Y. C. & Park, S. -G. 1989. Primary productivity and nitrogenous nutrient dynamics in the East Sea of Korea. J. Oceanol. Soc. Korea 24:52–61.
- Chung, C. -S. & Yang, D. -B. 1991. On the primary productivity in the southern sea of Korea. J. Oceanol. Soc. Korea 26:242–254.
- Chung, K. H. 1996. Deterministic factors of the marine basal production in the coastal waters of Korea. Ph.D. dissertation, Inha University, Incheon, Korea, 191 pp.
- Cloern, J. E. 1987. Turbidity as a control on phytoplankton biomass and productivity in estuaries. Cont. Shelf Res. 7:1367–1381.
- Côté, B. & Platt, T. 1983. Day-to-day variations in the spring-summer photosynthetic parameters of coastal marine phytoplankton. Limnol. Oceanogr. 28:320–344.
- Eom, S. H., Jeong, H. J., Ok, J. H., Park, S. A., Kang, H. C., You, J. H., Lee, S. Y., Yoo, Y. D., Lim, A. S. & Lee, M. J. 2021. Interactions between common heterotrophic protists and the dinoflagellate *Trixos furca*: implication on the long duration of its red tides in the South Sea of Korea in 2020. Algae 36:25–36.
- Eppley, R. W. 1972. Temperature and phytoplankton growth in the sea . Fish. Bull. 70:1063–1085.
- Eppley, R. W., Stewart, E., Abbott, M. R. & Heyman, U. 1985. Estimating ocean primary production from satellite chlorophyll: introduction to regional differences and statistics for the Southern California Bight. J. Plankton Res. 7:57–70.
- Falkowski, P. G., Koblitzek, M., Gorbunov, M. & Kolber, Z. 2004. Development and application of variable chlorophyll fluorescence techniques in marine ecosystems. In Papageorgiou, G. C. & Govindjee (Eds.) *Chlorophyll a Fluorescence: A Signature of Photosynthesis*. Springer, Dordrecht, pp. 757–778.
- Field, C. B., Behrenfeld, M. J., Randerson, J. T. & Falkowski, P. 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. Science 281:237–240.
- Gregg, W. W., Conkright, M. E., Ginoux, P., O'Reilly, J. E. & Casey, N. W. 2003. Ocean primary production and climate: global decadal changes. Geophys. Res. Lett. 30:1809.
- Gregg, W. W. & Rousseaux, C. S. 2019. Global ocean primary production trends in the modern ocean color satellite record (1998–2015). Environ. Res. Lett. 14:124011.
- Han, I. -S. & Lee, J. -S. 2020. Change the annual amplitude of sea surface temperature due to climate change in a recent decade around the Korean Peninsula. J. Korean Soc. Mar. Environ. Saf. 26:233–241.
- Han, M. -S., Jang, D. -H. & Yang, H. -S. 1998. The ecosystem of the southern coastal water of the East Sea, Korea, II. Primary productivity in and around cold water mass. J. Korean Soc. Oceanogr. 33:196–204.
- Holmes, R. W. 1970. The Secchi disk in turbid coastal waters. Limnol. Oceanogr. 15:688–694.
- Hwang, D. -W., Lee, Y. -W. & Kim, G. 2005. Large submarine groundwater discharge and benthic eutrophication in Bangdu Bay on volcanic Jeju Island, Korea. Limnol. Oceanogr. 50:1393–1403.
- Hyun, J. -H., Kim, D., Shin, C. -W., Noh, J. -H., Yang, E. -J., Mok, J. -S., Kim, S. -H., Kim, H. -C. & Yoo, S. 2009. Enhanced phytoplankton and bacterioplankton production coupled to coastal upwelling and an anticyclonic eddy in the Ulleung Basin, East Sea. Aquat. Microb. Ecol. 54:45–54.
- Jang, H. K., Kang, J. J., Lee, J. H., Kim, M., Ahn, S. H., Jeong, J. -Y., Yun, M. S., Han, I. -S. & Lee, S. H. 2018. Recent primary production and small phytoplankton contribution in the Yellow Sea during the summer in 2016. Ocean Sci. J. 53:509–519.
- Jang, H. -K., Youn, S. -H., Joo, H., Kim, Y., Kang, J. -J., Lee, D., Jo, N., Kim, K., Kim, M. -J., Kim, S. & Lee, S. -H. 2021. First

- concurrent measurement of primary production in the Yellow Sea, the South Sea of Korea, and the East/Japan Sea, 2018. *J. Mar. Sci. Eng.* 9:1237.
- Jeong, H. J., Kang, H. C., Lim, A. S., Jang, S. H., Lee, K., Lee, S. Y., Ok, J. H., You, J. H., Kim, J. H., Lee, K. H., Park, S. A., Eom, S. H., Yoo, Y. D. & Kim, K. Y. 2021. Feeding diverse prey as an excellent strategy of mixotrophic dinoflagellates for global dominance. *Sci. Adv.* 7:eabe4214.
- Jeong, H. J., Lim, A. S., Lee, K., Lee, M. J., Seong, K. A., Kang, N. S., Jang, S. H., Lee, K. H., Lee, S. Y., Kim, M. O., Kim, J. H., Kwon, J. E., Kang, C. H., Kim, J. S., Yih, W., Shin, K., Jang, P. K., Ryu, J. -H., Kim, S. Y., Park, J. Y. & Kim, K. Y. 2017. Ichthyotoxic *Cochlodinium polykrikoides* red tides offshore in the South Sea, Korea in 2014: I. Temporal variations in three-dimensional distributions of red-tide organisms and environmental factors. *Algae* 32:101–130.
- Jeong, W. -G. & Cho, S. -M. 2018. Estimation of primary production of the waters around rack oyster farm at Wando, Korea. *Fish. Aquat. Sci.* 21:9.
- Jeong, W. G., Cho, S. -M. & Lee, S. J. 2009. Studies for the sustainable management of oyster farms in Pukman Bay, Korea: estimate of primary production. *Fish. Aquat. Sci.* 12:111–117.
- Joo, H., Park, J. W., Son, S., Noh, J. -H., Jeong, J. -Y., Kwak, J. H., Saux-Picart, S., Choi, J. H., Kang, C. -K. & Lee, S. H. 2014. Long-term annual primary production in the Ulleung Basin as a biological hot spot in the East/Japan Sea. *J. Geophys. Res. Oceans* 119:3002–3011.
- Kang, J. J., Jang, H. K., Lim, J. -H., Lee, D., Lee, J. H., Bae, H., Lee, C. H., Kang, C. -K. & Lee, S. H. 2020. Characteristics of different size phytoplankton for primary production and biochemical compositions in the Western East/Japan Sea. *Front. Microbiol.* 11:560102.
- Kang, J. W., Moon, S. -R., Park, S. -J. & Lee, K. -H. 2009. Analyzing sea level rise and tide characteristics change driven by coastal construction at Mokpo Coastal Zone in Korea. *Ocean Eng.* 36:415–425.
- Kang, Y. S., Choi, J. K., Chung, K. H. & Park, Y. C. 1992. Primary productivity and assimilation number in the Kyonggi bay and the mid-eastern coast of Yellow Sea. *J. Oceanol. Soc. Korea* 27:237–246.
- Kim, D., Lee, Y. -J., Kang, H. Y., Kwon, K. -Y., Lee, W. -C. & Kwak, J. H. 2019. Seasonal variations in primary productivity and biomass of phytoplankton in Geoje-Hansan Bay on the southern coast of Korea. *Ocean Sci. J.* 54:213–227.
- Kim, J. & Kim, T. -H. 2017. Distributions and sources of dissolved organic matter in seawaters surrounding aqua farms on the Haengwon-ri in Jeju-island in summer 2015. *Ocean Polar Res.* 39:35–43.
- Kim, J. B., Lee, W. -C., Kim, H. C. & Hong, S. 2016. An assessment of primary productivity determined by stable isotopes and diving-PAM in the *Pyropia* sea farms of the Manho (Jindo-Haenam) region on the Southwestern coast of the Korean peninsula. *Korean J. Environ. Biol.* 34:18–29.
- Kim, T. -W., Lee, K., Lee, C. -K., Jeong, H. -D., Suh, Y. -S., Lim, W. -A., Kim, K. Y. & Jeong, H. -J. 2013. Interannual nutrient dynamics in Korean coastal waters. *Harmful Algae* 30(Suppl 1):S15–S27.
- Ko, E. H. 2014. Comparison of chlorophyll fluorescence method and ¹⁴C method to estimate primary production in Korean waters. M.S. thesis, University of Science and Technology, Daejeon, Korea, 75 pp.
- Ko, E., Park, J., Gorbunov, M. Y. & Yoo, S. 2019. Uncertainties in variable fluorescence and ¹⁴C methods to estimate primary production: a case study in the coastal waters off the Korean peninsula. *Mar. Ecol. Prog. Ser.* 627:13–31.
- Koh, H. -J., Park, S. -E., Cha, H. -K., Chang, D. -S. & Koo, J. -H. 2013. Coastal eutrophication caused by effluent from aquaculture ponds in Jeju. *J. Korean Soc. Mar. Environ. Saf.* 19:315–326.
- KOSIS. 2022. Korean Statistical Information Service Web. Available from: <http://kosis.kr>. Accessed Aug 4, 2022.
- Kwak, J. H., Hwang, J., Choy, E. J., Park, H. J., Kang, D. -J., Lee, T., Chang, K. -I., Kim, K. -R. & Kang, C. -K. 2013a. High primary productivity and f-ratio in summer in the Ulleung basin of the East/Japan Sea. *Deep-Sea Res Part I Oceanogr. Res. Pap.* 79:74–85.
- Kwak, J. H., Lee, S. H., Park, H. J., Choy, E. J., Jeong, H. D., Kim, K. R. & Kang, C. K. 2013b. Monthly measured primary and new productivities in the Ulleung Basin as a biological “hot spot” in the East/Japan Sea. *Biogeosciences* 10:4405–4417.
- Kwon, H. K., Yang, H. S., Yoon, Y. H., Choi, O. I., Choi, I. H. & Oh, S. J. 2015. Characteristics of marine environment and primary productivity of phytoplankton in the seaweed bed of Northwestern coast of Jeju island during autumn 2014. *The Sea* 20:180–191.
- Lee, E. -Y. & Park, K. -A. 2020. Validation of satellite sea surface temperatures and long-term trends in Korean coastal regions over past decades (1982–2018). *Remote Sens.* 12:3742.
- Lee, J. -B., Jwa, J. -H., Go, Y. -B. & Choe, Y. -C. 1993. Bioecological studies of the eastern coastal area in Cheju Island (II) phytoplankton dynamics and primary productivity around U-do. *J. Korean Earth Sci. Soc.* 14:458–466.
- Lee, S. H., Joo, H., Lee, J. H., Lee, J. H., Kang, J. J., Lee, H. W., Lee, D. & Kang, C. K. 2017. Seasonal carbon uptake rates

- of phytoplankton in the northern East/Japan Sea. Deep-Sea Res. Part II Top. Stud. Oceanogr. 143:45–53.
- Lee, S. Y., Jeong, H. J., Kang, H. C., Ok, J. H., You, J. H., Park, S. A. & Eom, S. H. 2021. Comparison of the spatial-temporal distributions of the heterotrophic dinoflagellates *Gyrodinium dominans*, *G. jinhaense*, and *G. moestrupii* in Korean coastal waters. Algae 36:37–50.
- Lee, W. J. & Choi, J. K. 2000. The role of heterotrophic protists in the planktonic community of Kyeonggi Bay, Korea. J. Kor. Soc. Oceanogr. 35:46–55.
- Lee, Y. -J., Kim, M. -S., Won, E. -J. & Shin, K. H. 2006. An application of ¹³C tracer for the determination of size fractionated primary productivity in upper stream of Lake Shihwa. Korean J. Limnol. 39:93–99.
- Lee, Y. -J., Min, J. -O., Shin, Y., Kim, S. -H. & Shin, K. -H. 2011. Temporal and spatial variations of primary productivity in estuary of Youngsan River and Mokpo coastal areas. Korean J. Limnol. 44:327–336.
- Luo, Z., Wang, N., Mohamed, H. F., Liang, Y., Pei, L., Huang, S. & Gu, H. 2021. *Amphidinium stirisquamatum* sp. nov. (Dinophyceae), a new marine sand-dwelling dinoflagellate with a novel type of body scale. Algae 36:241–261.
- Ministry of Oceans and Fisheries and Korea Marine Environment Management Corporation. 2022. *Annual report on marine environment monitoring in Korea 2021*. Korea Marine Environment Management Corporation, Seoul.
- Min, J. O., Ha, S. Y., Choi, B. H., Chung, M. H., Yoon, W. D., Lee, J. S. & Shin, K. H. 2011. Primary productivity and pigments variation of phytoplankton in the Seomjin River estuary during rainy season in summer. Korean J. Limnol. 44:303–313.
- Min, J. O., Ha, S. Y., Chung, M. H., Choi, B. H., Lee, Y. J., Youn, S. H., Yoon, W. D., Lee, J. S. & Shin, K. H. 2012. Seasonal variation of primary productivity and pigment of phytoplankton community structure in the Seomjin estuary. Korean J. Limnol. 45:139–149.
- Moon, P. 1940. Proposed standard solar-radiation curves for engineering use. J. Frankl. Inst. 230:583–617.
- Morel, A. & Berthon, J. -F. 1989. Surface pigments, algal biomass profiles, and potential production of the euphotic layer: relationships reinvestigated in view of remote-sensing applications. Limnol. Oceanogr. 34:1545–1562.
- Morel, A. & Smith, R. C. 1974. Relation between total quanta and total energy for aquatic photosynthesis. Limnol. Oceanogr. 19:591–600.
- Muller-Karger, F. E., Varela, R., Thunell, R., Luerssen, R., Hu, C. & Walsh, J. J. 2005. The importance of continental margins in the global carbon cycle. Geophys. Res. Lett. 32:L01602.
- National Institute of Fisheries Science. 2020. *Harmful algal blooms in the Korean coastal waters in 2019*. National Institute of Fisheries Science, Busan, 135 pp.
- Nielsen, E. S. 1975. *Marine photosynthesis with special emphasis on the ecological aspects*. Elsevier Scientific Publishing Company, Amsterdam, 141 pp.
- Oh, Y. H., Kim, Y., Park, S. R., Lee, T., Son, Y. B., Park, S. -E., Lee, W. C., Im, D. -H. & Kim, T. -H. 2021. Spatiotemporal change in coastal waters caused by land-based fish farm wastewater-borne nutrients: results from Jeju Island, Korea. Mar. Pollut. Bull. 170:112632.
- Ok, J. H., Jeong, H. J., Kang, H. C., Park, S. A., Eom, S. H., You, J. H. & Lee, S. Y. 2021. Ecophysiology of the kleptoplastic dinoflagellate *Shimiella gracilenta*: I. spatiotemporal distribution in Korean coastal waters and growth and ingestion rates. Algae 36:263–283.
- Park, J. G. 1996. Photosynthetic properties and longterm variation of phytoplankton biomass related to environmental changes in the southern east sea of Korea. Ph.D. dissertation, Seoul National University, Seoul, Korea, 174 pp.
- Park, J. -G. & Kim, E. -K. 2010. Time integration algorithm for the estimation of daily primary production. The Sea 15:124–132.
- Park, J. G., Shim, J. H. & Lee, J. B. 1998. Long-term variation of phytoplankton biomass and implication in the East and the South Sea, Korea. Algae 13:123–133.
- Park, J. -S., Kang, C. -K. & An, K. -H. 1991. Community structure and spatial distribution of phytoplankton in the polar front region off the east coast of Korea in summer. Bull. Korean Fish. Soc. 24:237–247.
- Raven, J. A. & Falkowski, P. G. 1999. Oceanic sinks for atmospheric CO₂. Plant Cell Environ. 22:741–755.
- Salgado-Hernanz, P. M., Regaudie-De-Gioux, A., Antoine, D. & Basterretxea, G. 2022. Pelagic primary production in the coastal Mediterranean Sea: variability, trends, and contribution to basin-scale budgets. Biogeosciences 19:47–69.
- Shim, J. H. 2003. *Plankton ecology*. SNU Press, Seoul, 382 pp.
- Shim, J. H., Lee, W. H. & Park, S. -Y. 1985. Studies on the plankton in the southwestern waters of the East Sea (Sea of Japan) (II): phytoplankton - standing crop, nanofraction, and primary production. J. Oceanol. Soc. Korea 20:37–54.
- Shim, J. H. & Park, Y. C. 1986. Primary productivity measurement using carbon-14 and nitrogenous nutrient dynamics in the southeastern sea of Korea. J. Oceanol. Soc. Korea 21:13–24.
- Shim, J. H., Yeo, H. G. & Park, J. G. 1992. Primary production system in the southern waters of the East Sea, Korea. I. Biomass and productivity. J. Oceanol. Soc. Korea 27:91–

- 100.
- Smyth, T. J., Pemberton, K. L., Aiken, J. & Geider, R. J. 2004. A methodology to determine primary production and phytoplankton photosynthetic parameters from Fast Repetition Rate Fluorometry. *J. Plankton Res.* 26:1337-1350.
- Son, S., Campbell, J., Dowell, M., Yoo, S. & Noh, J. 2005. Primary production in the Yellow Sea determined by ocean color remote sensing. *Mar. Ecol. Prog. Ser.* 303:91-103.
- Spector, M. & Edwards, M. S. 2020. Species-specific biomass drives macroalgal benthic primary production on temperate rocky reefs. *Algae* 35:237-252.
- Tiselius, P., Belgrano, A., Andersson, L. & Lindahl, O. 2016. Primary productivity in a coastal ecosystem: a trophic perspective on a long-term time series. *J. Plankton Res.* 38:1092-1102.
- Turner, J. T. 2004. The importance of small planktonic copepods and their roles in pelagic marine food webs. *Zool. Stud.* 43:255-266.
- Valiela, I. 1984. *Marine ecological processes*. Springer Science + Business Media, LLC, New York, 547 pp.
- Yeo, H. -G., Shim, J. -H. & Park, J. -G. 1996. Variation of abiotic environmental factors and primary production in the pelagic ecosystem of Kori. *Korea J. Environ. Biol.* 14:113-120.
- Yoo, M. H., Youn, S. H., Oh, H. J. & Choi, J. K. 2018. Reaction characteristics of phytoplankton before and after the yellow dust event in Taean peninsula and yellow dust impact assessment. *J. Korean Soc. Mar. Environ. Saf.* 24:898-906.
- Yoo, S. & Shin, K. 1995. Primary productivity properties in the nearshore region of Taean Peninsula. *Ocean Res.* 17:91-99.
- Yoon, J. -E., Park, J. & Yoo, S. 2012. Comparison of primary productivity algorithms for Korean waters. *Ocean Sci. J.* 47:473-487.
- Yoshikawa, T. & Furuya, K. 2008. Phytoplankton photosynthetic parameters and primary production in Japan Sea and the East China Sea: Toward improving primary production models. *Cont. Shelf Res.* 28:962-976.