



Review

Algae 2023, 38(4): 203-215

<https://doi.org/10.4490/algae.2023.38.12.13>

Open Access



Oomycete pathogens, red algal defense mechanisms and control measures

Xianying Wen¹, Giuseppe C. Zuccarello², Tatyana A. Klochkova³ and Gwang Hoon Kim^{1,*}

¹Department of Biological Sciences, Kongju National University, Gongju 32588, Korea

²School of Biological Sciences, Victoria University of Wellington, Wellington 6140, New Zealand

³Department of Biology, Kamchatka State Technical University, Petropavlovsk-Kamchatsky 683003, Russia

Oomycete pathogens are one of the most serious threats to the rapidly growing global algae aquaculture industry but research into how they spread and how algae respond to infection is unresolved, let alone a proper classification of the pathogens. Even the taxonomy of the genera *Pythium* and *Olpidiopsis*, which contain the most economically damaging pathogens in red algal aquaculture, and are among the best studied, needs urgent clarification, as existing morphological classifications and molecular evidence are often inconsistent. Recent studies have reported a number of genes involved in defense responses against oomycete pathogens in red algae, including pattern-triggered immunity and effector-triggered immunity. Accumulating evidence also suggests that calcium-mediated reactive oxygen species signaling plays an important role in the response of red algae to oomycete pathogens. Current management strategies to control oomycete pathogens in aquaculture are based on the high resistance of red algae to abiotic stress, these have environmental consequences and are not fully effective. Here, we compile a revised list of oomycete pathogens known to infect marine red algae and outline the current taxonomic situation. We also review recent research on the molecular and cellular responses of red algae to oomycete infection that has only recently begun, and outline the methods currently used to control disease in the field.

Keywords: aquaculture; disease; oomycete; pathogen; red algae

Abbreviations: CDPK, calcium calmodulin-dependent protein kinase; DAMP, damage-associated molecular pattern; ETI, effector-triggered immunity; HR, hypersensitive response; LRR, leucine-rich repeat; PAMP, pathogen-associated molecular pattern; PRR, pattern recognition receptor; PTI, pattern-triggered immunity; ROS, reactive oxygen species; TIR, Toll/interleukin-1 receptor

INTRODUCTION

Algae are the most important primary producers, sequestering more than half of the world's net carbon and producing about 52 giga tons of organic carbon per year (Hallmann 2007, Ullmann and Grimm 2021). Among these, seaweeds, multicellular algae, are cultivated in many parts of the world, and in 2019 alone, 34.7 mil-

lion tons of seaweed were produced, valued at approximately US\$14.7 billion, and used to produce food, food additives (e.g., hydrocolloids), animal feed, nutritional supplement, cosmetics, and pigments (Borowitzka 2013, Cai et al. 2021, Ullmann and Grimm 2021). However, the sustainability of the seaweed aquaculture industry faces



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received November 12, 2023, Accepted December 13, 2023

*Corresponding Author

E-mail: ghkim@kongju.ac.kr

Tel: +82-41-850-8504, Fax: +82-41-850-8479

a number of challenges, including rising water temperatures and more frequent strong weather events, which can lead to damage to facilities and disease outbreaks (Cottier-Cook et al. 2021).

Disease outbreaks in aquaculture farms can be caused by pathogens introduced from outside the farms and conversely the farms can act as pathogen pools from which pathogens can spread to the surrounding environment (Tedesco et al. 2021, Roh and Kannimuthu 2023). In recent years, there has been a growing concern about the potential threat of pathogens spreading from aquaculture farms and affecting coastal seaweed populations (Bouwmeester et al. 2021), as well as the adverse effects that treatments used by aquaculture farms to control pathogens may have on the surrounding environment (Kim et al. 2014). Currently, a significant number of commercially important seaweeds belong to the red algae, and despite their much longer history of aquaculture compared to other algae, little is known about how pathogens interact with red algae to cause disease and which genes are involved in the red algal defense against pathogens (Im et al. 2019). As a result, the methods currently used to control disease in seaweed farms are not the result of systematic basic scientific research, but have been developed based on farmers' experience, and may have little effect and may even pose a risk to the surrounding environment (Kamoun et al. 2015).

In order to develop effective disease control measures, we need to better understand the cellular and molecular mechanisms of host-pathogen interactions in red algae and the scientific basis of currently used treatments (Fawke et al. 2015). In this article, we list the known pathogens that infect red algae, with a focus on oomycetes, and review recent research on the molecular mechanisms of host-pathogen interactions and disease control methods currently used in aquaculture.

OOMYCETE PATHOGENS THAT INFECT RED ALGAE

Oomycetes are a diverse group of fungus-like eukaryotic microorganisms, also known as water molds (Beakes et al. 2012). Because of their filamentous growth habit, nutrient supply by absorption, flagellate stages, and reproduction by spores, oomycetes were long considered “lower” fungi (Latijnhouwers et al. 2003). However, it is now clear that this group of organisms is not related to fungi but is more closely related to diatoms and brown algae in the clade Stramenopiles (Beakes and Thines 2017,

Adl et al. 2019).

Oomycetes are one of the most important aquatic pathogens, devastating important agricultural crops and natural ecosystems (Hardham 2007, Kamoun et al. 2015, Badis et al. 2019, 2020). Oomycetes are capable of infecting hosts ranging from algae, plants, and animals (Gachon et al. 2010, Saraiva et al. 2023). Several pathogenic oomycetes are responsible for massive destruction and losses in agriculture and aquaculture (Phillips et al. 2008, Kim et al. 2014). However, taxonomic studies of marine oomycetes began much later and are less well informed than those of other oomycetes. It was not until 1960s that Sparrow cataloged the oomycetes infecting red algae and classified the species based on morphology and life history (Sparrow 1960). He recorded a total of seven species infecting red algae: *Eurychasmidium tumefaciens*, *E. sacculum* (formerly known as *Eurychasma sacculus*), *Sirolpidium andreei* (formerly known as *Olpidiopsis andreei*), *Petersenia lobata*, *P. pollagaster*, *Pontisma lagenidioides*, and *Pythium marinum*. Only two of these species, *Pyt. marinum* and *Po. lagenidioides*, have been taxonomically studied using molecular methods (LeVesque and De Cock 2004, Hyde et al. 2014, Buaya et al. 2019).

Molecular data on *Pyt. marinum* were first reported by LeVesque and De Cock (2004). *Pyt. marinum* is morphologically distinguished from other *Pythium* species by a single linear antheridium that tapers abruptly (Sparrow 1934, 1960). The internal transcribed spacers (1 and 2) and the 5.8S gene of *Pyt. marinum* (strain number: CBS 750.96) were identical to *Pyt. coloratum*, *Pyt. lutarium*, and *Pyt. dissotocum* but differed by 1 bp from *Pyt. diclinum* (LeVesque and De Cock 2004). In further studies, the large and small subunits of nrRNA, the cytochrome c oxidase subunit 2 gene (*cox2*), and β -tubulin were also used in the same strain of *P. marinum* (Hyde et al. 2014). *Pyt. marinum* showed a little difference from other *Pythium* species in a phylogenetic analysis based on these sequences (Hyde et al. 2014). Questions have been raised as to whether this strain is the *Pyt. marinum* reported by Sparrow (1934), as the strain used in the molecular phylogenetic study was isolated from soil rather than a marine environment where the species was originally reported (Hyde et al. 2014), but where the strain was isolated may not be a major issue as there are reports that *Pyt. porphyrae*, a member of the same genus, can survive in both freshwater and seawater and can infect both terrestrial plants and marine red algae (Klochkova et al. 2017a). Recently, a *Pythium* sp. infecting the green algae *Ulva* spp. and terrestrial grass species in low salinity environments, was discovered in the natural environment in Norway,

supporting the above ideas that terrestrial oomycete pathogens can be introduced into intertidal algal communities or *vice versa* (Herrero et al. 2020).

After Sparrow (1960), fourteen new species or varieties of oomycetes infecting red algae were discovered. Two species were classified as new only morphologically (i.e., *Olpidiopsis antithamnionis*, *P. palmariae*) (Whittick and South 1972, Van der Meer and Pueschel 1985), and nine species were morphologically and molecularly identified (i.e., *Olpidiopsis porphyrae*, Sekimoto et al. 2008; *O. feldmanni*, Aleem 1952, Fletcher et al. 2015; *O. bostrychia*, Sekimoto et al. 2009; *O. pyropiae*, Klochkova et al. 2016; *O. heterosiphoniae*, Klochkova et al. 2017b; *O. palmariae*, Badis et al. 2019; *O. muelleri*, Badis et al. 2019; *Pythium chondricola*, Lee et al. 2015, 2017, Lee and Lee 2022; and *Pyt. porphyrae*, Takahashi et al. 1977, Diehl et al. 2017, Qiu et al. 2019). Three varieties were also identified (*O. muelleri* var. *polysiphoniae*, Badis et al. 2019; *O. porphyrae* var. *scotiae*, Badis et al. 2019; and *O. porphyrae* var. *koreanae*, Kwak et al. 2017). These varieties are not distinguished by the 18S rRNA sequence but are distinguished from other varieties morphologically and in their host infection characteristics. For example, *O. muelleri* var. *polysiphoniae* forms a sporangium cell wall with a honeycomb structure that is different from that of *O. muelleri* (Badis et al. 2019). The 18S rRNA sequences of *O. porphyrae* var. *porphyrae* and *O. porphyrae* var. *koreanae* were identical but differs in their conserved introns, while *O. porphyrae* var. *scotiae* differs in both conserved introns and 18S rRNA sequences (Badis et al. 2019).

Accumulating molecular phylogenetic information has led to a need to reclassify marine oomycetes, particularly the genus *Olpidiopsis*. An oomycete believed to be *Po. lagenidioides* was collected from the red algae *Ceramium rubrum* in Drøbak, Oslo Fjord, Norway in 2017 (Buaya et al. 2019) and was originally recorded as infecting *Ceramium* spp. (Sparrow 1960). Morphologically similar to *Po. lagenidioides* recorded by Sparrow (1960), 18S rDNA sequences were analyzed to construct molecular phylogenies (Buaya et al. 2019). The phylogeny results showed *Po. lagenidioides* was within rhodophyte-infecting members of *Olpidiopsis*, forming a clade but without support (Buaya et al. 2019). Within the revised genus *Pontisma*, only *Po. lagenidioides* is known to form large hyphal thalli, but the thallus segments in that species develop in a very similar manner as compared to the species in which the thallus remains smaller (Buaya et al. 2019, 2021). *Olpidiopsis saprolegeniae*, infecting another oomycete, was distantly related to the red algal pathogens (Buaya et al. 2019), and the genus *Pontisma* was proposed for these

pathogens (Buaya et al. 2019, Buaya and Thines 2020). *Po. lagenidioides* grouped together with other the red algal pathogens (i.e., *O. feldmanni*, *O. heterosiphoniae*, *O. pyropiae*, *O. porphyrae*, and *O. porphyrae* var. *koreanae*), but this relationship was not supported by 18S rDNA sequences (Buaya et al. 2019, 2021). More work is needed before any taxonomic changes are made.

In some cases, oomycetes that have been considered distinct species are not well supported by molecular phylogenetic evidence. For example, *Pyt. porphyrae* is the first oomycete pathogen identified in seaweed farms and is the causative agent of red rot disease, one of the most common and severe diseases in *Pyropia* farms in China, Korea, and Japan (Arasaki 1947, Takahashi et al. 1977, Park et al. 2001, Ding and Ma 2005) (Fig. 1A & D). The morphology of *Pyt. porphyrae* and *Pyt. chondricola* is very similar and there is controversy about their separation into distinct species (Park et al. 2000, Diehl et al. 2017). *Pyt. porphyrae* and *Pyt. chondricola* both have filamentous, non-inflated sporangia, similar sexual structures, and both infect *Pyropia* (LeVesque and De Cock 2004, Lee et al. 2015). The main differences are the considerably lower cardinal temperatures for growth, larger oogonia, and aplerotic oospores in *Pyt. chondricola* (LeVesque and De Cock 2004). Sequences of the internal transcribed spacer region, currently used in oomycetes taxonomy, are 100% identical (LeVesque and De Cock 2004). The 18S rRNA sequence identified in the recently discovered genome of *Pyt. chondricola* was 100% identical to that of *Pyt. porphyrae* (Nguyen et al. 2022). Although there are differences in the *cox1* sequence, it is smaller than the regional differences between *Pyt. porphyrae* from Japan and Korea, so this is ultimately considered insufficient to distinguish between species using *cox1* (Lee et al. 2015, 2017, Lee and Lee 2022). In the event of a major outbreak of red rot disease in aquaculture farms, the names of these two species are sometimes used interchangeably, leading to confusion about the need for different treatments for the two pathogens, so it is necessary to clarify their taxonomic status.

Olpidiopsis is a genus of obligate holocarpic endobiotic oomycetes (Buaya et al. 2019), with most species reported to be parasitic within red algal cells, and several of which cause significant economic damage to seaweed farms in Far Eastern countries (Kim et al. 2014). To date, seven non-oosporic *Olpidiopsis* species have been described as parasites of marine red algae both morphologically and phylogenetically (Table 1). The morphological taxonomy of *Olpidiopsis* species is based upon a few morphological features such as shape and size of the thallus, the num-




Fig. 1. Oomycete pathogens that infect red algae. (A) Gametophyte of *Pyropia yezoensis* infected with *Pythium porphyrae* (red arrow) and *Olpidopsis porphyrae* var. *koreanae* (green arrow). (B) Zoosporangia of *Olpidopsis heterosiphoniae* (black arrows) developing in the axial cells of *Dasysiphonia japonica*. (C) Zoospores (black arrow) release through a discharge tube in *Olpidopsis bostrychiae* infecting *Bostrychia moritziana*. (D) Mycelium (black arrow) of *P. porphyrae* growing on the gametophyte of *P. yezoensis*. (E) Zoosporangia of *O. pyropiae* (black arrow) developing in *P. yezoensis* cells. (F) Zoosporangium of *O. bostrychiae* (black arrow) inside a sporophyte (conchocelis-stage) cell of *Porphyra pulchella*. (G) Zoosporangium of *O. heterosiphoniae* (black arrow) in the apical cell of *D. japonica*. Scale bars represent: A, 1 cm; B & D, 50 µm; C & E–G, 20 µm.

Table 1. Pathogenic species of oomycetes on red algae

Species	Host algae	Geographic distribution	References
<i>Eurychasmodium tumefaciens</i> (Magnus) Sparrow, 1936	<i>Ceramium</i> spp.	British	Magnus (1872), Sparrow (1960)
<i>Eurychasmodium sacculum</i> (H. E. Petersen) M. W. Dick, 2001 (previous name is <i>Eurychasma sacculus</i>)	<i>Devaleraea ramentacea</i> , <i>Palmaria palmata</i>	Greenland	Petersen (1905), Sparrow (1960)
<i>Sirolopidium andreei</i> M. W. Dick, 2001 (previous name is <i>Olpidiopsis andreei</i>)	<i>Ceramium diaphanum</i> (additional brown algae host: <i>Ectocarpus</i> spp., <i>Striaria attenuate</i> ; additional green algae host: <i>Acrosiphonia incurva</i> , <i>A. hystrix</i> , <i>Acrosiphonia</i> spp., <i>Spongomorpha</i> sp.)	Europe	Sparrow (1960)
<i>Petersenia lobata</i> (Petersen) Sparrow, 1934 (previous name is <i>Pleotrachelus lobatus</i>)	<i>Ceramium</i> spp., <i>Callithamnion</i> spp., <i>Spermothamnion</i> spp., <i>Seirospora</i> spp., <i>Aglaothamnion</i> sp., <i>Herposiphonia tenella</i> , <i>Polysiphonia urceolata</i> , <i>Pylaiella littoralis</i> (additional brown algae host: <i>Ectocarpus siliculosus</i>)	France, Algeria	Lagerheim (1899), Sparrow (1960)
<i>Petersenia pollagaster</i> (Petersen) Sparrow, 1943	<i>Ceramium</i> spp., <i>Chondrus crispus</i>	Denmark, Canada	Petersen (1905), Sparrow (1960), Craigie and Shacklock (1995)
<i>Pontisma lagenidioides</i> Petersen, 1905	<i>Ceramium</i> spp. (additional green algae host: <i>Chaetomorpha media</i>)	America, Denmark, Sweden, India	Petersen (1905), Sparrow (1960)
<i>Pythium marinum</i> Sparrow, 1934	<i>Ceramium rubrum</i> , <i>Porphyra</i>	Denmark, Pacific and Atlantic coasts of North America	Sparrow (1934), Klochkova et al. (2017a)
<i>Petersenia palmariae</i> Pueschel & Vandermeer, 1985	<i>Palmaria mollis</i>	Canada	Van der Meer and Pueschel (1985)
<i>Olpidiopsis antithamnionis</i> Whittick & South, 1972	<i>Antithamnion floccosum</i>	Canada	Whittick and South (1972)
<i>Olpidiopsis porphyrae</i> Sekimoto et al., 2008	<i>Pyropia</i> spp., <i>Bangia</i> spp.	Japan	Sekimoto et al. (2008)
<i>Olpidiopsis feldmanni</i> Aleem, 1952	<i>Asparagopsis</i> sp.	France	Aleem (1952), Fletcher et al. (2015)
<i>Olpidiopsis bostrychia</i> Sekimoto et al., 2009	<i>Bostrychia</i> spp., <i>Dasyisiphonia japonica</i> , <i>Pyropia</i> spp., <i>Porphyra</i> , <i>Stictosiphonia intricata</i>	Madagascar	Sekimoto et al. (2009), Klochkova et al. (2017b)
<i>Olpidiopsis porphyrae</i> var. <i>koreanae</i> G. H. Kim & T. A. Klochkova, 2017	<i>Bangia</i> sp., <i>Pyropia</i> spp.	Korea	Kwak et al. (2017)
<i>Olpidiopsis pyropiae</i> G. H. Kim & T. A. Klochkova, 2016	<i>Pyropia</i> spp.	Korea	Klochkova et al. (2016)
<i>Olpidiopsis heterosiphoniae</i> G. H. Kim & T. A. Klochkova, 2017	<i>Heterosiphonia</i> spp., <i>Dasya</i> sp., <i>Dasyisiphonia chejuensis</i> , <i>Pyropia tenera</i>	Korea	Klochkova et al. (2017b)
<i>Olpidiopsis porphyrae</i> var. <i>scotiae</i> Y. Badis et al., 2019	<i>Porphyra</i> sp., <i>Pyropia</i> spp.	UK	Badis et al. (2019)
<i>Olpidiopsis palmariae</i> Y. Badis & C. M. M. Gachon, 2019	<i>Palmaria palmata</i> (additional brown algae host: <i>Ectocarpus</i> sp.)	UK	Badis et al. (2019)
<i>Olpidiopsis muelleri</i> Y. Badis & C. M. M. Gachon, 2019	<i>Porphyra</i> sp.	UK	Badis et al. (2019)
<i>Olpidiopsis muelleri</i> var. <i>polysiphoniae</i> Y. Badis & C. M. M. Gachon, 2019	<i>Polysiphonia</i> sp.	UK	Badis et al. (2019)
<i>Pythium chondricola</i> De Cock 1986	<i>Chondrus crispus</i> , <i>Pyropia</i> spp.	Netherlands, Korea	De Cock (1986), Lee et al. (2015)
<i>Pythium porphyrae</i> M. Takah & M. Sasaki 1977	A total of 16 genera and 21 species, including <i>Porphyra</i> spp., <i>Pyropia</i> spp., <i>Gracilaria</i> spp.	Korea, USA, Japan, New Zealand	Takahashi et al. (1977), Jiang et al. (2012), Diehl et al. (2017)

ber of discharge tubes, plus host specificity (Klochkova et al. 2012). *O. porphyrae* var. *porphyrae* (Sekimoto et al. 2008) and *O. porphyrae* var. *koreanae* (Fig. 1A) (Kwak et al. 2017) infects species of the genus *Pyropia* and *Bangia*, developing spherical-shaped holocarpic thalli within the cytoplasm of its host (Fig. 1E), and produces monoplanetic, subapically-inserted biflagellate zoospores. The released spores immediately infect neighboring cells, and the disease spreads very quickly (Kim et al. 2014, Klochkova et al. 2016, Kwak et al. 2017), and both varieties are often found together on the same *Pyropia* blade (Fig. 1A & E). *O. porphyrae* var. *scotiae* reported from Scotland infects species of *Porphyra* (Badis et al. 2019). *Olpidiopsis* species also infect Ceramialean algae such as *Bostrychia moritziana* (Fig. 1C), *Heterosiphonia pulchra*, and *Dasy-siphonia japonica* (Fig. 1B). Species of *Olpidiopsis*, which develop zoosporangia inside the cells of their hosts, are usual highly host-specific, but can also have multiple hosts, differing only in the rate at which the infection spreads (Klochkova et al. 2012). *Olpidiopsis bostrychiae* can infect a variety of red algal genera (i.e., *Bostrychia* spp., *Dasy-siphonia japonica*, *Pyropia* spp., and *Porphyra* sp.) and grows best on the genus from which was isolated, *Bostrychia* (Sekimoto et al. 2009). *Olpidiopsis heterosiphoniae* can also infect a variety of red algal genera (i.e., *Heterosiphonia* spp., *Dasya* sp., *Dasy-siphonia chejuensis*, and *Pyropia tenera*), but can coexist for long periods of time without completely consuming the host (Klochkova et al. 2017b).

RED ALGAL DEFENSE AGAINST OOMYCETE PATHOGEN

In response to pathogen infection, hosts develop diverse immune responses. An immune response are the molecular steps that organisms use to defend themselves against foreign invaders (Coll et al. 2011). Disease develops when the pathogen is not able to evade these host defenses. Plants and animals, share striking similarities in their innate immune systems, some of which are likely due to convergent evolution (Jones and Dangl 2006, Mayor et al. 2007). Immune responses discriminate self from non-self and activate tightly regulated pre- and post-invasion defense responses to minimize the damage caused by harmful agents (Chakraborty et al. 2017). In both plants and animals, the first line of defense is provided by pattern recognition receptors (PRRs) on the plasma membrane that recognize conserved pathogen-associated molecular patterns (PAMPs) or microbe-asso-

ciated molecular patterns that activate pattern-triggered immunity (PTI) (Mayor et al. 2007, Shine et al. 2019, Ding et al. 2022). After infection, plants recognize effectors secreted by pathogens via resistance (R) proteins and induce effector-triggered immunity (ETI) (Coll et al. 2011, Chakraborty et al. 2017).

Plants have PRRs that can recognize molecular signatures that identify whole classes of microbes (such as chitin for fungi or peptidoglycan for bacteria) but are absent from the host (Boller and Felix 2009). The best-studied class of plant PRRs are receptor-like kinases, which have an ectodomain of leucine-rich repeats (LRRs). Toll-like receptors are one of the earliest PRRs discovered in animals (Iwasaki and Medzhitov 2010). Since then, many PRRs and their corresponding ligands have been discovered. Recognition of PAMPs by PRRs at the onset of infection activates PTI, which involves Ca^{2+} signaling, mitogen-activated protein kinase cascades, and reactive oxygen species (ROS) signaling, and can non-specifically control colonization of most pathogens (Ding et al. 2022). However, various pathogens have evolved “effector” proteins that can suppress PTI signaling leading to effector-triggered susceptibility (Bari and Jones 2009). These effector proteins are known as “virulence factors” (Chakraborty et al. 2017).

As genomic and transcriptomic research progresses in red algae, specific pathogen receptor genes homologous to those of other systems have been discovered (Brawley et al. 2017, de Oliveira et al. 2017, Tang et al. 2019). Plant-typical PRR genes (LRR domain containing genes) were identified in *Laurencia dendroidea* (de Oliveira et al. 2017), *Pyropia yezoensis* (Tang et al. 2019), and *P. tenera* (Im et al. 2019). Animal typical PRR genes (C-type lectin domains containing genes) have been found in *Porphyra umbilicalis* (Brawley et al. 2017), *P. yezoensis* (Tang et al. 2019), and *P. tenera* (Im et al. 2019). However, typical plant R proteins with a conserved protein structure of TNL (Toll/interleukin-1 receptor [TIR]-NBS-LRR or nTNL [NBS-LRR]) was not identified in red algae (Ortiz and Dodds 2018). R domain containing proteins were also identified in *Porphyra umbilicalis* (Brawley et al. 2017), *P. yezoensis* (Tang et al. 2019), and *P. tenera* (Im et al. 2019). It has been suggested that HSP90 proteins in *P. yezoensis* may play a role similar to R proteins in plants (van Bentem et al. 2005, Tang et al. 2019). The defense mechanisms of red algae against oomycete pathogens identified to date are summarised in Fig. 2.

Pathogen receptor genes in red algae are expressed differently between species and depending on the type of pathogen. A transcriptomic study of *P. yezoensis* found




Fig. 2. Schematic diagram of the defense mechanism in red algae. Pattern-triggered immunity (PTI) is triggered by pattern recognition receptors (PRRs). Receptor-like kinases are PRRs characterized by an extracellular domain (leucine-rich repeat [LRR] or C-type lectin), a single-pass transmembrane (TM) domain, and an intracellular kinase domain. Downstream signaling components include RBOH, Ca²⁺ permeable channels, and mitogen-activated protein (MAP) kinases. An increase in intracellular Ca²⁺ concentration triggers phosphorylation of RBOH by calcium calmodulin-dependent protein kinases (CDPKs), resulting in the production of reactive oxygen species. Oomycete pathogens deploy effectors to inhibit the PTI response. When host cells recognize pathogen effectors using putative resistance (R) proteins, effector-induced immunity (ETI) signaling is activated, resulting in structural changes that ultimately trigger a hypersensitivity reaction (HR) or other defense response. PAMP, pathogen-associated molecular pattern; DAMP, damage-associated molecular pattern.

that PRR genes (genes containing C-type lectin domains) and five genes encoding a single R proteins domain (LRR, NBS, or TIR) were upregulated after infection by *Pyt. porphyrae* (Tang et al. 2019). Microarray studies of *P. tenera* in response to infection by two oomycete pathogens (*Pyt. porphyrae*, *Olpidiopsis pyropiae*) and a viral pathogen (PyroV1) show that PRR genes are specifically upregulated in response to oomycete pathogens (Im et al. 2019) and not the virus. For example, in *P. tenera*, a PRR gene was upregulated by infection with *Pyt. porphyrae*, and a gene with an LRR domain was slightly upregulated by infection with *P. porphyrae* and *O. pyropiae*. In addition, two genes encoding NBS domain-containing proteins were upregulated upon infection with these oomycete pathogens, but none of these genes was upregulated upon infection with

PyroV1. A gene encoding a TIR domain-containing protein was upregulated only upon infection with PyroV1 (Im et al. 2019).

A set of pathogen receptor genes, as found in land plants, have not yet been reported in red algae, but recent studies have reported a number of genes involved in defense responses, including PTI and ETI. In *Laurencia dendroidea*, genes involved in PTI, such as mitogen-activated protein kinase kinase, calcium calmodulin-dependent protein kinase, NADPH oxidase, and antioxidant enzyme genes, were upregulated in a time-dependent manner after inoculation with the marine bacterium *Vibrio madracius* (de Oliveira et al. 2017). In *P. tenera*, several defense-related genes typically considered PTIs in plants, such as heat shock proteins, cell wall-associated hydrolase, and

NADPH oxidase, were induced in response to pathogen infection (Im et al. 2019). Similar defense mechanisms have been reported in *P. yezoensis*, including protective enzymes, heat shock proteins, secondary metabolites, cellulases, protease inhibitors, NADPH-oxidases, and antioxidant enzymes, and these genes were upregulated from the onset of infection by *Pyt. porphyrae* (Tang et al. 2019). It has also been reported that the hypersensitive response (HR)-related genes such as metacaspase, endonuclease G, and cytochrome C, which are upregulated during ETI, are also induced by infection with oomycete pathogens (Tang et al. 2019).

Calcium-mediated ROS signaling mediates the fertilization and wound-healing responses of red algae and also plays an important role in response to pathogens (Weinberger et al. 2005, Moon et al. 2022, Shim et al. 2022). NADPH oxidase in the cell membrane (called respiratory burst oxidase homolog or RBOH) reduces oxygen to superoxide, which rapidly forms hydrogen peroxide and diffuses from cell to cell through the extracellular space (Torres et al. 2006, Perez and Brown 2014, Peláez-Vico et al. 2022). ROS along with NO (nitric oxide) induces HR-mediated cell death in plants (Swarupa et al. 2014). ROS produced by pathogen-exposed cells trigger ROS / calcium-activated calcium channels in neighboring cells, resulting in auto-propagation throughout the plant, activating systemic acquired adaptation with increased HR reaction (Choudhury et al. 2017, Ding et al. 2022). Accumulation of ROS was also observed in red algae in response to pathogen infection. When the red alga *Gracilaria conferta* was exposed to oligosaccharides produced by microbial degradation of the agar cell wall, an oxidative burst was observed in the cell membrane as the first physiological sign of a PAMP (Weinberger 1999, Weinberger et al. 2005). ROS accumulation was also observed in *P. tenera* in response to oomycete as well as virus infection (Im et al. 2019). However, more research is needed to understand when and how ROS signaling is involved in red algal responses to infection by oomycete pathogens.

CURRENT TREATMENT MEASURES FOR OOMYCETE DISEASES

Control methods for oomycete diseases were first developed by seaweed farmers in the field, particularly in *Pyropia* farming, and developed to industrial levels in Asia after the 1950s. Most of the control methods currently used are based on *Pyropia*'s high tolerance to abiotic

stress (Kim et al. 2014). Sea farmers inhibit the growth of pathogens by exposing aquaculture nets to air for several hours and letting them dry in sunlight. Since the 1970s, a frozen-net method has been developed, in which young blades attached to aquaculture nets are air-dried, frozen at -20°C, and then reintroduced to farms when seawater is around 10°C (e.g., Fujita and Migita 1980, Ding and Ma 2005, and Klochkova et al. 2012). While the above methods are effective in reducing the damage caused by oomycete pathogens to some extent, it is difficult to avoid some reduction in production due to shorter cultivation periods or slower growth (Kim et al. 2023).

Acid washing of cultivation nets is the most commonly used method in *Pyropia* farms to reduce epiphytic green algae, diatoms, and oomycete pathogens (Park et al. 2001). A boat equipped with a large tub passes underneath the cultivation net and the net is immersed in an acid solution in the tub while the boat slowly moves forward. As *Pyropia* tolerates acid better than other organisms this treatment has been regarded as useful to clean the cultivation nets, but this method has little effect on *Olpidiopsis*-blight and is a burden on the environment (Kim et al. 2014). As regulations on aquaculture methods are strengthened, chemical treatments are currently limited to a few organic acids or concentrated salts that have been approved through government review and testing in the field (Kim et al. 2023). Recently, a technology has been developed that can more effectively control oomycete pathogens by washing aquaculture nets with calcium propionate instead of organic acids and is awaiting government review (Kim et al. 2023). Despite all these attempts, oomycete pathogens are still a major issue for *Pyropia* farms in Far Eastern countries and more research on their control is needed.

CONCLUSIONS AND PERSPECTIVES

Many questions still need to be addressed, or clarified, to improve our understanding of the pathogens of seaweed farms and how their impact can be mitigated. A better understanding of the diversity of pathogens found in the ocean and around seaweed farms is still needed as exploring the environmental DNA in seawater is still uncovering oomycetes related to known red algal pathogens (Badis et al. 2019).

Phylogenetic results show that rhodophyte-infecting members of *Olpidiopsis*, do not form a supported clade based on standard markers (Buaya et al. 2019). It is very likely that the genus *Olpidiopsis* currently contains spe-

cies that are phylogenetically divergent, and more information on host specificity or cell ultrastructure, together with the development of higher-resolution molecular markers, is needed to understand this diversity. *Pythium chondricola* and *Pyt. porphyrae* were originally named on the basis of host specificity, but current molecular markers make it very difficult to distinguish between them, leading to the two names being used interchangeably for pathogens found in *Pyropia* aquaculture in East Asia. Development of higher resolution molecular markers to discriminate between them using strains with host specificity consistent with the original description of the two species is required.

Although aspects of *Pyropia* spp. innate immunity were shown after infection with oomycetes, research on innate immunity in other red algae is needed. Several economic red algae species have been reported to be infected with oomycetes and warrant further study (Table 1). ROS are important molecules in innate immunity that are also found in red algae (Potin et al. 2002, Weinberger et al. 2005, Im et al. 2019, Tang et al. 2019). Current treatment measures for oomycete diseases mainly involve increasing resistance to abiotic stress. Since abiotic stress is also related to the production of ROS, it is expected that research on ROS signaling will be important in future red algal oomycete diseases.

ACKNOWLEDGEMENTS

This work was supported by the management of Marine Fishery Bio-resources Center (2023) funded by the National Marine Biodiversity Institute of Korea (MABIK) and the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2022R1A2C1091633) and the development of technology for biomaterialization of marine fisheries by-products of Korea institute of Marine Science & Technology Promotion (KIMST) funded by the Ministry of Oceans and Fisheries (KIMST-20220128). The sample of *Dasysiphonia japonica* used in this study was obtained from the culture collection of the National Marine Biodiversity Institute of Korea (MABIK) (No. KNU000166).

CONFLICTS OF INTEREST

The authors declare that they have no potential conflicts of interest.

REFERENCES

- Adl, S. M., Bass, D., Lane, C. E., Lukeš, J., Schoch, C. L., Smirnov, A., Agatha, S., Berney, C., Brown, M. W., Burki, F., Cárdenas, P., Čepička, I., Chistyakova, L., Del Campo, J., Dunthorn, M., Edvardsen, B., Eglit, Y., Guillou, L., Hampl, V., Heiss, A. A., Hoppenrath, M., James, T. Y., Karnkowska, A., Karpov, S., Kim, E., Kolisko, M., Kudryavtsev, A., Lahr, D. J. G., Lara, E., Le Gall, L., Lynn, D. H., Mann, D. G., Massana, R., Mitchell, E. A. D., Morrow, D., Park, J. S., Pawlowski, J. W., Powell, M. J., Richter, D. J., Rueckert, S., Shadwick, L., Shimano, S., Spiegel, F. W., Torruella, G., Youssef, N., Zlatogursky, V. & Zhang, Q. 2019. Revisions to the classification, nomenclature, and diversity of eukaryotes. *J. Eukaryot. Microbiol.* 66:4–119.
- Aleem, A. A. 1952. *Olpidiopsis feldmannii* sp. nov., Champignon marin parasite d'Algues de la famille des Bonnemaisoniacees. *C. R. Acad. Sci. Paris* 235:1250–1252.
- Arasaki, S. 1947. Studies on the rot of *Porphyra tenera* by a *Pythium*. *Nippon Suisan Gakkaishi* 13:74–90. (in Japanese)
- Badis, Y., Klochkova, T. A., Brakel, J., Arce, P., Ostrowski, M., Tringe, S. G., Kim, G. H. & Gachon, C. M. M. 2020. Hidden diversity in the oomycete genus *Olpidiopsis* is a potential hazard to red algal cultivation and conservation worldwide. *Eur. J. Phycol.* 55:162–171.
- Badis, Y., Klochkova, T. A., Strittmatter, M., Garvetto, A., Murúa, P., Sanderson, J. C., Kim, G. H. & Gachon, C. M. M. 2019. Novel species of the oomycete *Olpidiopsis* potentially threaten European red algal cultivation. *J. Appl. Phycol.* 31:1239–1250.
- Bari, R. & Jones, J. D. G. 2009. Role of plant hormones in plant defence responses. *Plant Mol. Biol.* 69:473–488.
- Beakes, G. W., Glockling, S. L. & Sekimoto, S. 2012. The evolutionary phylogeny of the oomycete “fungi”. *Protoplasma* 249:3–19.
- Beakes, G. W. & Thines, M. 2017. Hyphochytriomycota and Oomycota. In Archibald, J., Simpson, A. & Slamovits, C. (Eds.) *Handbook of the Protists*. Springer, Cham, pp. 435–505.
- Boller, T. & Felix, G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu. Rev. Plant Biol.* 60:379–406.
- Borowitzka, M. A. 2013. High-value products from microalgae: their development and commercialisation. *J. Appl. Phycol.* 25:743–756.
- Bouwmeester, M. M., Goedknecht, M. A., Poulin, R. & Thieltgies, D. W. 2021. Collateral diseases: aquaculture impacts on wildlife infections. *J. Appl. Ecol.* 58:453–464.

- Brawley, S. H., Blouin, N. A., Ficko-Blean, E., Wheeler, G. L., Lohr, M., Goodson, H. V., Jenkins, J. W., Blaby-Haas, C. E., Helliwell, K. E., Chan, C. X., Marriage, T. N., Bhattacharya, D., Klein, A. S., Badis, Y., Brodie, J., Cao, Y., Collén, J., Dittami, S. M., Gachon, C. M. M., Green, B. R., Karpowicz, S. J., Kim, J. W., Kudahl, U. J., Lin, S., Miche, G., Mittag, M., Olson, B. J. S. C., Pangilinan, J. L., Peng, Y., Qiu, H., Shu, S., Singer, J. T., Smith, A. G., Sprecher, B. N., Wagner, V., Wang, W., Wang, Z.-Y., Yan, J., Yarish, C., Zäuner-Riek, S., Zhuang, Y., Zou, Y., Lindquist, E. A., Grimwood, J., Barry, K. W., Rokhsar, D. S., Schmutz, J., Stiller, J. W., Grossman, A. R. & Prochnik, S. E. 2017. Insights into the red algae and eukaryotic evolution from the genome of *Porphyra umbilicalis* (Bangiophyceae, Rhodophyta). *Proc. Natl. Acad. Sci. U. S. A.* 114:E6361–E6370.
- Buaya, A. T., Ploch, S., Inaba, S. & Thines, M. 2019. Holocarpic oomycete parasitoids of red algae are not *Olpidiopsis*. *Fung. Syst. Evol.* 4:21–31.
- Buaya, A. T., Scholz, B. & Thines, M. 2021. *Sirolopidium bryopsidis*, a parasite of green algae, is probably conspecific with *Pontisma lagenidioides*, a parasite of red algae. *J. Syst. Evol.* 7:223–231.
- Buaya, A. T. & Thines, M. 2020. Diatomophthoraceae: a new family of *Olpidiopsis*-like diatom parasitoids largely unrelated to *Ectrogella*. *Fungal Syst. Evol.* 5:113–118.
- Cai, J., Lovatelli, A., Aguilar-Manjarrez, J., Cornish, L., Dabbadie, L., Desrochers, A., Diffey, S., Garrido Gamarro, E., Geehan, J., Hurtado, A., Lucente, D., Mair, G., Miao, W., Potin, P., Przybyla, C., Reantaso, M., Roubach, R., Tauati, M. & Yuan, X. 2021. *Seaweeds and microalgae: an overview for unlocking their potential in global aquaculture development*. FAO Fisheries and Aquaculture Circular. No. 1229. Food and Agriculture Organization, Rome, 48 pp.
- Chakraborty, S., Moeder, W. & Yoshioka, K. 2017. Plant immunity. In Roitberg, B. D. (Ed.) *Reference Module in Life Sciences*. Elsevier, Amsterdam, pp. 1–8.
- Choudhury, F. K., Rivero, R. M., Blumwald, E. & Mittler, R. 2017. Reactive oxygen species, abiotic stress and stress combination. *Plant J.* 90:856–867.
- Coll, N. S., Epple, P. & Dangl, J. L. 2011. Programmed cell death in the plant immune system. *Cell Death Differ.* 18:1247–1256.
- Cottier-Cook, E. J., Nagabhatla, N., Asri, A., Beveridge, M., Bianchi, P., Bolton, J., Bondad-Reantaso, M. G., Brodie, J., Buschmann, A., Cabarubias, J., Campbell, I., Chopin, T., Critchley, A., De Lombaerde, P., Doumeizel, V., Gachon, C. M. M., Hayashi, L., Hewitt, C. L., Huang, J., Hurtado, A., Kambey, C., Kim, G. H., Le Masson, V., Lim, P. E., Liu, T., Malin, G., Matoju, I., Montalescot, V., Msuya, F. E., Potin, P., Puspita, M., Qi, Z., Shaxson, L., Pinto, I. S., Stentiford, G. D., Suvo, J. & Yarish, C. 2021. *Ensuring the sustainable future of the rapidly expanding global seaweed aquaculture industry: a vision*. United Nations University (Institute on Comparative Regional Integration Studies) and Scottish Association for Marine Science Press, Hamilton, ON, 14 pp.
- Craigie, J. S. & Shacklock, P. F. 1995. Culture of Irish moss. In Boghen, A. D. (Ed.) *Cold-water Aquaculture in Atlantic Canada*. University of Moncton Press, Moncton, pp. 243–270.
- De Cock, A. W. A. M. 1986. Marine *Pythiaceae* from decaying seaweeds in the Netherlands. *Mycotaxon* 25:101–110.
- de Oliveira, L. S., Tschoeke, D. A., Magalhães Lopes, A. C. R., Sudatti, D. B., Meirelles, P. M., Thompson, C. C., Pereira, R. C. & Thompson, F. L. 2017. Molecular mechanisms for microbe recognition and defense by the red seaweed *Laurencia dendroidea*. *mSphere* 2:e00094-17.
- Diehl, N., Kim, G. H. & Zuccarello, G. C. 2017. A pathogen of New Zealand *Pyropia plicata* (Bangiales, Rhodophyta), *Pythium porphyrae* (Oomycota). *Algae* 32:29–39.
- Ding, H. & Ma, J. 2005. Simultaneous infection by red rot and chytrid diseases in *Porphyra yezoensis* Ueda. *J. Appl. Phycol.* 17:51–56.
- Ding, L.-N., Li, Y.-T., Wu, Y.-Z., Li, T., Geng, R., Cao, J., Zhang, W. & Tan, X.-L. 2022. Plant disease resistance-related signaling pathways: recent progress and future prospects. *Int. J. Mol. Sci.* 23:16200.
- Fawke, S., Doumane, M. & Schornack, S. 2015. Oomycete interactions with plants: infection strategies and resistance principles. *Microbiol. Mol. Biol. Rev.* 79:263–280.
- Fletcher, K., Uljević, A., Tsigotgi, A., Antolić, B., Katsaros, C., Nikolić, V., van West, P. & Küpper, F. C. 2015. New record and phylogenetic affinities of the oomycete *Olpidiopsis feldmanni* infecting *Asparagopsis* sp. (Rhodophyta). *Dis. Aquat. Org.* 117:45–57.
- Fujita, Y. & Migita, S. 1980. Death of parasitic *Pythium porphyrae* by drying and freeze-preservation of red rot infected thalli of *Porphyra yezoensis*. *Bull. Fac. Fish. Nagasaki Univ.* 49:11–16.
- Gachon, C. M. M., Sime-Ngando, T., Strittmatter, M., Chambouvet, A. & Kim, G. H. 2010. Algal diseases: spotlight on a black box. *Trends Plant Sci.* 15:633–640.
- Hallmann, A. 2007. Algal transgenics and biotechnology. *Transgenic Plant J.* 1:81–98.
- Hardham, A. R. 2007. Cell biology of plant-oomycete interactions. *Cell. Microbiol.* 9:31–39.
- Herrero, M.-L., Brurberg, M. B., Ojeda, D. I. & Roleda, M. Y. 2020. Occurrence and pathogenicity of *Pythium* (Oomy-

- cota) on *Ulva* species (Chlorophyta) at different salinities. *Algae* 35:79–89.
- Hyde, K. D., Nilsson, R. H., Alias, S. A., Ariyawansa, H. A., Blair, J. E., Cai, L., de Cock, A. W. A. M., Dissanayake, A. J., Glockling, S. L., Goonasekara, I. D., Gorczak, M., Hahn, M., Jayawardena, R. S., van Kan, J. A. L., Laurence, M. H., Lévesque, C. A., Li, X., Liu, J.-K., Maharachchikumbura, S. S. N., Manamgoda, D. S., Martin, F. N., McKenzie, E. H. C., McTaggart, A. R., Mortimer, P. E., Nair, P. V. R., Pawłowska, J., Rintoul, T. L., Shivas, R. G., Spies, C. F. J., Summerell, B. A., Taylor, P. W. J., Terhem, R. B., Udayanga, D., Vaghefi, N., Walther, G., Wilk, M., Wrzosek, M., Xu, J.-C., Yan, J. & Zhou, N. 2014. One stop shop: backbone trees for important phytopathogenic genera: I (2014). *Fungal Divers.* 67:21–125.
- Im, S. H., Klochkova, T. A., Lee, D. J., Gachon, C. M. M. & Kim, G. H. 2019. Genetic toolkits of the red alga *Pyropia tenera* against the three most common diseases in *Pyropia* farms. *J. Phycol.* 55:801–815.
- Iwasaki, A. & Medzhitov, R. 2010. Regulation of adaptive immunity by the innate immune system. *Science* 327:291–295.
- Jiang, Y. N., Haudenschild, J. S. & Hartman, G. L. 2012. Characterization of *Pythium* spp. from soil samples in Illinois. *Can. J. Plant Pathol.* 34:448–454.
- Jones, J. D. G. & Dangl, J. L. 2006. The plant immune system. *Nature* 444:323–329.
- Kamoun, S., Furzer, O., Jones, J. D. G., Judelson, H. S., Ali, G. S., Dalio, R. J. D., Roy, S. G., Schena, L., Zambounis, A., Panabières, F., Cahill, D., Ruocco, M., Figueiredo, A., Chen, X.-R., Hulvey, J., Stam, R., Lamour, K., Gijzen, M., Tyler, B. M., Grünwald, N. J., Mukhtar, M. S., Tomé, D. F. A., Tör, M., Van Den Ackerveken, G., McDowell, J., Daayf, F., Fry, W. E., Lindqvist-Kreuzer, H., Meijer, H. J. G., Petre, B., Ristaino, J., Yoshida, K., Birch, P. R. J. & Govers, F. 2015. The Top 10 oomycete pathogens in molecular plant pathology. *Mol. Plant Pathol.* 16:413–434.
- Kim, G. H., Moon, K.-H., Kim, J.-Y., Shim, J. & Klochkova, T. A. 2014. A revaluation of algal diseases in Korean *Pyropia* (*Porphyra*) sea farms and their economic impact. *Algae* 29:249–265.
- Kim, Y. T., Kim, R.-W., Shim, E., Park, H., Klochkova, T. A. & Kim, G. H. 2023. Control of oomycete pathogens during *Pyropia* farming and processing using calcium propionate. *Algae* 38:71–80.
- Klochkova, T. A., Jung, S. & Kim, G. H. 2017a. Host range and salinity tolerance of *Pythium porphyrae* may indicate its terrestrial origin. *J. Appl. Phycol.* 29:371–379.
- Klochkova, T. A., Kwak, M. S. & Kim, G. H. 2017b. A new endoparasite *Olpidiopsis heterosiphoniae* sp. nov. that infects red algae in Korea. *Algal Res.* 28:264–269.
- Klochkova, T. A., Shim, J. B., Hwang, M. S. & Kim, G. H. 2012. Host–parasite interactions and host species susceptibility of the marine oomycete parasite, *Olpidiopsis* sp., from Korea that infects red algae. *J. Appl. Phycol.* 24:135–144.
- Klochkova, T. A., Shin, Y. J., Moon, K.-H., Motomura, T. & Kim, G. H. 2016. New species of unicellular obligate parasite, *Olpidiopsis pyropiae* sp. nov., that plagues *Pyropia* sea farms in Korea. *J. Appl. Phycol.* 28:73–83.
- Kwak, M. S., Klochkova, T. A., Jeong, S. & Kim, G. H. 2017. *Olpidiopsis porphyrae* var. *koreanae*, an endemic endoparasite infecting cultivated *Pyropia yezoensis* in Korea. *J. Appl. Phycol.* 29:2003–2012.
- Lagerheim, G. 1899. Om växt- och djurlämningarna i Andrées polarboj. *Swedish Soc. Anthropol. Geogr.* 19:425–443.
- Latijnhouwers, M., de Wit, P. J. G. M. & Govers, F. 2003. Oomycetes and fungi: similar weaponry to attack plants. *Trends Microbiol.* 11:462–469.
- Lee, S. J., Hwang, M. S., Park, M. A., Baek, J. M., Ha, D.-S., Lee, J. E. & Lee, S.-R. 2015. Molecular identification of the algal pathogen *Pythium chondricola* (Oomycetes) from *Pyropia yezoensis* (Rhodophyta) using ITS and *cox1* markers. *Algae* 30:217–222.
- Lee, S. J., Jee, B. Y., Son, M.-H. & Lee, S.-R. 2017. Infection and *cox2* sequence of *Pythium chondricola* (Oomycetes) causing red rot disease in *Pyropia yezoensis* (Rhodophyta) in Korea. *Algae* 32:155–160.
- Lee, S. J. & Lee, S.-R. 2022. Rapid detection of red rot disease pathogens (*Pythium chondricola* and *P. porphyrae*) in *Pyropia yezoensis* (Rhodophyta) with PCR-RFLP. *Plant Dis.* 106:30–33.
- LeVesque, C. A. & De Cock, A. W. A. M. 2004. Molecular phylogeny and taxonomy of the genus *Pythium*. *Mycol. Res.* 108:1363–1383.
- Magnus, P. 1872. Ueber *Chytridium tumefaciens* n. sp. in den Wurzelhaaren von *Ceramium flabelligerum* und acanthonotum und andere in Meeresalgen lebende. *Sber. Ges. Naturf. Freunde Berl.* 1872:87–90.
- Mayor, A., Martinon, F., De Smedt, T., Pétrilli, V. & Tschopp, J. 2007. A crucial function of SGT1 and HSP90 in inflammasome activity links mammalian and plant innate immune responses. *Nat. Immunol.* 8:497–503.
- Moon, J.-S., Hong, C.-Y., Lee, J.-W. & Kim, G.-H. 2022. ROS signaling mediates directional cell elongation and somatic cell fusion in the red alga *Griffithsia monilis*. *Cells* 11:2124.
- Nguyen, H. D. T., Dodge, A., Dadej, K., Rintoul, T. L., Ponomareva, E., Martin, F. N., de Cock, A. W. A. M., Lévesque, C. A., Redhead, S. A. & Spies, C. F. J. 2022. Whole genome sequencing and phylogenomic analysis show support

- for the splitting of genus *Pythium*. *Mycologia* 114:501–515.
- Ortiz, D. & Dodds, P. N. 2018. Plant NLR origins traced back to green algae. *Trends Plant Sci.* 23:651–654.
- Park, C., Sakaguchi, K., Kakinuma, M. & Amano, H. 2000. Comparison of the morphological and physiological features of the red rot disease fungus *Pythium* sp. isolated from *Porphyra yezoensis* from Korea and Japan. *Fish Sci.* 66:261–269.
- Park, C. S., Kakinuma, M. & Amano, H. 2001. Detection and quantitative analysis of zoospores of *Pythium porphyrae*, causative organism of red rot disease in *Porphyra*, by competitive PCR. *J. Appl. Phycol.* 13:433–441.
- Peláez-Vico, M. Á., Fichman, Y., Zandalinas, S. I., Van Breusegem, F., Karpiński, S. M. & Mittler, R. 2022. ROS and redox regulation of cell-to-cell and systemic signaling in plants during stress. *Free Radic. Biol. Med.* 193:354–362.
- Perez, I. B. & Brown, P. J. 2014. The role of ROS signaling in cross-tolerance: from model to crop. *Front. Plant Sci.* 5:754.
- Petersen, H. E. 1905. Contributions a la connaissance des Phycomycetes marins (Chytridineae Fischer). *Oversigt Kgl. Danske Vidensk. Selskabs Forhandl.* 5:439–488.
- Phillips, A. J., Anderson, V. L., Robertson, E. J., Secombes, C. J. & van West, P. 2008. New insights into animal pathogenic oomycetes. *Trends Microbiol.* 16:13–19.
- Potin, P., Bouarab, K., Salaün, J.-P., Pohnert, G. & Kloareg, B. 2002. Biotic interactions of marine algae. *Curr. Opin. Plant Biol.* 5:308–317.
- Qiu, L., Mao, Y., Tang, L., Tang, X. & Mo, Z. 2019. Characterization of *Pythium chondricola* associated with red rot disease of *Pyropia yezoensis* (Ueda) (Bangiales, Rhodophyta) from Lianyungang, China. *J. Oceanol. Limnol.* 37:1102–1112.
- Roh, H. & Kannimuthu, D. 2023. Assessments of epidemic spread in aquaculture: comparing different scenarios of infectious bacteria incursion through spatiotemporal hybrid modeling. *Front. Vet. Sci.* 10:1205506.
- Saraiva, M., Ściślak, M. E., Ascurra, Y. T., Ferrando, T. M., Zic, N., Henard, C., van West, P., Trusch, F. & Vleeshouwers, V. G. A. A. 2023. The molecular dialog between oomycete effectors and their plant and animal hosts. *Fungal Biol. Rev.* 43:100289.
- Sekimoto, S., Klochkova, T. A., West, J. A., Beakes, G. W. & Honda, D. 2009. *Olpidiopsis bostrychiae* sp. nov.: an endoparasitic oomycete that infects *Bostrychia* and other red algae (Rhodophyta). *Phycologia* 48:460–472.
- Sekimoto, S., Yokoo, K., Kawamura, Y. & Honda, D. 2008. Taxonomy, molecular phylogeny, and ultrastructural morphology of *Olpidiopsis porphyrae* sp. nov. (Oomycetes, straminipiles), a unicellular obligate endoparasite of *Bangia* and *Porphyra* spp. (Bangiales, Rhodophyta). *Mycol. Res.* 112:361–374.
- Shim, E., Lee, J. W., Park, H., Zuccarello, G. C. & Kim, G. H. 2022. Hydrogen peroxide signalling mediates fertilization and post-fertilization development in the red alga *Bostrychia moritziana*. *J. Exp. Bot.* 73:727–741.
- Shine, M. B., Xiao, X., Kachroo, P. & Kachroo, A. 2019. Signaling mechanisms underlying systemic acquired resistance to microbial pathogens. *Plant Sci.* 279:81–86.
- Sparrow, F. K. 1934. Observations on marine phycomycetes collected in Denmark. *Dansk. Bot. Arkiv.* 8:1–24.
- Sparrow, F. K. 1960. *Aquatic Phycomycetes*. 2nd ed. University of Michigan Press, Ann Arbor, MI, 1187 pp.
- Swarupa, V., Ravishankar, K. V. & Rekha, A. 2014. Plant defense response against *Fusarium oxysporum* and strategies to develop tolerant genotypes in banana. *Planta* 239:735–751.
- Takahashi, M., Ichitani, T. & Sasaki, M. 1977. *Pythium porphyrae* Takahashi et Sasaki, sp. nov. causing red rot of marine red algae *Porphyra* spp. *Trans. Mycol. Soc. Jpn.* 18:279–285.
- Tang, L., Qiu, L., Liu, C., Du, G., Mo, Z., Tang, X. & Mao, Y. 2019. Transcriptomic insights into innate immunity responding to red rot disease in red alga *Pyropia yezoensis*. *Int. J. Mol. Sci.* 20:5970.
- Tedesco, P., Saraiva, M., Sandoval-Sierra, J. V., Fioravanti, M. L., Morandi, B., Dieguez-Urbeondo, J., van West, P. & Galuppi, R. 2021. Evaluation of potential transfer of the pathogen *Saprolegnia parasitica* between farmed salmonids and wild fish. *Pathogens* 10:926.
- Torres, M. A., Jones, J. D. G. & Dangel, J. L. 2006. Reactive oxygen species signaling in response to pathogens. *Plant Physiol.* 141:373–378.
- Ullmann, J. & Grimm, D. 2021. Algae and their potential for a future bioeconomy, landless food production, and the socio-economic impact of an algae industry. *Org. Agric.* 11:261–267.
- van Bentem, S. F., Vossen, J. H., de Vries, K. J., van Wees, S., Tameling, W. I. L., Dekker, H. L., de Koster, C. G., Haring, M. A., Takken, F. L. W. & Cornelissen, B. J. C. 2005. Heat shock protein 90 and its co-chaperone protein phosphatase 5 interact with distinct regions of the tomato I-2 disease resistance protein. *Plant J.* 43:284–298.
- Van der Meer, J. P. & Pueschel, C. M. 1985. *Petersenia palmariae* n. sp. (Oomycetes): a pathogenic parasite of the red alga *Palmaria mollis* (Rhodophyceae). *Can. J. Bot.* 63:404–408.
- Weinberger, F. 1999. Epiphyte-host interactions: *Gracilaria conferta* (Rhodophyta) and associated bacteria. Ph.D.

- dissertation, Christian Albrechts Universität, Kiel, Germany, 150 pp.
- Weinberger, F., Leonardi, P., Miravalles, A., Correa, J. A., Lion, U., Kloareg, B. & Potin, P. 2005. Dissection of two distinct defense-related responses to agar oligosaccharides in *Gracilaria chilensis* (Rhodophyta) and *Gracilaria conferta* (Rhodophyta). J. Phycol. 41:863–873.
- Whittick, A. & South, G. R. 1972. *Olpidiopsis antithamnionis* n. sp. (Oomycetes, Olpidiopsidaceae), a parasite of *Antithamnion floccosum* (O. F. Müll.) Kleen from Newfoundland. Arch. Mikrobiol. 82:353–360.