

# LECTURE 1

RUNLIN ZHANG

## CONTENTS

4	Notation	1
5	1. Lecture 1, Dirichlet's theorem, badly approximable numbers and Khintchin's zero-one law	1
6	1.1. Foreword	1
7	1.2. The beginning	2
8	1.3. Badly approximable numbers	2
9	1.4. Littlewood conjecture	3
10	1.5. Khintchine's zero-one law	3
11	1.6. Proof of the convergence part	4
12	1.7. Proof of the divergence part	4
13	1.8. Cassels' zero-one law	4
14	1.9. Partial converse to Borel–Cantelli	5
15	1.10. A reduction	6
16	1.11. Quasi-independence	7
17	1.12. Average of Euler's totient function	8
18	References	9

## NOTATION

The set of positive integers is denoted by  $\mathbb{Z}^+$ . For a real number  $x$ , let  $\langle x \rangle$  denote the distance to the nearest integer, namely,  $\langle x \rangle = \inf_{n \in \mathbb{Z}} |x - n|$ .  $\text{Leb}$  denotes the standard Lebesgue measure on  $\mathbb{R}^n$  where the  $n$  is understood from the context.

For  $x, y \in \mathbb{Z}$  non-zero, we let  $\gcd(x, y) \in \mathbb{Z}^+$  to be the greatest common divisor of  $|x|$  and  $|y|$ . If one of them is zero but the other is not, we set  $\gcd(x, y)$  to be the absolute value of the non-zero one. Also,  $\gcd(0, 0) := 0$ . Two integers are said to be coprime iff  $\gcd(x, y) = 1$ .

Abbreviate "infinitely many" as "i.m."; "almost every" as "a.e.".

## 1. LECTURE 1, DIRICHLET'S THEOREM, BADLY APPROXIMABLE NUMBERS AND KHintCHIN'S ZERO-ONE LAW

References: I am mostly following [Zaf17, Cas50]. One may also consult the survey [BRV16] (available on arxiv).

**1.1. Foreword.** Number theory provides a huge amount of interesting problems. Besides "elementary" methods, tools from different branches of math are introduced to solve them. Assuming Galois theory and ring theory, one can give an introduction to number fields. Assuming complex analysis, one can study Riemann zeta functions, Dirichlet L functions or modular forms.

This course is concerned with so-called "Diophantine approximation" problems, which are concerned with approximating real numbers by rational numbers. Actually, we will focus on a specific (still unsolved!) problem: Littlewood conjecture. We will present the work of Einsiedler–Katok–Lindenstrauss on this conjecture, showing the exception set has dimension zero. They use tools coming from dynamics, which will be introduced later.

As this course is supposed to be introductory, we will start with some basics before discussing the deep work of EKL.

1.2. **The beginning.** The starting point of Diophantine approximation is the following:

**Fact 1.1.** *The set of rational numbers  $\mathbb{Q}$  is dense in real numbers  $\mathbb{R}$ . In other words, for every  $x \in \mathbb{R}$  and  $\varepsilon > 0$ , there exist two integers  $(p, q)$  with  $q > 0$  such that  $\left|x - \frac{p}{q}\right| < \varepsilon$ .*

To have a better approximation of  $x \in \mathbb{R}$ , one should use rational numbers with large denominators. How large it has to be?

**Theorem 1.2** (Dirichlet). *For every  $x \in \mathbb{R}$  and  $N \in \mathbb{Z}^+$ , there exists  $(p, q) \in \mathbb{Z}^2$  with  $0 < q \leq N$  such that*

$$\left|x - \frac{p}{q}\right| < \frac{1}{Nq}.$$

The proof is based on “drawer’s principle” (or pigeon-hole principle).

*Proof.* For  $k = 1, 2, \dots, N$  find  $n_k \in \mathbb{Z}$  such that  $kx - n_k \in [0, 1)$ . Write

$$\{kx - n_k, k = 1, \dots, N\} = \{x_1 \leq x_2 \leq \dots \leq x_N\}.$$

Thus, one of the numbers

$$\{x_1, x_2 - x_1, x_3 - x_2, \dots, x_N - x_{N-1}, 1 - x_N\}$$

has to be strictly smaller than  $\frac{1}{N}$ . Say  $x_{i_0} - x_{i_0-1} < \frac{1}{N}$ . By convention,  $x_0 := 0 = 0 \cdot x$  and  $x_{N+1} := 1 = 0 \cdot x + 1$ . Therefore, for some integers  $\{k_1 < k_2\} \subset \{0, \dots, N\}$ , one has for some  $p \in \mathbb{Z}$ ,

$$|k_2x - k_1x - p| < \frac{1}{N}$$

Let  $q := k_2 - k_1$ , then

$$\left|x - \frac{p}{q}\right| < \frac{1}{Nq},$$

proving the assertion.  $\square$

1.3. **Badly approximable numbers.** As a corollary of the above theorem, one gets

**Corollary 1.3.** *For every  $x \in \mathbb{R}$ , there exist infinitely many pairs of integers  $(p, q)$  such that*

$$\left|x - \frac{p}{q}\right| < \frac{1}{q^2}.$$

*In other words,  $\langle qx \rangle < \frac{1}{q}$  for infinitely many  $q \in \mathbb{Z}^+$ .*

**Definition 1.4.** *A real number  $x$  is said to be **badly approximable** iff there exists  $c > 0$  such that*

$$\left|x - \frac{p}{q}\right| > \frac{c}{q^2}, \quad \forall (p, q) \in \mathbb{Z}^2, q > 0.$$

*Or in other words,  $\langle qx \rangle q > c$  for all  $q \in \mathbb{Z}^+$ . We will let **BAD** denote the set of badly approximable numbers. If an irrational number  $x \in \mathbb{R} \setminus \mathbb{Q}$  is not badly approximable, we say that  $x$  is **well-approximable**.*

This definition is non-trivial in the sense that there are badly approximable numbers as well as well-approximable numbers.

**Example 1.5.**  $\sqrt{2}$  is badly approximable.

*Proof.* Take  $\varepsilon \in (0, 1)$ . Assume that there are integers  $p, q$  with  $q > 0$  such that

$$q \left|q\sqrt{2} - p\right| < \varepsilon.$$

Thus

$$\left|q\sqrt{2} + p\right| < \varepsilon/q + 2q\sqrt{2} < 4q.$$

Multiplying the above two together gives

$$q |2q^2 - p^2| < 4q\varepsilon \implies |2q^2 - p^2| < 4\varepsilon.$$

But  $2q^2 - p^2$  is a non-zero integer, so  $|2q^2 - p^2| \geq 1$ . Thus  $1 < 4\varepsilon$ . This finishes the proof, showing  $q\langle q\sqrt{2} \rangle \geq \frac{1}{4}$  for every  $q \in \mathbb{Z}^+$ .  $\square$

**Conjecture 1.6.** *Algebraic numbers that are not contained in a quadratic number field are not bad.*

1 So far no single example seems known about this conjecture. For instance, it is un-  
 2 known whether  $\sqrt[3]{2}$  is badly approximable or not.

3 **Remark 1.7.** *However, for every  $\varepsilon > 0$  and irrational algebraic number  $x$ , there exists*  
 4  *$c = c(x, \varepsilon) > 0$  such that  $q^{1+\varepsilon} \langle qx \rangle > c(x, \varepsilon)$  for every  $q \in \mathbb{Z}^+$ . This is a theorem of*  
 5 *Roth. Its generalization by Schmidt, known as subspace theorem, has applications to*  
 6 *other problems in number theory.*

7 **Example 1.8.** *0.10100001000000001... ( the  $n$ -th group of 0's consists of  $n + m$  consec-*  
 8 *utive zeros if there are  $m$  digits in front of it) is well-approximable.*

#### 9 1.4. Littlewood conjecture.

10 **Conjecture 1.9** (Littlewood). *For every pair  $(x, y)$  of real numbers, for every  $\varepsilon > 0$ ,*  
 11 *there exists  $q \in \mathbb{Z}^+$  such that*

$$q \langle qx \rangle \langle qy \rangle < \varepsilon.$$

12 *Equivalently,*

$$\inf_{q \in \mathbb{Z}^+} q \langle qx \rangle \langle qy \rangle = 0. \quad (1)$$

13 **Remark 1.10.** *Note that if one of  $x$  or  $y$  does not belong to **BAD**, then Equa.(1) holds.*

**Theorem 1.11** (Einsiedler–Katok–Lindenstrauss [EKL06]).

$$\dim\{(x, y) \in \mathbb{R}^2, \text{ Equa.(1) fails } \} = 0.$$

14 This will be proved in later lectures. We will soon prove that

15 **Theorem 1.12.**  $\text{Leb}(\mathbf{BAD}) = 0$ . *Consequently,*

$$\text{Leb}\{(x, y) \in \mathbb{R}^2, \text{ Equa.(1) fails } \} = 0.$$

16 The following two theorems will not be proved, but I find it healthy to compare them  
 17 with Littlewood conjecture and EKL's work.

**Theorem 1.13** (Gallagher).

$$\text{Leb}\{(x, y) \in \mathbb{R}^2, \inf_{q \in \mathbb{Z}^+} q \langle qx \rangle \langle qy \rangle \cdot (\log q)^2 = 0\} = 0.$$

**Theorem 1.14** (Badziahin [Bad13]).

$$\dim\{(x, y) \in \mathbb{R}^2, \inf_{q \in \mathbb{Z}^+} q \langle qx \rangle \langle qy \rangle \cdot \log q \log \log q = 0\} = 2.$$

18 We will sometimes restrict our attention to numbers in the interval  $[0, 1)$  without loss  
 19 of generality.

20 **1.5. Khintchine's zero-one law.** Let  $\psi : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$  be a sequence of positive real  
 21 numbers (for instance  $\psi(q) := q^{-1}$ ). Define

$$\begin{aligned} W(\psi) &:= \left\{ x \in [0, 1), \left| x - \frac{p}{q} \right| < q^{-1} \psi(q) \text{ for i.m. } q \in \mathbb{Z}^+, p \in \mathbb{Z} \right\} \\ &= \{ x \in [0, 1), \langle qx \rangle < \psi(q) \text{ for i.m. } q \in \mathbb{Z}^+ \} \end{aligned} \quad (2)$$

22 **Theorem 1.15** (Khintchin). *Assume  $\psi$  is non-increasing. Then,*

$$\text{Leb}(W(\psi)) = \begin{cases} 0 & \text{if } \sum \psi(n) < +\infty \\ 1 & \text{if } \sum \psi(n) = +\infty \end{cases}.$$

23 **Remark 1.16.** *The assumption that  $\psi$  is non-increasing is necessary.*

24 That  $\text{Leb}(\mathbf{BAD}) = 0$  follows directly from this theorem.

25 *Proof of Theorem 1.12 assuming Theorem 1.15.* For every  $c > 0$  and  $q \in \mathbb{Z}^+$ , let  $\psi_c(q) :=$   
 26  $cq^{-1}$ . Then, **BAD** is the complement in  $[0, 1)$  of the union of  $W(\psi_{n^{-1}})$  as  $n$  ranges over  
 27 positive integers. Thus it suffices to show that  $\text{Leb}(W(\psi_c)) = 1$  for every  $c > 0$ . By  
 28 Theorem 1.15, this follows from the fact that  $\sum_{n \in \mathbb{Z}^+} cn^{-1} = +\infty$ .  $\square$

1 **1.6. Proof of the convergence part.** In this subsection, we explain the convergence  
 2 part of Theorem 1.15. Namely, we assume  $\sum \psi(n) < +\infty$  and prove  $\text{Leb}(W(\psi)) = 0$ .  
 3 For this one uses the Borel–Cantelli lemma:

4 **Lemma 1.17.** *Let  $(E_n)_{n \in \mathbb{Z}^+}$  be a sequence of measurable subsets of  $[0, 1)$  such that*  
 5  *$\sum \text{Leb}(E_n) < +\infty$ . Then*

$$\text{Leb}(\{x \in E_n \text{ for i.m. } n\}) = 0.$$

6 **Remark 1.18.** *The set  $\{x \in E_n \text{ for i.m. } n\}$  is sometimes written as  $\limsup E_n$ .*

7 Let

$$W_n(\psi) := \{x \in [0, 1) \mid \langle nx \rangle < \psi(n)\}.$$

8 In light of Lemma 1.17, it suffices to show that  $\sum \text{Leb}(W_n(\psi)) < \infty$ . Indeed, for  $n$  large  
 9 enough (such that  $\psi(n) < 0.5$ ),

$$\begin{aligned} W_n(\psi) &= \bigcup_{i=0,1,\dots,n-1} \{x \in W_n(\psi) \mid nx - i \in [0, 1)\} \\ &= \bigcup_{i=0,1,\dots,n-1} \left\{ x \in \left[\frac{i}{n}, \frac{i+1}{n}\right), nx \in [i, i + \psi(n)) \cup ((i+1) - \psi(n), i+1) \right\} \\ &= \bigcup_{i=0,1,\dots,n-1} \left[ \frac{i}{n}, \frac{i}{n} + \frac{\psi(n)}{n} \right) \cup \left( \frac{i+1}{n} - \frac{\psi(n)}{n}, \frac{i+1}{n} \right). \end{aligned}$$

10 Hence,

$$\text{Leb}(W_n(\psi)) = \sum_{i=0,1,\dots,n-1} \frac{2\psi(n)}{n} = 2\psi(n).$$

11 Thus the divergence of  $\sum \text{Leb}(W_n(\psi))$  follows.

12 **1.7. Proof of the divergence part.** From now on assume  $\sum \psi(n) = +\infty$  and we wish  
 13 to show  $\text{Leb}(W(\psi)) = 1$ . The proof will consist of two steps:  $\text{Leb}(W(\psi)) > 0$  and  
 14  $\text{Leb}(W(\psi)) > 0 \implies \text{Leb}(W(\psi)) = 1$ .

15 **1.8. Cassels' zero-one law.** In this subsection we prove

16 **Theorem 1.19.**  $\text{Leb}(W(\psi)) = 0$  or  $1$ .

17 Though we use the non-increasing feature of  $\psi$  below, this assumption can be removed  
 18 without much effort.

19 Choose a bijection  $n \mapsto \lambda_n$  from  $\mathbb{Z}^+$  to  $\mathbb{Q}$ . For a rational number  $x$ , find coprime  
 20 integers  $p, q$  with  $q > 0$  (if  $p = 0$ , we set  $q := 1$ ) such that  $x = \frac{p}{q}$  and define  $\Psi_{\text{red}}(x) :=$   
 21  $q^{-1}\psi(q)$ . One can check that

$$W(\psi) = W_{\text{red}}(\psi) := \{x \in [0, 1) \mid |x - \lambda_n| < \Psi_{\text{red}}(\lambda_n) \text{ for i.m. } n\}.$$

22 For  $k, N \in \mathbb{Z}^+$ , let

$$\begin{aligned} E_k &:= \left\{ x \in [0, 1) \mid |x - \lambda_n| < \frac{1}{k} \Psi_{\text{red}}(\lambda_n) \text{ for i.m. } n \right\}, \\ E_k^N &:= \left\{ x \in [0, 1) \mid |x - \lambda_n| < \frac{1}{k} \Psi_{\text{red}}(\lambda_n) \text{ for some } n > N \right\}. \end{aligned}$$

23 Also let  $E_\infty := \bigcap E_k$ . So  $E_1 = W_{\text{red}}(\psi)$  and  $E_k = \bigcap_{N=1}^\infty E_k^N$ . Theorem 1.19 would follow  
 24 from the following three lemmas.

25 For a positive integer  $n$  and  $x \in [0, 1)$ , define  $T_n(x)$  to be the unique element in  $[0, 1)$   
 26 such that  $T_n(x) - nx \in \mathbb{Z}$ .

27 **Lemma 1.20.** *For every  $k \in \mathbb{Z}^+$ ,  $T_k(E_\infty) \subset E_1$ .*

28 *Proof.* For  $x \in E_k$  with  $|x - \lambda_n| < k^{-1}\Psi_{\text{red}}(\lambda_n)$ , then  $|kx - k\lambda_n| < \Psi_{\text{red}}(\lambda_n) \leq \Psi_{\text{red}}(k\lambda_n)$ .  
 29 Thus  $kx \in E_1$ .  $\square$

30 **Lemma 1.21.** *For every measurable set  $E \subset [0, 1)$  with  $\text{Leb}(E) > 0$ , for every  $\varepsilon > 0$ ,  
 31 there exists  $N \in \mathbb{Z}^+$  such that*

$$\text{Leb}(T_N(E)) > 1 - \varepsilon.$$

1 Let  $I(x, \delta) := (x - \delta, x + \delta)$ . We will need Lebesgue's density theorem (see e.g. chapter 3,  
2 Theorem 1.4 of Stein's book "real analysis") for characteristic functions of Borel subsets.

3 **Theorem 1.22** (Lebesgue density theorem). *Let  $f$  be an integrable function on  $[0, 1]$ .  
4 Then for Lebesgue almost every  $x \in [0, 1]$ , one has*

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\varepsilon} \int_{x-\varepsilon}^{x+\varepsilon} f(t) \text{Leb}(t) = f(x).$$

5 *Proof.* By Lebesgue density theorem, find  $\theta^* \in E$  such that for any  $\varepsilon > 0$ , there exists  
6  $\eta^* > 0$  such that for every  $0 < \eta < \eta^*$ ,

$$\frac{\text{Leb}(E \cap I(\theta^*, \eta))}{2\eta} > 1 - \varepsilon.$$

7 Now choose  $\eta = \frac{1}{2N}$  for  $N \in \mathbb{Z}^+$  large such that the above inequality holds. Then

$$\text{Leb}\left(NE \cap \left(N\theta^* - \frac{1}{2}, N\theta^* + \frac{1}{2}\right)\right) > 1 - \varepsilon.$$

8 By reducing modulo  $\mathbb{Z}$ , we get  $\text{Leb}(T_N(E)) > 1 - \varepsilon$ .  $\square$

9 **Lemma 1.23.** *For every  $k \in \mathbb{Z}^+$ ,  $\text{Leb}(E_1 \setminus E_k) = 0$ . Consequently,  $\text{Leb}(E_\infty) > 0$  if  
10  $\text{Leb}(E_1) > 0$ .*

11 *Proof.* To save notation write  $\alpha_n := \Phi_{\text{red}}(\lambda_n)$  in the proof. Assume  $\text{Leb}(E_1 \setminus E_k) > 0$ ,  
12 find  $N \in \mathbb{Z}^+$  large enough such that  $\text{Leb}(E_1 \setminus E_k^N) > 0$ .

13 By Lebesgue density theorem again, we find  $\theta_{\neq 0}^* \in E_1$  such that for every  $\varepsilon > 0$ , there  
14 exists  $\eta^*(\varepsilon) > 0$  such that for every  $0 < \eta < \eta^*(\varepsilon)$ , one has

$$\text{Leb}(I(\theta^*, \eta) \cap E_k^N) < \varepsilon \text{Leb}(I(\theta^*, \eta)). \quad (3)$$

15 We take  $\varepsilon := \frac{1}{2(k+1)}$  and write  $\eta^* := \eta^*(\varepsilon)$ . Take  $n$  sufficiently large (that is,  $n > N$  and  
16  $2\alpha_n < \eta^*$ ) such that

$$|\theta^* - \lambda_n| < \alpha_n.$$

17 By definition, one has  $I(\lambda_n, \frac{1}{k}\alpha_n) \subset E_k^N$ . Let  $\eta := |\theta^* - \lambda_n| + \frac{1}{k}\alpha_n$ , which is smaller than  
18  $\eta^*$ . Also,  $I(\lambda_n, \frac{1}{k}\alpha_n) \subset I(\theta^*, \eta)$ . Hence,

$$\begin{aligned} \frac{\text{Leb}(I(\lambda_n, \frac{\alpha_n}{k}))}{\text{Leb}(I(\theta^*, \eta))} &= \frac{\alpha_n/k}{\eta} > \frac{\alpha_n/k}{\alpha_n + \alpha_n/k} = \frac{1}{1+k} \\ \implies \text{Leb}(I(\theta^*, \eta) \cap E_k^N) &> \frac{1}{1+k} \text{Leb}(I(\theta^*, \eta)), \end{aligned}$$

19 which is a contradiction against Equ.(3).  $\square$

20 *Proof of Theorem 1.19.* Assume  $\text{Leb}(E_1) > 0$  and want to show  $\text{Leb}(E_1) = 1$ . By Lemma  
21 1.23,  $\text{Leb}(E_\infty) > 0$ . Apply Lemma 1.21 to  $E = E_\infty$ , we get  $\text{Leb}(\bigcup_{n \in \mathbb{Z}^+} T_n(E_\infty)) = 1$ .  
22 But this set is contained in  $E_1$  by Lemma 1.20. So we obtain  $\text{Leb}(E_1) = 1$  and we are  
23 done.  $\square$

24 **1.9. Partial converse to Borel–Cantelli.** The proof of the divergence part is more  
25 difficult partly because the converse to the Borel–Cantelli lemma is not true. However,  
26 we do have a partial converse assuming certain independence properties for the sequence  
27 of sets  $(E_n)$ .

28 **Lemma 1.24.** *Let  $(E_n)$  be a sequence of measurable subsets of  $[0, 1]$ . Then for every  
29 pair of integers  $0 < m < n$ , we have*

$$\text{Leb}\left(\bigcup_{i=m}^n E_i\right) \geq \frac{\left(\sum_{i=m}^n \text{Leb}(E_i)\right)^2}{\sum_{i=m}^n \sum_{j=m}^n \text{Leb}(E_i \cap E_j)}. \quad (4)$$

30 *Proof.* This is a consequence of Cauchy-Schwarz.

$$\left(\int_0^1 \left(\sum_{i=m}^n \mathbf{1}_{E_i}\right) \cdot (\mathbf{1}_{\bigcup_{i=m}^n E_i}) \text{Leb}\right)^2 \leq \int_0^1 \left(\sum_{i=m}^n \mathbf{1}_{E_i}\right)^2 \text{Leb} \cdot \int_0^1 \mathbf{1}_{\bigcup_{i=m}^n E_i}^2 \text{Leb}$$

1 For the left hand side one has:

$$\left( \int_0^1 \left( \sum_{i=m}^n \mathbf{1}_{E_i} \right) \cdot (\mathbf{1}_{\cup_{i=m}^n E_i}) \text{Leb} \right)^2 = \left( \sum_{i=m}^n \text{Leb}(E_i) \right)^2,$$

2 and for the right hand side:

$$\int_0^1 \left( \sum_{i=m}^n \mathbf{1}_{E_i} \right)^2 \text{Leb} \cdot \int_0^1 \mathbf{1}_{\cup_{i=m}^n E_i}^2 \text{Leb} = \sum_{i,j=m}^n \text{Leb}(E_i \cap E_j) \cdot \text{Leb} \left( \bigcup_{i=m}^n E_i \right).$$

3 Putting them together finishes the proof.  $\square$

4 From this lemma, one can easily prove the following converse to Borel–Cantelli.

5 **Lemma 1.25.** *Let  $(E_n)_{n \in \mathbb{Z}^+}$  be a sequence of measurable subsets of  $[0, 1]$  such that*  
 6  *$\sum \text{Leb}(E_n) < \infty$ . Assume furthermore that  $\text{Leb}(E_i \cap E_j) = \text{Leb}(E_i) \text{Leb}(E_j)$  for ev-*  
 7 *ery  $i \neq j$  (namely,  $E_n$ 's are independent from each other). Then*

$$\text{Leb}(\{x \in E_n \text{ for i.m. } n\}) = 1.$$

8 However, our sets are not independent. Nevertheless, we will be able to find a lower  
 9 bound for the RHS of Equa.(4), which shows that  $W(\psi)$  has positive Lebesgue measure.  
 10 Our proof is then complete by invoking Theorem 1.19.

11 **1.10. A reduction.** Let  $\psi_1(n) := \min\{\psi(n), \frac{1}{n}\}$ . As  $W(\psi_1) \subset W(\psi)$ , it suffices to show  
 12 that  $\text{Leb}(W(\psi_1)) = 1$ .

13 **Lemma 1.26.**  $\sum \psi_1(n) = +\infty$ .

14 **Remark 1.27.** *It is not true in general that for two non-increasing sequence  $(a_n)$  and*  
 15  *$(b_n)$  of positive real numbers,  $\sum a_n = \sum b_n = +\infty$  would imply  $\sum \min\{a_n, b_n\} = +\infty$ .*

16 *Proof.* Assuming  $\sum \psi_1(n) < +\infty$ , we will show that  $\sum \psi(n) < +\infty$ , which is a contra-  
 17 diction.

18 Decompose  $\mathbb{Z}^+ \setminus \{1\} = I \sqcup J$  such that

$$i \in I \iff \psi(n) \leq \frac{1}{n}, \quad i \in J \iff \psi(n) > \frac{1}{n}.$$

19 Thus  $\sum_I \psi(n) < +\infty$  and  $\sum_J \frac{1}{n} < +\infty$ . Decompose  $J = \bigsqcup_{i \in \mathbb{Z}^+} J_i$  where  $J_i = \{a_i, a_i +$   
 20  $1, \dots, b_i\}$  and  $b_i + 1 < a_{i+1}$ . Therefore

$$+\infty > \sum_{n \in J} \frac{1}{n} > \sum_{i \in \mathbb{Z}^+} \int_{a_i}^{b_i+1} \frac{1}{x} dx = \log \left( \frac{b_i + 1}{a_i} \right).$$

21 On the other hand,

$$\sum_{n \in J} \psi(n) = \sum_i \sum_{j \in J_i} \psi(n) \leq \sum_i \sum_{j \in J_i} \psi(a_i - 1) = \sum_i \sum_{j \in J_i} \frac{1}{a_i - 1} = \sum_i \frac{b_i - (a_i - 1)}{a_i - 1}.$$

22 Define  $\lambda_i := \frac{b_i}{a_i - 1} - 1 > 0$  for every  $i \in \mathbb{Z}^+$ , then  $\sum_{n \in J} \psi(n) \leq \sum \lambda_i$ , which will be shown  
 23 to be convergent.

24 Note that (for  $a_i > 1$ )

$$\frac{b_i + 1}{a_i} - 1 > \frac{1}{2} \left( \frac{b_i}{a_i - 1} - 1 \right).$$

25 Indeed for  $\frac{p}{q} > 1$  with  $q > 1$ , one has  $\frac{p+1}{q+1} - 1 > \frac{1}{2} \left( \frac{p}{q} - 1 \right)$ . Since this, after the denominators  
 26 are cleared, is equivalent to  $(q-1)(p+q) > 0$ .

27 So we have  $\sum \log(1 + \frac{1}{2} \lambda_i)$  is convergent. This implies that  $\sum \lambda_i$  is convergent by  
 28 Lemma 1.28.  $\square$

29 **Lemma 1.28.** *Let  $(\lambda_n)$  be a sequence of non-negative real numbers, one has that*

$$\sum \lambda_n < +\infty \iff \sum \ln(1 + \lambda_n) < +\infty.$$

30 *Proof.* Note that we may assume that  $(\lambda_n)$  tends to 0 for otherwise both sides are diver-  
 31 gent. For  $x \geq 0$ ,  $\ln(1 + x) \leq x$ . Conversely, for  $x$  sufficiently small,  $\ln(1 + x) > \frac{1}{2}x$ . So  
 32 we are done.  $\square$

33 In light of Lemma 1.26, we will assume  $\psi(n) \leq \frac{1}{2n}$  in the next subsection.

1 **1.11. Quasi-independence.** For this subsection, define

$$E_n := \bigcup_{q=2^{n-1}}^{2^n-1} \bigcup_{p \in \{1, \dots, q\}, \gcd(p, q)=1} I\left(\frac{p}{q}, \frac{\psi(2^n)}{2^n}\right).$$

2 As  $\psi(n) \leq \frac{1}{2n}$ , one can check that for every two distinct indices  $(p, q), (p', q')$  appearing  
3 above,

$$I\left(\frac{p}{q}, \frac{\psi(2^n)}{2^n}\right) \cap I\left(\frac{p'}{q'}, \frac{\psi(2^n)}{2^n}\right) = \emptyset.$$

4 Also, since  $\frac{\psi(2^n)}{2^n} \leq \frac{\psi(q)}{q}$ , the set  $E_n$  is contained in

$$W_n(\psi) := \left\{ x \in [0, 1) \mid \left| x - \frac{p}{q} \right| < \Psi_{\text{red}}\left(\frac{p}{q}\right) \text{ for some } 2^{n-1} \leq q \leq 2^n - 1 \right\}.$$

5 Thus, if  $x$  belongs to  $E_n$  infinitely many  $n$ 's, then  $x$  belongs to  $W(\psi)$ . Therefore, it  
6 suffices to prove that

7 • RHS of Equa.(4) for such  $(E_i)$  has a lower bound independent of  $M$  and for  $N$   
8 large enough, i.e., there exists  $C > 0$  such that for every  $M$ , for  $N$  large enough

$$\sum_{i, j=M}^N \text{Leb}(E_i \cap E_j) \leq C \left( \sum_{i=M}^N \text{Leb}(E_i) \right)^2;$$

9 •  $\sum \text{Leb}(E_n) = +\infty$ .

10 Let  $\phi$  be Euler's totient function. Namely, for a positive integer  $N$ ,  $\phi(N) := |(\mathbb{Z}/N\mathbb{Z})^\times|$   
11 is the number of integers in  $\{1, \dots, N\}$  that are coprime to  $N$ . Firstly we have

$$\text{Leb}(E_n) = \left( 2 \cdot \frac{\psi(2^n)}{2^n} \right) \cdot \sum_{q=2^{n-1}}^{2^n-1} \phi(q). \quad (5)$$

12 Then estimate the Lebesgue measure of  $E_m \cap E_n$  for  $m < n$ . For  $(a, b)$  (resp.  $(c, d)$ )  
13 appearing in the index of  $E_m$  (resp.  $E_n$ ), one has

$$\text{Leb}\left(I\left(\frac{a}{b}, \frac{\psi(2^m)}{2^m}\right) \cap I\left(\frac{c}{d}, \frac{\psi(2^n)}{2^n}\right)\right) \leq \text{Leb}\left(I\left(\frac{c}{d}, \frac{\psi(2^n)}{2^n}\right)\right) = 2 \frac{\psi(2^n)}{2^n}.$$

14 For distinct  $(c_1, d_1), (c_2, d_2)$  appearing in the index of  $E_n$ , one has

$$\left| \frac{c_1}{d_1} - \frac{c_2}{d_2} \right| = \left| \frac{c_1 d_2 - c_2 d_1}{d_1 d_2} \right| \geq \frac{1}{d_1 d_2} \geq \frac{1}{2^{2n}}.$$

15 Thus, for every fixed  $(a, b)$  appearing in the index of  $E_m$ , the number of  $(c, d)$  appearing  
16 in the index of  $E_n$  such that  $I\left(\frac{a}{b}, \frac{\psi(2^m)}{2^m}\right) \cap I\left(\frac{c}{d}, \frac{\psi(2^n)}{2^n}\right) \neq \emptyset$  is at most

$$\frac{2 \frac{\psi(2^m)}{2^m}}{\frac{1}{2} \frac{1}{2^{2n}}} + 2 = 4 \cdot 2^{2n} \cdot \frac{\psi(2^m)}{2^m} + 2.$$

17 Therefore,

$$\text{Leb}(E_m \cap E_n) \leq \left( 2 \frac{\psi(2^n)}{2^n} \right) \cdot \left( 4 \cdot 2^{2n} \cdot \frac{\psi(2^m)}{2^m} + 2 \right) \cdot \left( \sum_{q=2^{m-1}}^{2^m-1} \phi(q) \right)$$

18 Combining with Equa.(5), one has (for  $m < n$ )

$$\text{Leb}(E_m \cap E_n) \leq 2 \cdot \frac{2^{2n}}{\sum_{q=2^{n-1}}^{2^n-1} \phi(q)} \cdot \text{Leb}(E_n) \text{Leb}(E_m) + \left( 4 \frac{\psi(2^n)}{2^n} \right) \cdot \left( \sum_{q=2^{m-1}}^{2^m-1} \phi(q) \right). \quad (6)$$

19 Before proceeding further, note two consequences of Lemma 1.29 to be presented in  
20 the next subsection. There exists a constant  $C > 0$  such that for all positive integers  $k$ ,

$$2^{2k} \leq C \cdot \sum_{q=2^{k-1}}^{2^k-1} \phi(q);$$

21

$$\sum_{q=1}^{2^{k-1}-1} \phi(q) \leq C \cdot \sum_{q=2^{k-1}}^{2^k-1} \phi(q).$$

1 The first inequality and Equa.(5) imply that

$$\sum_{n=1}^N \text{Leb}(E_n) \geq \sum_{n=1}^N 2C^{-1}2^n\psi(2^n) \geq 2C^{-1} \sum_{n=1}^N \sum_{q=2^n}^{2^{n+1}-1} \psi(q) = \sum_{q=2}^{2^{N+1}-1} \psi(q)$$

2 which diverges to  $+\infty$ .

3 Take two positive integers  $M < N$ .

4 Now we go back to Equa.(6) and sum over  $m < n, m, n = M, \dots, N$ . The first summand  
5 in Equa.(6) is bounded from above by

$$\sum_{m < n, m, n = M, \dots, N} 2C \text{Leb}(E_m) \text{Leb}(E_n)$$

6 whereas the second summand is

$$\sum_{n=M}^N 4 \frac{\psi(2^n)}{2^n} \sum_{q=2^{M-1}}^{2^{n-1}-1} \phi(q) \leq \sum_{n=M}^N C \text{Leb}(E_n).$$

7 Consequently,

$$\begin{aligned} \sum_{m, n = M, \dots, N} \text{Leb}(E_m \cap E_n) &\leq 2C \sum_{m, n = M, \dots, N} \text{Leb}(E_m) \text{Leb}(E_n) + 3C \sum_{n=M}^N \text{Leb}(E_n) \\ &= 2C \left( \sum_{n=M}^N \text{Leb}(E_n) \right)^2 + 3C \sum_{n=M}^N \text{Leb}(E_n). \end{aligned}$$

8 Since  $\sum \text{Leb}(E_n)$  diverges, there exists  $C' > 0$  (independent of  $M$ ) such that

$$\sum_{n=M}^N \text{Leb}(E_n) < C' \left( \sum_{n=M}^N \text{Leb}(E_n) \right)^2$$

9 for all  $N$  large enough. Thus

$$\sum_{m, n = M, \dots, N} \text{Leb}(E_m \cap E_n) \leq C'' \left( \sum_{n=M}^N \text{Leb}(E_n) \right)^2$$

10 for some  $C'' > 0$  and  $N$  large enough, completing the proof.

## 11 1.12. Average of Euler's totient function.

12 **Lemma 1.29.** *For any integer  $N \geq e$ , one has*

$$\left| \sum_{n=1}^N \phi(n) - \frac{1}{2\zeta(2)} N^2 \right| \leq 5N \ln N$$

13 where  $\zeta(s) := \sum_{n \in \mathbb{Z}^+} \frac{1}{n^s}$  is the usual Riemann zeta function.

14 One may note that  $\sum \phi(n)$  is counting primitive integral vectors in a cone.

15 The proof is based on ‘‘Fubini’’, ‘‘change of variable’’ and the Mobius function:

16 **Definition 1.30.** *Decompose a positive integer  $n \neq 1$  into products of distinct prime*  
17 *numbers  $n = \prod_{i=1}^k p_i^{d_i}$  with  $d_i \in \mathbb{Z}^+$  and  $k \in \mathbb{Z}^+$ . Define the Mobius function  $\mu : \mathbb{Z}^+ \rightarrow$*   
18  *$\{-1, 0, 1\}$  by*

$$\mu(n) = \begin{cases} (-1)^k & \text{if } n \neq 1, d_i = 1 \text{ for every } i; \\ 1 & \text{if } n = 1; \\ 0 & \text{otherwise.} \end{cases}$$

19 **Lemma 1.31.** *For  $n \in \mathbb{Z}^+$ , one has*

$$\sum_{d|n} \mu(n) = \begin{cases} 0 & n \neq 1 \\ 1 & n = 1 \end{cases}$$

20 *Proof.*  $0 = (1-1)^n = \sum \binom{n}{j} (-1)^j = \sum_{d|n} \mu(n).$

□



**Lemma 1.32.**

$$\sum_{d \in \mathbb{Z}^+} \frac{\mu(d)}{d^2} \cdot \sum_{n \in \mathbb{Z}^+} \frac{1}{n^2} = 1, \text{ or equivalently, } \sum_{d \in \mathbb{Z}^+} \frac{\mu(d)}{d^2} = \zeta(2)^{-1}.$$

1 *Proof.* Expand the product and apply the lemma above. □

*Proof of Lemma 1.29.*

$$\begin{aligned} \sum_{n=1}^N \phi(n) &= \sum_{n=1}^N \sum_{m=1, \dots, n; (m,n)=1} 1 = \sum_{n=1}^N \sum_{m=1}^n \sum_{d|(m,n)} \mu(d) \\ &= \sum_{n=1}^N \sum_{d|n} \sum_{m=1, \dots, n; d|m} \mu(d) = \sum_{n=1}^N \sum_{d|n} \frac{n}{d} \mu(d) \\ &= \sum_{\{(m,d), md \leq N\}} m \mu(d) = \sum_{d=1}^N \mu(d) \sum_{m=1}^{\lfloor \frac{N}{d} \rfloor} m \\ &= \sum_{d=1}^N \mu(d) \left( \frac{1}{2} \frac{N^2}{d^2} + \text{error}_1(d) \right) \end{aligned}$$

2 where

$$|\text{error}_1(d)| = \left| \int_0^{\lfloor \frac{N}{d} \rfloor} x dx + \int_{\lfloor \frac{N}{d} \rfloor}^{\frac{N}{d}} x dx - \sum_{m=1}^{\lfloor \frac{N}{d} \rfloor} m \right| \leq \frac{1}{2} \frac{N}{d} + \frac{N}{d} \leq 2 \frac{N}{d}.$$

3 So if  $N \geq e$ ,

$$\left| \sum_{d=1}^N \text{error}_1(d) \right| \leq \sum_{d=1}^N 2 \frac{N}{d} \leq 2N \left( 1 + \sum_{d=2}^N \frac{1}{d} - \int_1^N \frac{1}{x} dx \right) \leq 2N(\ln(N) + 1) \leq 4N \ln(N).$$

4 Therefore,

$$\sum_{n=1}^N \phi(n) = \frac{N^2}{2} \sum_{d=1}^{\infty} \frac{\mu(d)}{d^2} + \text{error}_2(d)$$

5 with

$$|\text{error}_2(d)| \leq 4N \ln(N) + \frac{N^2}{2} \sum_{d=N+1}^{\infty} \frac{1}{d^2} \leq 4N \ln(N) + \frac{N^2}{2} \int_N^{\infty} \frac{1}{x^2} dx \leq 5N \ln(N)$$

6 if  $N \geq e$ . □

7

8

## REFERENCES

- 9 [Bad13] Dzmitry A. Badziahin, *On multiplicatively badly approximable numbers*, *Mathematika* **59**  
10 (2013), no. 1, 31–55. MR 3028170 3  
11 [BRV16] Victor Beresnevich, Felipe Ramírez, and Sanju Velani, *Metric Diophantine approximation: as-*  
12 *pects of recent work*, Dynamics and analytic number theory, London Math. Soc. Lecture Note  
13 Ser., vol. 437, Cambridge Univ. Press, Cambridge, 2016, pp. 1–95. MR 3618787 1  
14 [Cas50] J. W. S. Cassels, *Some metrical theorems in Diophantine approximation. I*, Proc. Cambridge  
15 Philos. Soc. **46** (1950), 209–218. MR 36787 1  
16 [EKL06] Manfred Einsiedler, Anatole Katok, and Elon Lindenstrauss, *Invariant measures and the set of*  
17 *exceptions to Littlewood’s conjecture*, Ann. of Math. (2) **164** (2006), no. 2, 513–560. MR 2247967  
18 3  
19 [Zaf17] Agamemnon Zafeiropoulos, *Inhomogeneous diophantine approximation, m\_0 sets and projec-*  
20 *tions of fractals.*, Ph.D. thesis, University of York, 2017. 1  
21 Email address: zhangrunlin@outlook.com