

Feedback — Problem Set-3

[Help Center](#)

You submitted this quiz on **Thu 5 Nov 2015 10:47 PM PST**. You got a score of **5.00** out of **5.00**.

Question 1

How many different minimum cuts are there in a tree with n nodes (ie. $n - 1$ edges) ?

Your Answer	Score	Explanation
<input type="radio"/> $\binom{n}{2}$		
<input checked="" type="radio"/> $n - 1$	✓ 1.00	Each edge defines a distinct minimum cut (with one crossing edge).
<input type="radio"/> n		
<input type="radio"/> $2^n - 2$		
Total	1.00 / 1.00	

Question 2

Let "output" denote the cut output by Karger's min cut algorithm on a given connected graph with n vertices, and let $p = \frac{1}{\binom{n}{2}}$. Which of the following statements are true? For hints on this question, you might want to watch the short optional video on "Counting Minimum Cuts".

Your Answer	Score	Explanation
<input checked="" type="checkbox"/> For every graph G with n nodes, there exists a min cut (A, B) of G such that $Pr[out = (A, B)] \geq p$.	✓ 0.20	This is even weaker than what we proved in lecture.
<input type="checkbox"/> For every graph G with n nodes and every	✓ 0.20	Recall the first problem of this

min cut (A, B) ,			problem set.
$Pr[out = (A, B)] \leq p.$			
<input type="checkbox"/> For every graph G with n nodes, there exists a min cut (A, B) such that	✓	0.20	Recall the first problem of this problem set.
$Pr[out = (A, B)] \leq p.$			
<input checked="" type="checkbox"/> For every graph G with n nodes and every min cut (A, B) of G ,	✓	0.20	This is exactly what we proved in lecture.
$Pr[out = (A, B)] \geq p.$			
<input checked="" type="checkbox"/> There exists a graph G with n nodes and a min cut (A, B) of G such that	✓	0.20	An n -cycle works.
$Pr[out = (A, B)] \leq p.$			
Total		1.00 / 1.00	

Question 3

Let $.5 < \alpha < 1$ be some constant. Suppose you are looking for the median element in an array using RANDOMIZED SELECT (as explained in lectures). What is the probability that after the first iteration the size of the subarray in which the element you are looking for lies is $\leq \alpha$ times the size of the original array?

Your Answer	Score	Explanation
<input type="radio"/> $1 - \frac{\alpha}{2}$		
<input checked="" type="radio"/> $2\alpha - 1$	✓ 1.00	Precisely, as the pivot selected should lie between $1 - \alpha$ and α times the original array.
<input type="radio"/> $\alpha - \frac{1}{2}$		
<input type="radio"/> $1 - \alpha$		
Total	1.00 / 1.00	

Question 4

Let $0 < \alpha < 1$ be a constant, independent of n . Consider an execution of RSelect in which you always manage to throw out at least a $1 - \alpha$ fraction of the remaining elements before you recurse. What is the maximum number of recursive calls you'll make before terminating?

Your Answer	Score	Explanation
<input type="radio"/> $-\frac{\log(n)}{\log(1-\alpha)}$		
<input type="radio"/> $-\frac{\log(n)}{\alpha}$		
<input type="radio"/> $-\frac{\log(n)}{\log(\frac{1}{2}+\alpha)}$		
<input checked="" type="radio"/> $-\frac{\log(n)}{\log(\alpha)}$	✓ 1.00	Equivalently, log base 1/alpha of n.
Total	1.00 / 1.00	

Question 5

The minimum s-t cut problem is the following. The input is an undirected graph, and two distinct vertices of the graph are labelled "s" and "t". The goal is to compute the minimum cut (i.e., fewest number of crossing edges) that satisfies the property that s and t are on different sides of the cut. Suppose someone gives you a subroutine for this s-t minimum cut problem via an API. Your job is to solve the original minimum cut problem (the one discussed in the lectures), when all you can do is invoke the given min s-t cut subroutine. (That is, the goal is to reduce the min cut problem to the min s-t cut problem.)

Now suppose you are given an instance of the minimum cut problem -- that is, you are given an undirected graph (with no specially labelled vertices) and need to compute the minimum cut. What is the minimum number of times that you need to call the given min s-t cut subroutine to guarantee that you'll find a min cut of the given graph?

Your Answer	Score	Explanation
<input type="radio"/> 2^n		



$n - 1$



1.00

Call an arbitrary vertex s , let t range over all other $n-1$ vertices, and return the best of the s - t min cuts founds.

☐ $\binom{n}{2}$

☐ n

Total

1.00 /

1.00