



C Shared and distinct transcriptional responses

	KLF4-OE	IGFL1
KLF4	+1.37***	+1.68***
KRT14	+7.85***	+3.48
KRT17	+3.50***	+2.39***
PDGFRA	-2.65***	-0.86***
VIM	-0.09	+0.77***
EPCAM	-1.91	-4.77**
TACSTD2	-4.74	-6.56***
TGFA	+4.51***	+4.57***
HBEGF	+2.37***	+3.97***
EGFR	+0.80**	+0.81***
DUSP6	+1.10***	+3.48***
NFKBIA	-0.90**	+0.95***
MKI67	-0.32	-2.24***
PCNA	-0.60	-1.07***
FGF7	+1.02***	-2.41***




KLF4-OE: 2 vs 2 libraries; IGFL1: 2 vs 3 libraries.

* FDR <0.05; ** <0.01; *** <0.001 (gene-wide BH).

IGFL1 raw counts in the KLF4 experiment: 0 / 0 / 0 / 0.

D Pathway-level transcriptional responses




Numbers: NES. Filled symbols: full-library FDR <0.05.

Open symbols: FDR ≥0.05; GO + KEGG corrected jointly.

IGFL1: NF-κB FDR=0.0233; cell cycle FDR=0.000409.

E Patient-level associations within Fib_K14




One point per patient (n=20); samples averaged per patient.

Gene expression: mean log1p(CP10K).

Module: KRT6A/B, KRT16/17, LAMC2, LAMB3, TACSTD2.

Spearman correlations; BH across all tested associations.

F Ligand and candidate receptor distribution



Parentheses: patients per cell state; patient-weighted means.

Fib_K14: IGFLR1 detected in 0.92% of cells.

Detection: >0 counts; ≥20 cells per sample–state group.