

# Are local-mins Rare? \*

Thomas Jefferson

## Abstract

Analyze nonconvex optimization from topology.

## 1 Test Section

This is a percent %. Should stay here after running the cleaning file.

We add a figure for testing.



If the code is correct, then the following figure should not appear in the cleaned tex file ( if the code is wrong in the sense that it only removes % but not the commands after it, then the non-included figure would appear in the cleaned tex file).

We can also check inputting other Latex Files. This is a file in a sub-folder “figures”, and we are supposed to see one figure, but not two figures.

---

\*This is an informal note.

## 2 Model

This part is a test that the standard math formulas will not be affected after running the cleaning file.

Definition (LSC property):  $\forall \bar{\alpha} \in G, \bar{x} \in L_f(\bar{\alpha}) \triangleq \{x : f(x) \leq \bar{\alpha}\}$ , and  $\forall$  sequence  $\{\alpha_i\} \rightarrow \bar{\alpha}$ , we have

$$\exists x^i \in L_f(\alpha_i), \text{ s.t. } \{x^i\} \rightarrow \bar{x}.$$

Definition (GC property): For any two points  $f(\tilde{x}) < f(\bar{x})$ , there exists a sequence  $\{x^i\} \in C$  converging to  $\bar{x}$  such that

$$f(x^i) \leq \frac{1}{i}f(\tilde{x}) + (1 - \frac{1}{i})f(\bar{x}). \quad (1)$$