

Rupadarshi's MS thesis notebook

Summer holidayrs will start from Wed 7 May, '25 and end on Mon 4 Aug, '25.

39 days left in summer holidays of 90 days

57.07% complete

MS thesis period will start from Mon 4 Aug, '25 and end on Sat 4 Apr, '26 tentatively.

39 days to go

FYI: It is a 244 days of MS thesis year.

contents

- *on $\mathbb{R}^n$*
 - Hilbert transform
- *general theory*
 - Ergodicity and mixing
 - spectral theory and ergodicity
 - Locally compact groups
 - Actions of locally compact groups
- *examples*
 - $SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$

references

- main references
 - Coifman and Weiss - Transference Methods in Analysis
 - Amos Nevo - Pointwise Ergodic Theorems for Actions of Groups
 - Einsiedler and Ward - Ergodic Theory
- secondary references
 - sub-Riem_notes - 2024 - September.pdf
- tertiary references: long-term textbooks
 - Sigurdur Helgason - Groups and geometric analysis
 - operator theory
 - Eisner, Farkas, Haase, Nagel - Operator Theoretic Aspects of Ergodic Theory-Springer (2015).pdf
- notes/lectures
 - Uri Bader - A course in linear groups and ergodic theory

order of presentations

- Presentation 1 (19 June '25)
 - singular integral transforms on $\mathbb{R}^n$, transference methods ^[1]
- Meet (26 June '25)

- Ergodic transformations, spectral invariants of ergodic transformations
- Presentation 2
 - Amenable groups, examples and equivalent definitions
- Presentation 3
 - proof of ergodicity of geodesic flow
- Presentation 4
 - proof of ergodicity of hyperbolic linear maps on torus
- Presentation 5
 - Back to Amos Nevo:
- Presentation 6
- Presentation 7

1. [Coifman and Weiss - Transference Methods in Analysis > singular integral transforms](#) sections 1 and *stated* section 2
↩

spectral theory and ergodicity

spectral theorems

 **(Spectral theorem for bounded self-adjoint operators on separable Hilbert spaces, operator-valued integration version)**  If $A \in \mathcal{B}(H)$ is **self-adjoint** then $\exists$! projection valued measure μ_A on the Borel space $\sigma(A) \subset \mathbb{C}$ with values in projections on H such that

$$\int_{\lambda \in \sigma(A)} \lambda \, \delta \mu_A = A$$

[Brian C. Hall - Quantum Theory for Mathematicians-Springer-Verlag New York \(2013\), p.141](#)

 **(Spectral theorem for bounded self-adjoint operators on separable Hilbert spaces, direct integration version)**  If $A \in \mathcal{B}(H)$ is **self-adjoint** then there $\exists$ σ -finite measure $\mu : \sigma(A) \rightarrow [0, \infty]$ and a unitary map

$$U : H \rightarrow \int_{\lambda \in \sigma(A)}^{\oplus} H_{\lambda} \, \delta \mu$$

such that $UAU^{-1}(s)(\lambda) = \lambda s(\lambda)$ for all sections s in the direct integral.

[Brian C. Hall - Quantum Theory for Mathematicians-Springer-Verlag New York \(2013\), p.147](#)

 **(Spectral theorem for bounded normal operators on separable Hilbert spaces, Hilbert bundle on the circle version)** If $A \in \mathcal{B}(H)$ is **normal** then there exists a σ -finite? Borel measure $\mu : \sigma(A) \rightarrow [0, \infty]$, a Borel measurable

$$M : \sigma(A) \rightarrow \mathbb{N} \cup \{\infty\}$$

and a unitary isomorphism

$$U : H \cong L^2(E(M) \rightarrow S^1)$$

that sends A to the operator $A(f) = \lambda f(\lambda)$ where $E(M) \rightarrow S^1$ is the Hilbert bundle with fibres satisfying

$$\begin{cases} \dim E_{\lambda} = M(\lambda) \\ E_{\lambda} \cong l^2(\mathbb{Z}) \end{cases} \quad \text{when } M(\lambda) = \infty$$

 **(Spectral theorem for bounded self-adjoint operators on separable Hilbert spaces, multiplier version)**  If $A \in \mathcal{B}(A)$ is **self-adjoint** then there exists a σ -finite measure space (X, μ) , a bounded measurable $h : X \rightarrow \mathbb{R}$ and a unitary isomorphism

$$U : (H, A) \rightarrow (L^2(X, \mu), h)$$

[Brian C. Hall - Quantum Theory for Mathematicians-Springer-Verlag New York \(2013\), p.147](#)

Theorem 18.4 (Spectral Theorem, Multiplier Form). *For a bounded, normal operator T on a Hilbert space H , the pair (H, T) is unitarily equivalent to $(L^2(\Omega, \mu), M)$, where μ is a positive Baire measure on a locally compact space Ω , and M is the multiplication operator associated with a $\sigma(T)$ -valued continuous function on Ω .*

Proof. We employ the terminology introduced above. For each α let $K_\alpha := K \times \{\alpha\}$ be a copy of $K = \sigma(T)$, so that the sets K_α are pairwise disjoint. Let $\Omega := \bigcup_\alpha K_\alpha$ with the direct sum topology and the direct sum measure $\mu := \bigoplus_\alpha \mu_{x_\alpha}$. Then we obtain

$$H = \bigoplus_\alpha Z(x_\alpha) \cong \bigoplus_\alpha L^2(K, \mu_{x_\alpha}) \cong L^2\left(\bigcup_\alpha K_\alpha, \bigoplus_\alpha \mu_{x_\alpha}\right) = L^2(\Omega, \mu).$$

Each K_α is compact and open in Ω , whence Ω is locally compact. On $Z(x_\alpha)$ the operator T acts as multiplication by z on $L^2(K, \mu_{x_\alpha})$, and hence T acts on H as multiplication on $L^2(\Omega, \mu)$ by a function m simply given by $m(z, \alpha) = z$ on K_α . This is a continuous function on Ω . $\square$

if it is not invertible). Thus for any measure-preserving transformation T , the associated operator U_T is an isometry, and if T is invertible then the associated operator U_T is a unitary operator, called the *associated unitary operator* of T or *Koopman operator* of T .

A property of a measure-preserving transformation is said to be a *spectral* or *unitary property* if it can be detected by studying the associated operator on L^2_μ .

Lemma 2.18. *A measure-preserving transformation T is ergodic if and only if 1 is a simple eigenvalue of the associated operator U_T . Hence ergodicity is a unitary property.*

PROOF. This follows from the proof of the equivalence of (2) and (5) in Proposition 2.14 or via Exercise 2.3.5 applied with $p = 2$: an eigenfunction for the eigenvalue 1 is a T -invariant function, and ergodicity is characterized by the property that the only T -invariant functions are the constants. $\square$

Example 3: Lebesgue spectrum. Consider the unitary operator U on $\ell^2(\mathbb{Z})$ defined by the left shift:

$$U(a_i) = (a_{i-1}).$$

Clearly this operator has no eigenvectors. What is its spectral measure?

Theorem 2.5 *The shift operator on $\ell^2(\mathbb{Z})$ has Lebesgue spectrum of multiplicity one.*

(This means its spectral measure (class) μ is given by Lebesgue measure $|d\lambda|$ on the unit circle S^1 .)

Proof. Observe that the map

$$(a_i) \mapsto \sum a_i z^i$$

gives an isomorphism between $\ell^2(\mathbb{Z})$ and $L^2(S^1)$. Since

$$zf(z) = z \sum a_i z_i = \sum a_{i-1} z^i,$$

this isomorphism sends U to the operator $L(f) = zf(z)$ acting on $L^2(S^1)$. Consequently the spectral measure of U is given simply by Lebesgue measure on S^1 , with multiplicity one. ■

spectrum of circle rotations

We have the unitary map $R^{\theta*}$ on $L^2(S^1)$ such that

$$R^{\theta*}(f)(z) := f(R^\theta(x))$$

On the Fourier basis $(e_k)_{k \in \mathbb{Z}}$ this acts by

$$R^{\theta*}(e_k) = \exp(2\pi i \theta)^k e_k$$

so it is diagonal with eigenvalues A_θ .

- If θ is rational, then its spectral measure is

$$\mu = \sum_{x \in A_\theta} \delta_x$$

with multiplicity $M \Big|_{A_\theta} \equiv \infty$.

- If θ is irrational then its spectral measure is

$$\mu = \sum_{x \in A_\theta} \delta_x$$

with multiplicity $M \Big|_{A_\theta} \equiv 1$

☞ **An irrational rotation of the circle has *pure point spectrum of multiplicity 1* supported on $\langle \lambda \rangle < S^1$ for some $\lambda \in S^1$ of infinite order that is either the rotation angle or its inverse. Thus, two irrational rotations by $\lambda_1, \lambda_2 \in S^1$ are isomorphic $\iff \lambda_1 = \lambda_2^\pm$.**

$SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$

[#Wiki/group-action/Lie](#)

- We observe

$$\Im \left(\frac{az + b}{cz + d} \right) = \frac{\Im(z)}{|cz + d|^2}$$

for $a, b, c, d \in \mathbb{R}$

 **Definition.** $SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$

The group $SL(2, \mathbb{R})$ acts on the upper-half plane $\mathbb{R}H^2$ by *Mobius transformations*

$$(x, y) \mapsto \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}^\top$$

for $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2, \mathbb{R})$ and $(x, y) \in \mathbb{R}H^2$, where the action by $PSL(2, \mathbb{R})$ is faithful.

group of complex 1-manifold automorphisms

- Let $x + iy \in H^2$. Then

$$\begin{bmatrix} \sqrt{y} & x/\sqrt{y} \\ 0 & 1/\sqrt{y} \end{bmatrix}$$

maps

$$i \mapsto \frac{\sqrt{y} i + \frac{x}{\sqrt{y}}}{\frac{1}{\sqrt{y}}} = x + iy$$

- Thus the action $SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$ is **transitive**.
- Now, the stabilizer of $i \in \mathbb{R}H^2$ must have

$$|ci + d|^2 = 1$$

by

- We observe

$$\Im \left(\frac{az + b}{cz + d} \right) = \frac{\Im(z)}{|cz + d|^2}$$

for $a, b, c, d \in \mathbb{R}$

- That means $(c, d) = (\sin \theta, \cos \theta)$ for some $\theta \in \mathbb{R}$, but

$$\begin{aligned} \frac{ai + b}{\sin \theta i + \cos \theta} &= i \\ \implies a, b &= \cos \theta, -\sin \theta \end{aligned}$$

- Thus

$$\text{stab}_{SL(2, \mathbb{R})}(i) = SO(2, \mathbb{R})$$

 The transitive, faithful action $PSL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$ by *Mobius transformations* is the connected component of isometry group of the hyperbolic metric on upper-half plane

$$\mathrm{AutMan}_O(\mathbb{R}H^2, g_{\mathrm{Hy}})_0 \cong PSL(2, \mathbb{R})$$

Therefore the action is symplectic or area preserving in the hyperbolic area 2-form.

Thus, we have

$$SL(2, \mathbb{R}) \curvearrowright (\mathbb{R}H^2, g_{\mathrm{Hy}}, \lambda_{\mathrm{Hy}})$$

Lie generators

For the transitive action $SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$ by *Mobius transformations*, the stabilizer of $i = (0, 1) \in H$ is $SO(2, \mathbb{R}) < SL(2, \mathbb{R})$. Thus

$$\frac{SL(2, \mathbb{R})}{SO(2, \mathbb{R})} \cong_{\mathrm{Man}} \mathbb{R}H^2$$

by orbit-stabilizer theorem.

sub-actions

- how subgroups act
 - $SL(2, \mathbb{Z}) \curvearrowright \mathbb{R}H^2$
 - Γ torsion-free discrete
 - ...

induced action on $T^\cup H^2$

$$A(z) = \frac{az + b}{cz + d}$$

then

- its *complex* derivative is

$$\begin{aligned} A'(z) &= \frac{a(cz + d) - c(az + b)}{(cz + d)^2} \\ &= \frac{\det A}{(cz + d)^2} \\ &= \frac{1}{(cz + d)^2} \end{aligned}$$

- which is precisely the multiplier for the linear map $\mathfrak{D}_z A$, making the the derivative

$$\begin{aligned} \mathfrak{D}A : T\mathbb{R}H^2 &\rightarrow T\mathbb{R}H^2 \\ (z, v) &\mapsto \left(\frac{az + b}{cz + d}, \frac{v}{(cz + d)^2} \right) \end{aligned}$$

where $T\mathbb{R}H^2 \cong \mathbb{R}H^2 \times U(1)$ naturally.

- The action preserves the hyperbolic metric because

$$\begin{aligned} \langle D_z A(v), D_z A(w) \rangle_{\mathrm{Hyp}} &= \left(\frac{y}{|cz + d|^2} \right)^{-2} \left(\frac{v}{(cz + d)^2}, \frac{w}{(cz + d)^2} \right) \\ &= \frac{1}{y^2} \langle v, w \rangle_{\mathrm{DOT}} \\ &= \langle v, w \rangle_{\mathrm{Hyp}} \end{aligned}$$

- As the inner product is preserved, hyperbolic unit vectors map to themselves, thus we restrict the action to

$$SL(2, \mathbb{R}) \curvearrowright T^u \mathbb{R}H^2$$

- Now consider the action of the stabilizers

$$A = \frac{i \cos \theta - \sin \theta}{i \sin \theta + \cos \theta}$$

of $i \in \mathbb{R}H^2$

$$(D_i A)(v) = \frac{v}{(i \sin \theta + \cos \theta)^2} = (\cos 2\theta - i \sin 2\theta)v$$

- So varying θ , we can map v to any element of $T_i^u \mathbb{R}H^2$.
- Thus, the action

$$SL(2, \mathbb{R}) \curvearrowright T^u \mathbb{R}H^2$$

- Now, say A is a stabilizer of (i, v) , then

$$\begin{aligned} 2\theta \bmod 2\pi &= 0 \\ \implies \theta &= \pi \mathbb{Z} \\ \implies A &= \pm I \end{aligned}$$

- Thus, the action

$$PSL(2, \mathbb{R}) \curvearrowright T^u \mathbb{R}H^2$$

is **transitive and free** giving us an isomorphism

$$\begin{aligned} PSL(2, \mathbb{R}) &\cong T^u \mathbb{R}H^2 \\ A &\mapsto A(i, i) \end{aligned}$$

 **Definition.** $SL(2, \mathbb{R}) \curvearrowright T\mathbb{R}H^2 \cong \mathbb{R}H^2 \times U(1)$

We have the action

$$\begin{aligned} SL(2, \mathbb{R}) &\curvearrowright T\mathbb{R}H^2 \cong \mathbb{R}H^2 \times U(1) \\ (z, v) &\xrightarrow{\begin{bmatrix} a & b \\ c & d \end{bmatrix}} \left(\frac{az + b}{cz + d}, \frac{v}{(cz + d)^2} \right) \end{aligned}$$

induced by $SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$ where the **action by $PSL(2, \mathbb{R})$ is free and transitive** giving us an isomorphism of left $PSL(2, \mathbb{R})$ -manifolds

$$\begin{aligned} PSL(2, \mathbb{R}) &\cong T^u \mathbb{R}H^2 \\ A &\mapsto A(i, i) \end{aligned}$$

geodesic flow

We have the geodesic flow

$$\exp(t-): T^u \mathbb{R}H^2 \rightarrow T^u \mathbb{R}H^2$$

of the hyperbolic metric on $\mathbb{R}H^2$

- We have

$$\exp(t(i, i)) = (e^t i, e^t i) = \mathfrak{D}_i \begin{bmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{bmatrix} (i)$$

- For the geodesic starting at $(z, v) = A(i, i)$ we have

$$\begin{aligned} \exp(t(z, v)) &= \mathfrak{D} A(\exp(t(i, i))) \\ &= \mathfrak{D} A\left(\mathfrak{D}_i \begin{bmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{bmatrix} (i)\right) \\ &= \mathfrak{D}_i \left(A \begin{bmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{bmatrix}^{-1} \right) (i) \end{aligned}$$

- Thus, under the identification

 **Definition.** $SL(2, \mathbb{R}) \curvearrowright T\mathbb{R}H^2 \cong \mathbb{R}H^2 \times U(1)$

We have the action

$$\begin{aligned} &SL(2, \mathbb{R}) \curvearrowright T\mathbb{R}H^2 \cong \mathbb{R}H^2 \times U(1) \\ &(z, v) \xrightarrow{\begin{bmatrix} a & b \\ c & d \end{bmatrix}} \left(\frac{az + b}{cz + d}, \frac{v}{(cz + d)^2} \right) \end{aligned}$$

induced by $SL(2, \mathbb{R}) \curvearrowright \mathbb{R}H^2$ where the action by $PSL(2, \mathbb{R})$ is free and transitive giving us an isomorphism of left $PSL(2, \mathbb{R})$ -manifolds

$$\begin{aligned} PSL(2, \mathbb{R}) &\cong T^u\mathbb{R}H^2 \\ A &\mapsto A(i, i) \end{aligned}$$

the geodesic flow corresponds to right multiplication by $\begin{bmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{bmatrix}^{-1}$ that is

$$A \xrightarrow{t} A \begin{bmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{bmatrix}^{-1}$$

Coifman and Weiss - Transference Methods in Analysis (1977)

Definition. Hilbert transform on $\mathbb{R}$

For $p \in [0, \infty]$, the **Hilbert transform** is the operator

$$H : L^p(\mathbb{R}) \rightarrow L^p(\mathbb{R})$$
$$f \mapsto Hf := \lim_{\epsilon \rightarrow 0+} \underbrace{\frac{1}{\pi} \int_{\epsilon \leq |t|} \frac{f(s-t)}{t}}_{H_\epsilon f}$$

- for $p = 1$, this operator is weak- L^1 bounded
- for $p > 1$, this operator is bounded

Proposition: For the Hilbert transform

Definition. Hilbert transform on $\mathbb{R}$

For $p \in [0, \infty]$, the **Hilbert transform** is the operator

$$H : L^p(\mathbb{R}) \rightarrow L^p(\mathbb{R})$$
$$f \mapsto Hf := \lim_{\epsilon \rightarrow 0+} \underbrace{\frac{1}{\pi} \int_{\epsilon \leq |t|} \frac{f(s-t)}{t}}_{H_\epsilon f}$$

if $1 < p < \infty$ we have

$$f \in L^p(\mathbb{R}) \implies \|H_\epsilon f\|_p \leq A_p \|f\|_p$$

singular integral transforms

Let $h \in L^1(\mathbb{R}^n)$. Then

$$\int_{\mathbb{R}^n} h = \int_{\hat{x} \in S^{n-1}} \left(\int_{r \in (0, \infty)} h(r\hat{x}) r^{n-1} \right)$$

Definition. Singular integral transform in $\mathbb{R}^n$ with homogeneous odd kernels

Let

$$k(y) := \frac{\Omega(\hat{y})}{|y|^n}$$

where $\Omega \in L^1(S^{n-1})$ and $\Omega(-\hat{y}) = -\Omega(\hat{y})$.

Then

$$\begin{aligned} (H_k^\epsilon f)(x) &:= \int_{\epsilon \leq |y| \leq \frac{1}{\epsilon}} k(y) f(x-y) \\ &= \int_{\hat{y} \in S^{n-1}} \Omega(\hat{y}) \int_{r \in (\epsilon, 1/\epsilon)} \frac{f(x-r\hat{y})}{r^n} r^{n-1} \\ &= \int_{\hat{y} \in S^{n-1}} \Omega(\hat{y}) \int_{r \in (\epsilon, 1/\epsilon)} \frac{f(x-r\hat{y})}{r} \end{aligned}$$

and

$$H_k f := \lim_{\epsilon \rightarrow 0} H_k^\epsilon$$

We define the transformations

$$\begin{aligned} U_{\hat{y}}^t &: \mathbb{R}^n \rightarrow \mathbb{R}^n \\ U_{\hat{y}}^t(x) &:= x + t\hat{y} \end{aligned}$$

for $\hat{y} \in S^{n-1}$ and $t \in \mathbb{R}$.

For

$$f : \mathbb{R}^n \rightarrow \mathbb{C}$$

we have

$$\begin{aligned} g_{\hat{y}}^x &: \mathbb{R} \rightarrow \mathbb{C} \\ g_{\hat{y}}^x(t) &:= f(U_{\hat{y}}^t(x)) = f(x + t\hat{y}) \end{aligned}$$

Proposition:

$$2(H_k^\epsilon f)(x) = \int_{\hat{y} \in S^{n-1}} (H_\epsilon g_{\hat{y}}^x)(0) \Omega(\hat{y})$$



Before describing this general situation, however, let us establish (1.6). We shall use polar coordinates in order to express certain integrals over $\mathbb{R}^n$: if $h \in L^1(\mathbb{R}^n)$ and $x = rx' \in \mathbb{R}^n$, where $x' \in \Sigma_{n-1}$ and $r = |x| \geq 0$, then

$$(1.8) \qquad \int_{\mathbb{R}^n} h(x) dx = \int_{\Sigma_{n-1}} \left\{ \int_0^\infty h(rx') r^{n-1} dr \right\} dx'$$

(dx' is the element of Lebesgue measure on the surface Σ_{n-1}). Thus, since $\Omega(ry') = \Omega(y')$ (Ω being homogeneous of degree 0)

$$\begin{aligned} 2(H_k^\epsilon f)(x) &= 2 \int_{\epsilon \leq |y| \leq 1/\epsilon} k(y) f(x-y) dy \\ &= 2 \int_{\Sigma_{n-1}} \Omega(y') \left\{ \int_\epsilon^{1/\epsilon} \frac{f(x-ry')}{r^n} r^{n-1} dr \right\} dy'. \end{aligned}$$

Since $\int_{\Sigma_{n-1}} \Phi(y') dy' = \int_{\Sigma_{n-1}} \Phi(-y') dy'$, whenever $\Phi \in L^1(\Sigma_{n-1})$, and since $\Omega(-y') = -\Omega(y')$ (by assumption), it follows that the last expression equals

$$\int_{\Sigma_{n-1}} \Omega(y') \left\{ \int_{\epsilon \leq |r| \leq 1/\epsilon} \frac{f(x-ry')}{r} dr \right\} dy' = \int_{\Sigma_{n-1}} \Omega(y') (H_\epsilon g_{y'}^x)(0) dy'$$

and (1.6) is established.



Proposition:

$$\|H_k^\epsilon f\|_p \leq B_p \|f\|_p$$

From

Proposition:

$$2(H_k^\epsilon f)(x) = \int_{\hat{y} \in S^{n-1}} (H_\epsilon g_{\hat{y}}^x)(0) \Omega(\hat{y})$$

$$\begin{aligned}
2\|H_k^\epsilon f\|_p &= \left(\int_{x \in \mathbb{R}^n} \left| \int_{\hat{y} \in S^{n-1}} (H_\epsilon g_{\hat{y}}^x)(0) \Omega(\hat{y}) \right|^p \right)^{1/p} \\
&\leq \int_{\hat{y} \in S^{n-1}} \left(\int_{x \in \mathbb{R}^n} |(H_\epsilon g_{\hat{y}}^x)(0) \Omega(\hat{y})|^p \right)^{1/p}
\end{aligned}$$

- Now the integral inside is for a fixed $\hat{y} \in S^{n-1}$, so we write

$$x = s\hat{y} + w$$

for $w \in T_{\hat{y}}S^{n-1} \hookrightarrow \mathbb{R}^n$ and split the integral

$$\begin{aligned}
&\int_{x \in \mathbb{R}^n} |(H_\epsilon g_{\hat{y}}^x)(0) \Omega(\hat{y})|^p \\
&= |\Omega(\hat{y})|^p \int_{w \in T_{\hat{y}}S^{n-1}} \int_{s \in \mathbb{R}} |(H_\epsilon g_{\hat{y}}^x)(0)|^p \\
&= |\Omega(\hat{y})|^p \int_{w \in T_{\hat{y}}S^{n-1}} \int_{s \in \mathbb{R}} \left| \int_{\epsilon \leq |t| \leq \frac{1}{\epsilon}} \frac{f((s-t)\hat{y} + w)}{t} \right|^p \\
&\leq |\Omega(\hat{y})|^p \int_{w \in T_{\hat{y}}S^{n-1}} A_p^p \int_{s \in \mathbb{R}} |f(s\hat{y} + w)|^p \\
&\leq A_p^p |\Omega(\hat{y})|^p \|f\|_{L^p(\mathbb{R}^n)}^p
\end{aligned}$$

- Here, we took the estimate from one dimension

Proposition: For the Hilbert transform

📌 **Definition. Hilbert transform on $\mathbb{R}$**

For $p \in [0, \infty]$, the **Hilbert transform** is the operator

$$\begin{aligned}
H : L^p(\mathbb{R}) &\rightarrow L^p(\mathbb{R}) \\
f &\mapsto Hf := \lim_{\epsilon \rightarrow 0^+} \underbrace{\frac{1}{\pi} \int_{\epsilon \leq |t|} \frac{f(s-t)}{t}}_{H_\epsilon f}
\end{aligned}$$

if $1 < p < \infty$ we have

$$f \in L^p(\mathbb{R}) \implies \|H_\epsilon f\|_p \leq A_p \|f\|_p$$

applied it over lines $\{s\hat{y} + w \mid s \in \mathbb{R}\}$ in $\mathbb{R}^n$ and then averaged it over all $w \perp \hat{y}$: so averaging it over *all lines* in $\mathbb{R}^n$ in the direction $\hat{y}$.

- Thus

$$2\|H_k^\epsilon f\|_p \leq \left(\int_{\hat{y} \in S^{n-1}} |\Omega(\hat{y})| \right) A_p \|f\|_{L^p(\mathbb{R}^n)}$$

- This further averaged it over all $\hat{y} \in S^{n-1}$, essentially averaging the result over *all lines* in $\mathbb{R}^n$.
- Hence

$$B_p = \frac{1}{2} A_p \|\Omega\|_{L^1(S^{n-1})}$$

concludes the result.

on the circle

$$\mathbb{R} \curvearrowright S^1$$

by isometries

general transference results

 **Definition.** Continuous representation of an amenable group on L^p space of a σ -finite space

Let G is a locally compact group with the *amenable* property: for any compact $C \subset G$ and $\epsilon > 0$ there exists an open nbd V of identity having finite left Harr measure $\nu(V)$ such that

$$\frac{\nu(VC^{-1})}{\nu(V)} \leq 1 + \epsilon$$

Let

$$R : G \rightarrow \mathcal{B}(L^p(X, \mu))$$

be a continuous representation of G for (X, μ) a σ -finite measure space.

Let $k \in L^1(G)$ have compact support C and $N_p(k)$ the operator norm of the convolution operator $\|k * -\|_{L^p(G) \rightarrow L^p(G)}$ for $p \geq 1$.

We have the ***R-convolution with k*** operator

$$\begin{aligned} H_k : L^p(X, \mu) &\rightarrow L^p(X, \mu) \\ F &\mapsto (H_k F)(x) := \int_{u \in G} k(u) (R_{u^{-1}} F)(x) \end{aligned}$$

 For

 **Definition.** Continuous representation of an amenable group on L^p space of a σ -finite space

Let G is a locally compact group with the *amenable* property: for any compact $C \subset G$ and $\epsilon > 0$ there exists an open nbd V of identity having finite left Harr measure $\nu(V)$ such that

$$\frac{\nu(VC^{-1})}{\nu(V)} \leq 1 + \epsilon$$

Let

$$R : G \rightarrow \mathcal{B}(L^p(X, \mu))$$

be a continuous representation of G for (X, μ) a σ -finite measure space.

Let $k \in L^1(G)$ have compact support C and $N_p(k)$ the operator norm of the convolution operator $\|k * -\|_{L^p(G) \rightarrow L^p(G)}$ for $p \geq 1$.

We have the ***R-convolution with k*** operator

$$\begin{aligned} H_k : L^p(X, \mu) &\rightarrow L^p(X, \mu) \\ F &\mapsto (H_k F)(x) := \int_{u \in G} k(u) (R_{u^{-1}} F)(x) \end{aligned}$$

let there exists an uniform operator bound: $\exists c > 0$ such that

$$\|R_u F\| \leq c \|F\|_p$$

for all $F \in L^p(X, \mu), u \in G$. Then the operator H_k is well-defined and is bounded with

$$\|H_k\|_p \leq c^2 N_p(k)$$

Amos Nevo - Pointwise Ergodic Theorems for Actions of Groups (2006)

[Handbook of Dynamical Systems 1, Part B - Hasselblatt and A. Katok \(Eds.\) - \(2006\).pdf > page=884](#)

averaging operators of measurable preserving group actions

📌 Definition. LC2 σ -Borel G -space X

Let

- G be a locally compact 2-countable group (LC2)
- X LC2 space on which

$$G \curvearrowright X$$

acts by a group of homomorphisms

- $m_{X/G}$ be a G -invariant σ -finite Borel measure on X

This gives rise

- to a representation

$$\begin{aligned} \pi : G &\curvearrowright L^p(X, m_{X/G}) \\ f &\mapsto f \circ g^{-1} \end{aligned}$$

by isometries of $L^p(X)$

- for $p \in [1, \infty)$

$$G \rightarrow \text{Iso}(L^p(X))$$

is strongly continuous

- For any Borel probability measure μ on G we have **averaging operator**

$$\begin{aligned} P_g &:= \int_G \pi_g(-) : L^p(X, m_{X/G}) \rightarrow L^p(X, m_{X/G}) \\ f &\mapsto \int_G f(g^{-1}x) \delta\mu_g \\ &= \int_G (\pi_g f)(x) \delta\mu_g \end{aligned}$$

for $p \in [1, \infty]$, $g \in M$.

- $P_g(g)$ is G -invariant?

This is reminiscent of such operators in finite group representation theory!

regular representation

Let $X = G$ and the measure $m = \mu_G$ is the left-invariant Harr measure.

2.1.1. The regular representation and the action by convolutions. Consider the case where $X = G$, and the measure $m = m_G$ is left-invariant Haar measure. Then the operators $\lambda(g)f(x) = f(g^{-1}x)$ are isometric in every $L^p(G, m_G)$, and $g \mapsto \lambda(g)$ is the left regular representation. If μ is absolutely continuous with density $d\mu = b(g) dm_G(g)$, then the operator $\lambda(\mu)$ is the operator of left convolution by μ :

$$\lambda(\mu)f(x) = \int_G f(g^{-1}x)b(g) dm_G(g) = \mu * f(x).$$

We can similarly consider the right regular representation of G , given by $\rho(g)f(x) = f(xg)$. Of course, in general $\rho(g)$ is *not* an isometric operator in $L^p(G, m_G)$, unless the left Haar measure m_G is also right-invariant, namely unless G is unimodular. Note also that the operator $\rho(\mu)$ is given by $\rho(\mu)f(x) = \int_G f(xg)b(g) dm_G(g)$, and is equal to the convolution operator $f * \mu^\vee$ if and only if G is unimodular (here $\mu^\vee(A) = \mu(A^{-1})$). In particular if G is unimodular and the measure μ is symmetric (namely satisfies $\mu^\vee = \mu$), then $\rho(\mu)f = f * \mu$.

ergodic theorems

 **Definition.** Family of averaging operators on a probability G -space

Let G be a LC2 group, $(X, \mathfrak{M}, m)$ be a *probability space*

$$G \rightarrow \text{AutMeas}(X, \mathfrak{M}, m)$$

be a *group homomorphism*.

Let $\mu_t, t \in \mathbb{R}_+$ be a family of *probability measures* on G such that

$$\begin{aligned} \mu : \mathbb{R}_+ &\rightarrow M(G) \cong \mathcal{C}_0(G)^* \\ t &\mapsto \mu_t \end{aligned}$$

is w^* -continuous. These are regarded as a **family of averaging operators**

$$(\pi(\mu_t)f)(x) := \int_{g \in (G, \mu_t)} f(g^{-1}x)$$

producing a *sampling method* along the orbits Gx .

Usually each μ_t will have some compact support $A_t \subset G$.

Question

We want to establish a *pointwise ergodic theorem* in L^p for families of averaging operators μ_t on G for an ergodic action on X , that is

$$\lim_{t \rightarrow \infty} \pi(\mu_t)f(x) = \int_{(X, m)} f$$

for m -almost every $x \in X$ for $f \in L^p(X, m), 1 \leq p < \infty$.

Bug

The maximal function associated with a [family of averaging operators](#) is "defined" by

$$f \mapsto (M_\mu^* f)(x) := \sup_{t \in \mathbb{R}_+} |\pi(\mu_t) f(x)|$$

growth type of groups

[1]

Let G be a compactly generated, LC2 group, that is

$$V \subset G$$

is a compact set such that

$$\bigcup_{n \in \mathbb{N}} V^n = G$$

where $V^0 := \{i_G\}$.

- Clearly

$$A, B \subset AB \subset G$$

thus

$$k \in \mathbb{N} \implies V^n \subset V^{n+k} = V^n V^k$$

We define the distance

$$|g|_V := \min\{n \mid g \in V^n\}$$

The distance function is inversion invariant $\iff V$ is symmetric $V = V^{-1} \iff$ the function $d_V(g, h) := |g^{-1}h|_V$ is symmetric in the pair (h, g) . So we assume $V = V^{-1}$. Then

$$d_V : G \times G \rightarrow \mathbb{R}$$

is a left G -invariant metric on G called the **word metric** defined by V .

- G is LC2 $\implies$ there is some $n \in \mathbb{N}$ such that V^n contains an open set
- by

- Clearly

$$A, B \subset AB \subset G$$

thus

$$k \in \mathbb{N} \implies V^n \subset V^{n+k} = V^n V^k$$

we have

$$m_G(V^{n+k}) > 0$$

and

$$m_G(V^{n+m}) = m_G(V^n V^m) \leq ? m_G(V^n) m_G(V^m)$$

Pansu's 1983 theorem

In these examples the homogeneity of the metric on G , namely the fact that $B_t = \alpha_t(B_1)$ (where α_t is the dilation automorphism group) immediately implies that $m_G(B_t) = m_G(B_1)t^q$, where q is the homogeneous dimension. Thus in particular the polynomial volume growth for homogeneous metrics is exact. Riemannian metrics are not homogeneous in general, and here, under one further assumption, the following result was established by P. Pansu.

THEOREM 4.4 [123]. *Let G be a simply connected nilpotent Lie group admitting a lattice subgroup Γ . Then*

- (1) *G has exact polynomial volume growth w.r.t. any Riemannian metric on G which is Γ -invariant.*
- (2) *Any lattice Γ in G has exact polynomial volume growth with respect to any word metric.*

Pansu's theorem has the following two consequences [123] (see also the discussion in [42, Chapter VII.C]).

COROLLARY 4.5.

- (1) *Every discrete group containing a finitely generated nilpotent group of finite index has exact polynomial volume growth with respect to any word metric. Indeed, by a well-known result of Malcev, Γ can be embedded as a lattice in a simply connected nilpotent Lie group.*
- (2) *Every discrete group Γ with polynomial volume growth has exact polynomial volume growth with respect to any word metric. Indeed, by Gromov's theorem [63], Γ is virtually nilpotent, namely contains a nilpotent subgroup of finite index.*

We note that a sharper form of Pansu's theorem has been established by M. Stoll [137], namely that $|\#B_t - Ct^q| \leq Bt^{q-1}$ for 2-step nilpotent groups.

- [123] <https://www.cambridge.org/core/services/aop-cambridge-core/content/view/9352E615E18472BAE919B24621420F9B/S0143385700002054a.pdf/croissance-des-boules-et-des-geodesiques-fermees-dans-les-nilvarietes.pdf>

- [Croissance des boules et des géodésiques fermées dans les nilvariétés | Ergodic Theory and Dynamical Systems | Cambridge Core](#)

! [15] missing??

- generalization, 2007? - <https://arxiv.org/pdf/0704.0095#page=2.0>
- asymptotics, 2012 - <https://arxiv.org/pdf/1204.1613#page=2.05>
- [137] <https://sci-hub.se/10.1112/s0024610798006371>
- <https://ep455.user.srcf.net/pdfs/nilpoly.pdf>
- [42] (Chicago Lectures in Mathematics) Pierre de la Harpe - Topics in Geometric Group Theory- The University of Chicago Press (2000), p.204
- (Grundlehren der mathematischen Wissenschaften) Martin R. Bridson, André Häfliger - Metric Spaces of Non-Positive Curvature -Springer (2009), p.95
- (Colloquium publications (American Mathematical Society) 63) Bogdan, Nica, Druțu, Cornelia, Kapovich, Michael - Geometric group theory-American Mathematical Society (2018), p.562
- (Modern Birkhäuser Classics) Mikhail Gromov (auth.) - Metric Structures for Riemannian and Non-Riemannian Spaces-Birkhäuser Basel (2007), p.103
- https://www.imo.universite-paris-saclay.fr/~pierre.pansu/Pansu_Pisa_jun23_minicourse_beamer.pdf
- https://cvgmt.sns.it/media/doc/paper/5339/sub-Riem_notes.pdf#page=229.65
 - [Topics in geometry: Nilpotent groups and subriemannian geometry](#)
 - [SubRiemannian geometry course - YouTube](#)

Theorem 11.4.14 (Quantitative Pansu Theorem; [BL13]). *Let G be a simply connected s -step nilpotent Lie group equipped with a left-invariant sub-Finsler metric ρ_1 . With respect to some compatible linear grading, consider the Pansu limit metric ρ_0 on G and the contracted metrics ρ_ϵ as in (11.4.6). For every compact set $K \subseteq G$ there is a constant $C > 0$ such that*

$$|\rho_\epsilon(p, q) - \rho_0(p, q)| \leq C\epsilon^{1/s}, \quad \forall p, q \in K, \forall \epsilon \in (0, 1). \quad (11.4.15)$$

11.4.1 Pansu asymptotic structure

Given the nilpotent Lie algebra $\mathfrak{g}$, we consider the associated Carnot algebra $\mathfrak{g}_\infty$ of G , as in Definition 8.2.25. This is a stratifiable algebra, and $\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ is the first stratum of a stratification. Consider G_∞ to be the nilpotent simply connected Lie group with Lie algebra $\mathfrak{g}_\infty$, by recalling the existence by Birkhoff Theorem; see Proposition 8.4.7. We shall put a Carnot metric on it. Consider the projection $\pi : \mathfrak{g} \rightarrow \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ modulo $[\mathfrak{g}, \mathfrak{g}]$. Since Δ_{1_G} generates the whole Lie algebra and since $\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ is an abelian Lie algebra, we have $\pi(\Delta_{1_G}) = \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$. We define a norm $\|\cdot\|_\infty$, on $\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$, which we name *Pansu limit norm*, by imposing $B_{\|\cdot\|_\infty}(0, 1) = \pi(B_{\|\cdot\|}(0, 1))$. Let d_∞ be the subFinsler distance associated with $(G_\infty, \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}], \|\cdot\|_\infty)$. Thus (G_∞, d_∞) is a Carnot group. We call d_∞ the *Pansu limit metric* or the *asymptotic metric*.

Pansu Theorem 11.1.8 will follow from a more quantitative version as obtained in [BL13]:

Theorem 11.4.1 ([Pan83, BL13]; see Theorem 11.4.14). *Let (G, d) be a nilpotent simply connected Lie group equipped with a sub-Finsler left-invariant metric. Then, the associated Carnot group G_∞ equipped with the Pansu limit metric d_∞ is the asymptotic cone of (G, d) and there is a set-wise identification between G and G_∞ such that*

$$\left| d(p, q) - d_\infty(p, q) \right| = O\left(\max\{d(1, p), d(1, q)\}^{1-1/s}\right), \quad \text{as } p, q \rightarrow \infty, \quad (11.4.2)$$

where s denotes the nilpotency step of G .

Theorem 11.1.8 (Pansu; [Pan83]; see Theorem 11.4.1). *Let G be a nilpotent simply connected Lie group equipped with a left-invariant sub-Finsler distance. Then, the asymptotic cone of G exists and is a sub-Finsler Carnot group.*

appendix

[2]

-
1. <https://www.math.uchicago.edu/~may/VIGRE/VIGRE2009/REUPapers/Lim.pdf> ↩
 2. <https://arxiv.org/pdf/2105.02090> ↩

Einsiedler and Ward - Ergodic Theory

#book/series

- [Einsiedler and Ward - Ergodic Theory \(2011\).pdf](#)
- [Entropy | Books](#)
- [Homogeneous | Books](#)

Sigurdur Helgason - Groups and geometric analysis

§1. Harmonic Analysis on Homogeneous Spaces

1. General Problems

Let X be a locally compact Hausdorff space acted on transitively by a locally compact group G . We assume G leaves invariant a positive measure μ on X . Let T_x denote the (unitary) representation of G on the Hilbert space $L^2(X)$ defined by $(T_x(g)f)(x) = f(g^{-1} \cdot x)$ for $g \in G$,

Chapter 1: Integral geometry and Radon transforms

Invariant measures on coset spaces

Definition. Invariant forms

Let M be a smooth manifold and

$$\Phi : M \rightarrow M$$

be a diffeomorphism. A differential form $\omega \in \Omega(M)$ is **Φ -invariant** if

$$\Phi^* \omega = \omega$$

Definition. Left and right invariant forms on a Lie group

Let G be a Lie group with Lie algebra $\mathfrak{g}$. A differential form $\omega \in \Omega(G)$ is **left-invariant** if

$$x \in G \implies L_x^* \omega = \omega$$

and **right-invariant** if

$$x \in G \implies R_x^* \omega = \omega$$

Let $X_1, \dots, X_n$ be a basis of $\mathfrak{g}$ (left-invariant vector fields). Then we have n many 1-forms

$$\omega_1, \dots, \omega_n$$

such that

$$\omega_i(X_j) = \delta_{ij}$$

These 1-forms are left-invariant and thus

$$\lambda := \omega_1 \wedge \dots \wedge \omega_n$$

is also a left-invariant n -form on G and it is the unique left-invariant n -form on G up to a constant factor.

Radon transform on $\mathbb{R}^n$

Let

$$P(\mathbb{R}^{n*}) \cong \mathbb{R}P^{n-1}$$

be the compact smooth $n - 1$ -manifold of all hyperplanes in $\mathbb{R}^n$ passing through origin.

Each hyperplane ξ can be written

$$\xi = \{x \in \mathbb{R}^n \mid \langle x, \hat{n} \rangle = \alpha\}$$

This produces a *surjective* map onto the space of all hyperplanes $\boldsymbol{Pl}^n$ in $\mathbb{R}^n$

$$\begin{aligned} S^{n-1} \times \mathbb{R} &\rightarrow \boldsymbol{Pl}^n \\ (\hat{n}, \alpha) &\mapsto \xi = \{x \in \mathbb{R}^n \mid \langle x, \hat{n} \rangle = \alpha\} \end{aligned}$$

that is a smooth double covering(?) as

$$(\hat{n}, \alpha), (-\hat{n}, -\alpha)$$

produce the same plane. Thus we identify

$$\mathcal{C}^\bullet(\boldsymbol{Pl}^n) \leftrightarrow \{f \in \mathcal{C}^\bullet(S^{n-1} \times \mathbb{R}) \mid f(\hat{n}, \alpha) = f(-\hat{n}, -\alpha)\}$$

📌 Definition. Radon transform and its dual

Let

$$f : \mathbb{R}^n \rightarrow \mathbb{C}$$

be integrable on each hyperplane in $\mathbb{R}^n$. Let $\boldsymbol{P}^n$ denote the compact smooth n -manifold of all hyperplanes in $\mathbb{R}^n$. The **Radon transform** of f is

$$\begin{aligned} J(f) : \boldsymbol{P}^n &\rightarrow \mathbb{C} \\ \xi &\mapsto \hat{J}(f)(\xi) := \int_{(\xi, m_{\mathbb{R}^2})} f \end{aligned}$$

and the **dual Radon transform** of $\phi \in \mathcal{C}(\boldsymbol{P}^n)$

$$\begin{aligned} \mathbb{R}^n &\rightarrow \mathbb{C} \\ x &\mapsto \check{J}(\phi)(x) := \int_{x \in \xi} \phi \end{aligned}$$

Uri Bader - A course in linear groups and ergodic theory

$$SL(2, \mathbb{R}) \curvearrowright \mathbb{R}^2$$

Consider

$$\underbrace{SL(2, \mathbb{R})}_{=: G} \curvearrowright \underbrace{\mathbb{R}^2}_{=: X}$$

There are two orbits

$$\{0\}$$

and

$$\mathbb{R}^2 \setminus \{0\}$$

(disjoint of course).

By orbit-stabilizer theorem

$$\mathbb{R}^2 \setminus \{0\} \cong \frac{G}{\mathcal{U}}$$

where stabilizer is the subgroup

$$\mathcal{U} := \text{stab}_G \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \left\{ \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \right\} = \exp \left\{ \begin{bmatrix} 0 & * \\ 0 & 0 \end{bmatrix} \right\} \cong (\mathbb{R}, +)$$

$SL(2, \mathbb{R})$ -maps

 **Question**

Find all G -invariant maps

$$\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

that is

$$\forall g \in G, x \in \mathbb{R}^2, \phi(gx) = g\phi(x)$$

Because G -orbits will go to G -orbits under such a map we conclude

$$\phi(0) = 0$$

But if we choose where one point of a orbit will go, then we know where the other elements will go

$$\underbrace{g\phi(x)}_{\text{known}} = \underbrace{\phi(gx)}_{\text{unkown}}$$

But to know

$$\phi \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

we observe


$$g \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \implies \phi \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \phi(g \begin{pmatrix} 1 \\ 0 \end{pmatrix}) = g\phi \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

means

$$\mathcal{U} \subset \text{stab}_G(\phi \begin{pmatrix} 1 \\ 0 \end{pmatrix})$$

So we must look at *the orbit structure of $\mathcal{U} \curvearrowright \mathbb{R}^2$* to know what point does it *stabilizes*:

$$\begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix} \in \mathcal{U} \implies \begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix} + \alpha \begin{pmatrix} y \\ 0 \end{pmatrix}$$



✓ [flow](#)

✓

The dynamics is "tame".

So

$$\mathcal{U} \subset \text{stab}_G(\phi \begin{pmatrix} 1 \\ 0 \end{pmatrix}) \iff \phi \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \alpha \\ 0 \end{pmatrix}$$

Then

$$\phi(y) = \phi(g \begin{pmatrix} 1 \\ 0 \end{pmatrix}) = g\phi \begin{pmatrix} 1 \\ 0 \end{pmatrix} = g \begin{pmatrix} \alpha \\ 0 \end{pmatrix} = \alpha g \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \alpha y$$

Thus, every G -invariant map $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a scaling

$$\phi(x) = \alpha x$$

by some $\alpha \in \mathbb{R}$, which is invertible if $\alpha \in \mathbb{R}^\times$.

✎ Exercise

$$\text{Aut}_G \text{Set} \left(\frac{G}{H} \right) \cong \frac{N_G(H)}{H}$$

For our case $H = \mathcal{U}$

$$N_G(\mathcal{U}) = \left\{ \begin{bmatrix} * & * \\ 0 & * \end{bmatrix} \right\} \cong \mathbb{R}^\times \times \mathbb{R}$$

and

$$\frac{N_G(\mathcal{U})}{\mathcal{U}} \cong \mathbb{R}^\times$$

$SL(2, \mathbb{Z})$ -maps

We may ask

?

Question

Find all Borel maps

$$\mathbb{R}^2 \rightarrow \mathbb{R}^2$$

that commutes with $\Gamma := SL(2, \mathbb{Z})$ defined upto Lebesgue measure-zero sets.

whose answer is

Every Borel measurable $SL(2, \mathbb{Z})$ -map $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ is essentially a scaling upto Lebesgue measure-zero sets.

But what is special about Γ ? The fact that it is a "*lattice*" in G that is:

Proposition:

$\Gamma < G$ is discrete and the space

$$G \curvearrowright \frac{G}{\Gamma}$$

has a G -invariant probability measure.

The subgroup

$$A := \left\{ \begin{bmatrix} \lambda & \\ & \lambda^{-1} \end{bmatrix} \mid \lambda \in \mathbb{R}^\times \right\} < G = SL(2, \mathbb{R})$$

which acts by $\begin{bmatrix} -1 & \\ & -1 \end{bmatrix}$ along with the flow

$$\left\{ \exp \left(\alpha \begin{bmatrix} 1 & \\ & -1 \end{bmatrix} \right) \mid \alpha \in \mathbb{R} \right\}$$

✓

flow of [A](#)

✓

This dynamics is "*tame*" as in the orbits are "locally closed".

"Tame dynamics"	"Wild dynamics"
action by $\mathcal{U}, A$ above on $\mathbb{R}^2$	irrational torus rotations
	$\Gamma \curvearrowright \frac{G}{\mathcal{U}}$
	$\mathcal{U} \curvearrowright \frac{G}{\Gamma}$

There are no $SL(2, \mathbb{R})$ -invariant metric on $\mathbb{R}^2$

Every continuous

$$d : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow [0, \infty)$$
$$d(x, y) = d(y, x)$$
$$d(x, x) = 0$$

satisfying triangle inequality which is $SL(2, \mathbb{R})$ -invariant, that is,

$$g \in SL(2, \mathbb{R}) \implies d(gx, gy) = d(x, y)$$


is trivial, as in $d \equiv 0$.

☀️ Fix $\epsilon > 0$.

- The restriction $d(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, -) : \mathbb{R}^2 \rightarrow [0, \infty)$ is continuous so preimage of $(0, \epsilon)$ contains a standard ball say B_ϵ .
- Because $\mathcal{U}$ fixes $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$


$$d(gp, \begin{pmatrix} 1 \\ 0 \end{pmatrix}) = d(p, \begin{pmatrix} 1 \\ 0 \end{pmatrix}) < \epsilon$$

for any $p \in B_\epsilon$.



-
- Thus the $d(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, -)$ -preimage of $(0, \epsilon)$ actually contains a horizontal strip $U_\epsilon := \mathbb{R} \times (-\epsilon, \epsilon)$.
- Now pick $x, y \in \mathbb{R}^2$ and flow them along A so that they both reach the U_ϵ , so

$$gx \in U_\epsilon$$



-
- By triangle-inequality

$$d(x, \begin{pmatrix} 1 \\ 0 \end{pmatrix}) = d(gx, g \begin{pmatrix} 1 \\ 0 \end{pmatrix}) \leq d(gx, \begin{pmatrix} 1 \\ 0 \end{pmatrix}) + d(g \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}) \leq 2\epsilon$$

- By triangle-inequality

$$d(x,y) \leq d(x,\begin{pmatrix} 1 \\ 0 \end{pmatrix}) + d(y,\begin{pmatrix} 1 \\ 0 \end{pmatrix}) = 4\epsilon$$

- So

$$d(x,y) < 4\epsilon$$

for all $\epsilon > 0$. Thus $d \equiv 0$.

Action of $\mathcal{U}$ on G/Γ is ergodic

Corollary of

 Every continuous

$$\begin{aligned} d : \mathbb{R}^2 \times \mathbb{R}^2 &\rightarrow [0, \infty) \\ d(x,y) &= d(y,x) \\ d(x,x) &= 0 \end{aligned}$$

satisfying triangle inequality which is $SL(2,\mathbb{R})$ -invariant, that is,

$$g \in SL(2,\mathbb{R}) \implies d(gx,gy) = d(x,y)$$

is trivial, as in $d \equiv 0$.

 The action of $\mathcal{U}$ on G/Γ is ergodic

Definition. Ergodic actions

An action $G \curvearrowright X$ with an invariant measure class is **ergodic** if either of these is true

- $\forall A \subseteq X$ measurable and invariant A is either a null set or a full measure set
- $L^\infty(X)^G$ is one dimensional
- (if X has a G -invariant probability measure) $L^2(X)^G$ is one dimensional

We observe the action of G on G/Γ is ergodic because the action is transitive?



 Let X is a G -space, $\mathcal{U} < G$ and $\Gamma < G$ is a lattice. Then we have a G -invariant measure on G/Γ so we may consider the L^2 space on it. Then

$$L^2(G/\Gamma)^\mathcal{U} \cong \text{Hom}_G \left(\frac{G}{\mathcal{U}}, L^2(G/\Gamma) \right)$$

•  Every continuous

$$\begin{aligned} d : \mathbb{R}^2 \times \mathbb{R}^2 &\rightarrow [0, \infty) \\ d(x,y) &= d(y,x) \\ d(x,x) &= 0 \end{aligned}$$

satisfying triangle inequality which is $SL(2,\mathbb{R})$ -invariant, that is,

$$g \in SL(2, \mathbb{R}) \implies d(gx, gy) = d(x, y)$$

is trivial, as in $d \equiv 0$.

implying

$$\mathrm{Hom}_G \left(\frac{G}{\mathcal{U}}, L^2(G/\Gamma) \right) \cong L^2(G/\Gamma)^G \cong \mathbb{C}$$

by

We observe the action of G on G/Γ is ergodic because the action is transitive?

Ergodic group actions

Definition. Ergodic actions

An action $G \curvearrowright X$ with an invariant measure class is **ergodic** if either of these is true

- $\forall A \subseteq X$ measurable and invariant A is either a null set or a full measure set
- $L^\infty(X)^G$ is one dimensional
- (if X has a G -invariant probability measure) $L^2(X)^G$ is one dimensional

 Let X is a G -space, $\mathcal{U} < G$ and $\Gamma < G$ is a lattice. Then we have a G -invariant measure on G/Γ so we may consider the L^2 space on it. Then

$$L^2(G/\Gamma)^\mathcal{U} \cong \mathrm{Hom}_G \left(\frac{G}{\mathcal{U}}, L^2(G/\Gamma) \right)$$

duality principle

$$\begin{aligned} \mathrm{Hom}_\Gamma \left(\frac{G}{\mathcal{U}}, \mathbb{R}^2 \right) &\rightarrow \mathrm{Hom}_\mathcal{U} \left(\frac{G}{\Gamma}, \mathbb{R}^2 \right) \\ \Phi &\mapsto \tilde{\Phi}(g) := g\Phi \left(g^{-1} \right)? \end{aligned}$$