

## PRESENT STATUS OF MONKEYPOX IN INDIA

Deeksha Tiwari <sup>\*1</sup>, Mrs. Preeti Mishra <sup>1</sup>, Dilip Kumar Chanchal <sup>2</sup> and Vivek Pal <sup>2</sup>

Faculty of Pharmacy <sup>1</sup>, Raja Balwant Singh Engineering Technical Campus Bichpuri, Agra - 283105, Uttar Pradesh, India.

Department of Pharmacy <sup>2</sup>, Smt. Vidyawati College of Pharmacy, Jhansi - 284128, Uttar Pradesh, India.

### ABSTRACT

The monkeypox poxvirus is linked to smallpox and spreads via human-animal interaction. Previous smallpox protection provides minimal monkeypox protection. Slowly increasing monkeypox cases in Africa were generally ignored by the scientific community until last year, when more than 16,000 cases were recorded from nonendemic nations. WHO recently declared monkeypox a global health emergency. Although most instances are in males who have sex with other men, the disease threatens the whole population. A short feverish sickness with lymphadenopathy preceded the rash, which spreads radially outward. Recovery usually takes two to four weeks. Children, new parents, and those with weakened immune systems are prone to complications. PCR can diagnose viral DNA. Most treatment focuses on symptom relief. Tecovirimat is a good antiviral. High-risk populations, healthcare practitioners, and PCR labs should acquire smallpox immunizations. Symptom relief is usually enough. Tecovirimat fights viruses. Healthcare personnel and individuals in intimate social networks should be vaccinated against smallpox.

**Keywords:** Immunization, Monkeypox, Lesion, Revealed, Pathogenic

**Correspondence to Author:** Deeksha Tiwari, Faculty of Pharmacy 1, Raja Balwant Singh Engineering Technical Campus Bichpuri, Agra - 283105, Uttar Pradesh, India. Email: tiwarideeksha04108@gmail.com

## INTRODUCTION

The monkey pox virus is a zoonotic member of the family Poxviridae, which is composed entirely of Orthopox viruses. It's a DNA virus with two strands, like the one that caused smallpox before it was destroyed. The monkeypox virus it causes is mostly found in the western and central parts of the African continent<sup>1, 2</sup>. The illness has been termed "miniature smallpox" because its relatively tiny size. This virus has several reservoirs in endemic areas of Africa, including squirrels, Gambian pouched rats, and even certain kinds of monkeys, such the cynomolgus monkeys from whom it was initially identified. This zoonotic illness has the potential to spread to people and cause devastating outbreaks. West African and Congo Basin (Central African) clades of monkeypox virus are the only two

clades of monkeypox virus known to have existed at any point in history<sup>3</sup>.

The virus that causes monkeypox was first found in monkeys in a Danish lab in 1958, thus the name. Where smallpox had been declared eradicated in 1968, a 9-year-old kid in the Democratic Republic of the Congo became the first human case to be detected in 1970<sup>2, 4</sup>. Up until 1986, the majority of cases (15 out of 16) had been reported in West African countries, and 85 percent of all cases had originated in Central Africa. Human monkeypox has been documented in West Africa, where it thrives in the hot, humid forest climate and at lower altitudes than in the Congo Basin<sup>5</sup>. Only 13% of those who contracted the illness had Smallpox vaccination scars, hence it mostly struck immunologically unprepared people<sup>3</sup>. Primary exposure to a common animal source or

secondary transmission from human to human led to the reported instances, which occurred at the same time as peak human outdoor activity (mostly hunting and agriculture). Skinning and handling wild animals, close contact with a case, and not being vaccinated against smallpox were identified as risk factors for Monkeypox in a 1997 WHO epidemiological assessment in the Kasai area<sup>6</sup>.

During the last 40 years, there have been regular outbreaks in 11 different African nations. Over 500 cases have been suspected and 200 have been verified in Nigeria since 2017, with a case fatality ratio of 3%<sup>2</sup>. However, it is no longer exclusive to Africa alone. This brief overview of the recent resurgence of Monkeypox epidemics across the globe, particularly in India, highlights the emergency mindset necessary for mounting a coordinated response to prevent, engage, and protect vulnerable populations in society; to strengthen surveillance and infection control; and to promote research into more vaccines and therapeutics for zoonotic diseases. Preventative measures in public health are perhaps more crucial than ever before. India must be prepared to respond vigorously to any public health crisis involving the re-emerging zoonoses (monkeypox, cow-pox, camel-pox) that threaten to fill the ecological void left by the eradication of smallpox<sup>7</sup>. How lethal the illness is, and if its nature and epidemiological qualities may have major repercussions for the health system of India, which has seen a terrible breakdown of the healthcare system owing to COVID-19, are unknowns to the public health and community medicine specialists. This study seeks to rectify the inequitable distribution of the IDSP's new medications, vaccinations, and other interventions among medical practitioners.

## **2. Characteristics Of The Disease**

Suspects often present with vesicular rash, which may or may not be accompanied by additional symptoms and signs include fever, respiratory problems, and lymphadenopathy

(swollen lymph nodes). Rash outbreaks often start between the third and fifth day of a fever. It often takes two to four weeks for them to disappear after spreading from the face to the arms and legs and then the trunk. The oral mucous membranes, conjunctivae, cornea and genitalia are impacted in certain instances. Monkeypox may cause consequences including secondary infections, broncho pneumonia, sepsis, encephalopathy, and loss of eyesight. Lymphadenopathy and a low mortality rate distinguished Monkeypox from Smallpox in clinical presentation (3-6 percent). Lesions, bodily fluids, respiratory droplets, sexual contact, and contaminated items, even in moderate sickness, may all facilitate the transmission of monkeypox<sup>8</sup>.

Suspected cases are being isolated at global transit hubs for 21 days to avoid transmission; the incubation period typically lasts between 6 and 13 days, but may last anywhere from 5 to 21 days. The illness is thought to be self-limiting, although it may be rather severe in those with immune suppression, such as young children, pregnant women, and people with compromised health. Men who have sex with men (MSM) seeking treatment at primary and sexual health clinics have been shown to be at risk. Research conducted in the United Kingdom between April and June 2022 found that 98% of those infected were males and that 41% of them were HIV-positive<sup>9</sup>. Co-infection with hepatitis A, B, or C and monkeypox is prevalent in this population<sup>10</sup>. Health professionals and family members are at a higher risk of infection during human monkeypox epidemics due to their frequent close contact with affected patients. Isolating diseased patients, avoiding contact with potentially infectious animals and fomites, and practising diligent hand cleanliness are also critical preventative measures<sup>11</sup>.

## **3. PRESENT GLOBAL SCENARIO**

Public health officials throughout the world are paying more attention to the outbreak after hearing reports of numerous cases occurring in regions other than Africa. In 2003, the United States of America recorded the first incidence, which was traced back to exposure to sick prairie dogs kept as pets. Patients complained of high body temperatures and a rash that progressed from macule to papule to vesicle to pustule to crusty, dead skin that flaked off. The virus was verified by electron microscopy in both a human skin lesion and a lymph node from a "prairie dog." As a consequence, 79 instances were reported across 6 states, with 29 of those cases verified by laboratory tests<sup>3</sup>. Cases of monkeypox have been documented in tourists visiting Singapore in 2019, the United States of America in 2021, and Israel and the United Kingdom between 2018 and 2022. Cases of this condition have skyrocketed since 1990<sup>13</sup>. The animals that host the virus in Africa and elsewhere are themselves distinct.

Epidemiological research on monkeypox virus epidemics has shown that human expansion into formerly uninhabited forest regions, increased international travel, and the end of smallpox immunisation programmes are major contributing factors<sup>13</sup>. As a result of the massive scale of smallpox vaccination programmes that followed the disease's eradication<sup>14</sup>, the World Health Organization (WHO) named monkeypox a Global Health Emergency of International Concern on July 23, 2022<sup>12</sup>. Cases outside of an endemic area have been reported in 78 of the 194 WHO Member States so far, including 5 in India. As of July 26<sup>th</sup>, 2022, there have been around 18,880 confirmed cases of monkeypox worldwide, with 5 fatalities reported from Africa<sup>15, 16</sup>. Since new zoonotic viruses are on the rise across the globe, we need to pay special attention to the possibility of their spread inside India<sup>2, 17</sup>. The number of reported cases is quite high. The spread of the disease in these outbreak areas is being contained through a combination of contact tracing, clinical management, isolation,

laboratory investigation, genomic sequencing of DNA (where available), and infection prevention and control measures, as well as continuous detection of the virus by established surveillance mechanisms. The nations in question are moving swiftly to get vaccines, and IEC materials and infographics are being deployed to promote awareness. For the first time ever, a two-dose vaccination (MVA-BN) using the Ankara strain of attenuated vaccinia virus has been authorised for the prevention of monkeypox. Monkeypox virus genes found were mostly of the West African lineage.

Currently, two vaccinations (ACAM-2000 and MVA-BN) are in use as preventative measures for close contacts. However, only a few number are now available for purchase<sup>2, 3, 17</sup>. Only Bavarian Nordic has a supply of Monkeypox vaccinations, which industrialised countries like the United States and the United Kingdom have been purchasing. -NHS<sup>18</sup>. In order to expedite the development of a vaccine against the monkeypox virus, the Indian government has extended an invitation for vaccine and in-vitro diagnostic kit producers to submit expressions of interest (EOI) by August 10<sup>th</sup>, 2022. (MPXV)<sup>19</sup>.

#### **4. POSITION OF INDIA**

The World Health Organization (WHO) has established worldwide standards and case criteria for MPXV detection. It has suggested a concerted effort to halt the spread of the disease, include and protect the most vulnerable members of society, step up monitoring, improve infection control and management practises, and speed up the development of vaccines and medicines<sup>20</sup>. Suspected cases include people of any age who report in a non-endemic nation with symptoms including headache, sudden fever onset, lymphadenopathy, myalgia, asthenia, and back pain and an unexplained acute rash that has no recognised clinical aetiology.

A probable case of monkeypox is one in which there is evidence of an epidemiological

connection or direct contact, recent travel to an endemic nation, several sexual partners in the 21 days before to symptom start, hospitalisation, or a positive serological test. Confirmed instances of monkeypox have been found, thanks to the use of real-time polymerase chain reaction (PCR) to sequence the specific viral DNA that had first led to their detection<sup>4</sup>. The National Institute of Virology in Pune, India, has revised the case reporting form for fever with rashes, which may be found on the Integrated Disease Surveillance Programme website. In terms of the regional IDSP, one sample will be approved for MPXV testing at VRDL, another will be sent to NIV Pune, and a third will be preserved for future testing<sup>21</sup>.

When possible instances have been reported from India, the NCDC, MoHFW has stressed certain public health measures.

They are:

- i. People who are thought to have monkeypox or who are considered likely cases must be treated with extreme caution in medical institutions.
- ii. Isolation should be maintained for all suspected cases until all lesions have resolved OR until the treating physician chooses to lift isolation.
- iii. The IDSP District Surveillance Officer should be notified of any incidents of this kind.
- iv. The best possible measures should be taken to prevent the spread of infection at all times throughout patient care.
- v. Vesicle fluid, blood, and sputum samples are to be submitted to NIV Pune for testing; if a positive case is identified, the patient's contacts during the preceding 21 days must be identified and placed under quarantine<sup>12</sup>.

There are a lot of reasons which may make Monkeypox as a severe public health concern in India, beginning with the subtropical environment, and overpopulation with high

transmission rates. The first laboratory-confirmed case was recorded on July 14<sup>th</sup> 2022 in a 35 year-old guy from Thiruvananthapuram, Kerala, upon returning from UAE [22]. The next two cases were also confirmed from Kerala on July 18<sup>th</sup> and 22<sup>nd</sup><sup>23</sup>. New Delhi reported its first MPXV case in a 34 year-old man on July 24<sup>th</sup> with no history of travel, unlike the previous cases<sup>24</sup>. The area continues to be on high alert as a 47-year-old woman has approached the health department on July 26<sup>th</sup> and is currently the 5<sup>th</sup> in the growing list as a suspected case of MPXV<sup>25</sup>.

There is now a continuing monitoring in the community, primary care, fever clinics, sexual health clinics, infectious illness units, obstetrics and gynaecology, and dermatology clinics. Contact tracing should be undertaken as soon as there is probable case identification. The patients are to be interrogated to collect the names and contact information of any contacts, who should be informed within 24 hours after identification. The contacts would then be held in isolation for 21 days<sup>26</sup>. However, we cannot avoid talking about the obvious: the rapidity with which this sickness has been crossing across borders. Forty percent of India's population is comprised of people aged 13 to 35 who are not protected by the disease's vaccine<sup>27</sup>. There has to be a supply of the smallpox vaccine in case an epidemic occurs in the nation. This is not a very unique or changeable illness; thus, the likelihood of it becoming a major public health problem in the United States decreases if we are appropriately prepared. When caring for a patient diagnosed with Monkeypox, medical staff should always wear appropriate PPE. A medical mask should be worn whenever the patient is within a 1-meter radius of a healthcare provider or another patient. In 2022, Tecovirimat, an antiviral drug, will be available for the treatment of monkeypox, as reported by the World Health Organization. While this medication and the new vaccination MVA-BN are not yet widely accessible, India may be ready for an outbreak by keeping an eye on airports and other transportation hubs,

as well as educating its population about the epidemiology and warning symptoms of the illness<sup>4</sup>.

The International Committee for Medical Research (ICMR) has issued recommendations for the management of symptomatic treatment, including measures to prevent further damage to already damaged skin and mucous membranes, as well as rehydration and nutritional therapy<sup>26</sup>. The dynamic epidemiology of this resurging illness may only be understood via heightened monitoring and case detection efforts. One Health is more important today than it has ever been. The Institut Pasteur de Bangui in the Central African Republic is conducting a clinical trial known as "A One Health Study of Monkeypox Human Infection" (AFRIPOX)<sup>28</sup>. In order to combat smallpox in India, we need to better understand its zoonotic and human-to-human risk factors, animal reservoirs, post-vaccination immunity status of vaccinated individuals, and the differences between circulating strains in human and animal populations, as well as develop diagnostics and next-generation sequencing<sup>29</sup>.

The healthcare staff might utilise a technique called 'ring vaccination,' in which they vaccinate the close contacts of persons who have been infected with monkeypox, to prevent the spread of the disease. MoHFW collaborates with the Ministries of Agriculture, Fisheries, Animal Husbandry, and Dairy to provide a comprehensive response to any emergency. With regards to monkeypox research, WHO has suggested consulting with the Emergencies Social Science Technical Working Group, the Scientific Advisory Group for the Origins of Novel Pathogens (SAGO), the Strategic and Technical Advisory Group on Infectious Hazards (STAG-IH), the SAGE working group on smallpox and monkeypox vaccines, and the WHO Research & Development Blueprint<sup>30</sup>.

## CONCLUSION:

The poxvirus that causes monkeypox is closely linked to the virus that causes smallpox, and it spreads from monkeys to humans via intimate contact between the two species. The protection that was previously received against smallpox gives only a moderate degree of protection against monkeypox. The gradual increase in the number of cases of monkeypox that has been observed in Africa over the past few decades has been largely disregarded by the scientific community around the world up until this year, when more than 16,000 cases were documented from states that are not endemic for the disease. The World Health Organization (WHO) has only recently recognised monkeypox as a public health issue on a global scale. Although the majority of recent cases have been found in males who have had sexual contact with other males, the condition poses a greater threat to the entire population as a whole. The rash is preceded by a brief episode of feverish illness that is accompanied by lymphadenopathy. The rash develops from a lesion to a papule to a vesicle to a pustule, and then it spreads radially outward. The majority of people go back to normal in two to four weeks, depending on the severity of their condition. Complications are most prevalent in youngsters, mothers who have just given birth, and those whose immune systems are compromised.

For diagnostic purposes, the polymerase chain reaction (PCR) is utilised to precisely identify viral DNA. In the majority of instances, all that is required is the treatment of symptoms. Tecorivimat is effective in warding against viral infections. Those at a higher risk of contracting smallpox, including high-risk groups, healthcare practitioners, and labs that do polymerase chain reaction (PCR), are all strongly recommended to be vaccinated against the disease. In the majority of instances, all that is required is the treatment of symptoms. Tecorivimat is effective in warding against viral infections. Those who fall into one of the high-risk groups, such as those who work in the medical field or have close personal relationships, are strongly



encouraged to be vaccinated against smallpox using one of the vaccines now on the market.

## REFERENCES

- [1]. Moore M, Zahra F. Monkeypox. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022. [cited 2022 May 24]. Available: <http://www.ncbi.nlm.nih.gov/books/NBK574519/>
- [2]. WHO. Monkeypox [Internet]; 2022. Available: <https://www.who.int/newsroom/factsheets/detail/monkeypox>
- [3]. Georges AJ, Matton T, Courbot-Georges MC. Le monkey-pox, un paradigme de maladie émergente puis réémergente. Médecine Mal Infect. 2004; 34(1):12–9.
- [4]. Multi-country monkeypox outbreak in nonendemic countries [Internet]; 2022. Available: <https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON385>
- [5]. Levine RS, Peterson AT, Yorita KL, Carroll D, Damon IK, Reynolds MG. Ecological Niche and Geographic Distribution of Human Monkeypox in Africa. PLoS One. 2007;2(1):e176.
- [6]. Heymann DL, Szczeniowski M, Esteves K. Re-emergence of monkeypox in Africa: A review of the past six years. Br Med Bull. 1998;54(3):693–702.
- [7]. Cohen J. Is an old virus up to new tricks? Science. 1997;277(5324):312–3.
- [8]. BBC. As it happened: UK monkeypox cases rise to 57 [Internet]. BBC News; 2022.
- [9]. Thornhill JP, Barkati S, Walmsley S, Rockstroh J, Antinori A, Harrison LB, et al. Monkeypox Virus Infection in Humans across 16 Countries — April–June 2022. N Engl J Med. 2022;0(0):null.
- [10]. Liu X, Zhu Z, He Y, Lim JW, Lane B, Wang H, et al. Monkeypox claims new victims: the outbreak in men who have sex with men. Infect Dis Poverty. 2022;11(1):84.
- [11]. California Department of Public Health. Monkeypox [Internet]. [cited 2022 Jun 15]; 2022. Available: <https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Monkeypox.aspx>
- [12]. Resources: Integrated Disease Surveillance Programme (IDSP) [Internet]. May 20, 2022. Available: <https://idsp.nic.in/index4.php?lang=1&level=0&linkid=414&lid=1610&Background=Light>
- [13]. Bunge EM, Hoet B, Chen L, Lienert F, Weidenthaler H, Baer LR, et al. The changing epidemiology of human monkeypox—A potential threat? A systematic review. PLoS Negl Trop Dis. 2022;16(2):e0010141.
- [14]. WHO Director-General's statement at the press conference following IHR Emergency Committee regarding the multicountry outbreak of monkeypox - 23 July 2022 [Internet]. [cited 2022 Jul 27]. Available: <https://www.who.int/director-general/speeches/detail/who-director-general-s-statement-on-the-press-conference-following-IHR-emergency-committee-regarding-the-multi-country-outbreak-of-monkeypox--23-july-2022>
- [15]. Mathieu E, Dattani S, Ritchie H, Roser M. Monkeypox. Our World Data [Internet]. 2022. [cited 2022 Jun 15]. Available: <https://ourworldindata.org/monkeypox>
- [16]. WHO reports 14,000 cases of monkeypox globally, five deaths in Africa | Reuters [Internet]. [cited 2022 Jul 27]. Available: <https://www.reuters.com/business/healthcare-pharmaceuticals/who-reports-14000-cases-monkeypox-globally-five-deaths-africa-2022-07-20/>
- [17]. Multi-country monkeypox outbreak: situation update [Internet]; 2022 [cited 2022 Jun 20]. Available: <https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON392>
- [18]. Jul 20 SS| NR| CN|, 2022. WHO: 14,000 monkeypox cases worldwide, 5 deaths [Internet]. CIDRAP. [cited 2022 Jul 27]. Available: <https://www.cidrap.umn.edu/news-perspective/2022/07/who-14000-monkeypox-cases-worldwide-5-deaths>
- [19]. July 27 SMMSND, July 27 2022 UPDATED: Ist 2022 20:10. Centre floats tender to develop monkeypox vaccine [Internet]. India Today. [cited 2022 Jul 27]. Available: <https://www.indiatoday.in/india/story/govt-floats-tender-for-developing-vaccine-against-monkeypox-1980704-2022-07-27>
- [20]. Monkeypox declared a global health emergency by the World Health Organization [Internet]. UN News; 2022 [cited 2022 Jul 27]. Available: <https://news.un.org/en/story/2022/07/1123152>
- [21]. Indian Council of Medical Research, New Delhi. Instructions on MPXV\_26.07.2022.pdf; 2022.
- [22]. Bureau TH. Kerala reports India's first confirmed monkeypox case. The Hindu [Internet]. 2022 [cited 2022 Jul 27]. Available: <https://www.thehindu.com/scitech/health/kerala-reports-indias-first-confirmed-monkeypox-case/article65640571.ece>
- [23]. Explained: Third case of monkeypox confirmed in Kerala, is there a risk of an outbreak in India? [Internet]. The Indian Express; 2022 [cited 2022 Jul 27]. Available: <https://indianexpress.com/article/explained/second-case-monkeypox-kerala-outbreak-india-8037348/>
- [24]. India's 4th monkeypox case in Delhi, patient has no foreign travel history. The Times of India [Internet]; 2022 [cited 2022 Jul 27]; Available: <https://timesofindia.indiatimes.com/india/indias-4th-monkeypox-case-in-delhi-patient-has-no-foreign-travel-history/articleshow/93096145.cms>
- [25]. Dash N. Monkeypox in Noida: Suspected case reported, 5 th case in India [Internet]; 2022 [cited 2022 Jul 27]. Available: <https://www.indiatvnews.com/news/india/monkeypox-case-in-noida-uttar-pradesh-viral-zoonosis-2022-07-27-795321>
- [26]. Ministry of Health and Family Welfare, GOI. Guidelines for Management of Monkeypox Disease; 1970. Available: <https://main.mohfw.gov.in/diseases/alerts-0>
- [27]. Decent Work for Youth in India [Internet]; 2012 [cited 2022 Jun 20].

- Available:[http://www.ilo.org/newdelhi/info/WCMS\\_175936/lang--en/index.htm](http://www.ilo.org/newdelhi/info/WCMS_175936/lang--en/index.htm)
- [28]. Institut Pasteur. A One Health Study of Monkeypox Human Infection, Animal Reservoir, Disease Ecology and Diagnostic Tools [Internet]. [clinicaltrials.gov](https://clinicaltrials.gov); 2022 [cited 2022 May 23]. Report No.: NCT05058898. Available:<https://clinicaltrials.gov/ct2/show/NCT05058898>
- [29]. A One health study of Monkeypox: Human infection, animal reservoir, disease ecology and diagnostic tools [Internet]. Agencenationale de la recherche. [cited 2022 May 24]; 2022. Available:<https://anr.fr/Project-ANR-19-CE35-0011>
- [30]. Multi-country monkeypox outbreak: Situation update [Internet]; 2022 [cited 2022 Jun 22].

**How to Cite**

Tiwari D, Mishra P, Chanchal K D, Pal V. Present Status of Monkey pox In India. *International Journal of Pharmaceutical Science and Medicine* 2022; 1(1): 15-21.