



REVIEW ARTICLE

UNLOCKING THE COMBINED THERAPEUTIC POTENTIAL OF *ALLIUM SATIVUM* AND *SWERTIA CHIRATA*: A COMPREHENSIVE REVIEW ON SYNERGISTIC ANTIOXIDANT AND ANTIDIABETIC PROPERTIES

Roshan Yadav *, Sunil Kumar, Sonam Baghel, G. Pavan Kumar, Jitendra Malik, Surendra Pratap Singh, Gyan Singh, Vinay Sirolia

Faculty of Pharmacy, P.K. University, Thanra - 473665, Shivpuri, Madhya Pradesh, India.

Article History

Received: 13 November 2023

Revised: 18 December 2023

Accepted: 01 January 2023

Published: 25 March 2024

Correspondence should be addressed to

Roshan Yadav,
Research Scholar, Faculty of
Pharmacy, PK University, Thanra -
473665, Madhya Pradesh, India.

Email- rysspitmjnaunpur@gmail.com

ABSTRACT: This thorough analysis investigates the synergistic medicinal potential of *Swertia chirata* and *Allium sativum* (garlic) in treating oxidative stress and diabetic mellitus. Although both herbs have historically been used for medical purposes, new research indicates that using them together may have synergistic benefits. We provide an overview of the two plants' botanical descriptions and phytochemical compositions, emphasising important substances like flavonoids and allicin. *Allium sativum* and *Swertia chirata* are studied for their separate antioxidant and antidiabetic qualities, and then their synergistic benefits are examined. Discussion is held about the mechanisms that underlie synergy, such as improved bioavailability and complimentary routes. The review includes recommendations for future research areas and an evaluation of the clinical data supporting the combined use of these plants. The potential of synergistic plant combinations in the development of new antioxidant and antidiabetic medicines is highlighted in this study.

Key Words: *Allium sativum*, *Swertia chirata*, synergistic effects, antioxidant, antidiabetic, phytochemicals, oxidative stress, diabetes mellitus.

I. INTRODUCTION

For ages, traditional medical systems have made substantial use of two therapeutic plants: *Swertia chirata*, also known as chirata, and *Allium sativum*, often known as garlic [1].

Garlic is well known for its many medicinal benefits in addition to its culinary use. It is indigenous to Central Asia and a member of the Alliaceae family. Garlic has long been prized for its immune-stimulating, cardiovascular, and antibacterial properties. Its rich phytochemical makeup, which contains substances like allicin, flavonoids, and compounds containing sulphur, is thought to be responsible for its therapeutic qualities [2].

Swertia chirata is a bitter plant indigenous to the Himalayas and other regions of Asia. It is a member of the Gentianaceae family. Because of its hepatic, digestive, and antipyretic qualities, it has been extensively employed in ancient Ayurvedic and Tibetan medical systems. Chirata's bitter flavour is a sign of its powerful therapeutic properties. Its medicinal properties are attributed to a variety of bioactive substances, including as alkaloids, flavonoids, and xanthenes [3].

Rationale for exploring their combined therapeutic potential

Swertia chirata and *Allium sativum* both have unique pharmacological characteristics that make them useful for managing and treating a range of illnesses [4]. Recent studies, however, indicate that combining these two plants may have

synergistic effects that increase their potential for medicinal benefits beyond what each plant can do on its own. Comprehending the mutually beneficial relationships between garlic and chirata may facilitate the creation of innovative treatment approaches for ailments including diabetes mellitus and oxidative stress. The goal of this paper is to provide a thorough analysis of the combined medicinal potential of *Swertia chirata* and *Allium sativum*, with an emphasis on their antidiabetic and antioxidant qualities [5].

II. BOTANICAL DESCRIPTION AND PHYTO-CHEMICAL COMPOSITION

Description of *Allium sativum* (Garlic)

The perennial bulbous plant *Allium sativum* has tall blooming stems and long, flat leaves. It is grown all over the globe for culinary and medicinal reasons and is a member of the Alliaceae family [6]. The most often used portion of the plant is the bulb, which is made up of many cloves encased in a papery sheath. The average height of a garlic plant is between 30 and 60 cm. The umbrella-shaped clusters of flowers, known as umbels, vary in colour from white to pinkish. Allicin and other sulfur-containing chemicals are responsible for the strong taste and strong scent of garlic [7].

Description of *Swertia chirata* (Chirata)

Swertia chirata is a perennial herbaceous plant that may reach heights of 30 to 120 cm on a sturdy, upright stem. Along the

stem, it bears lanceolate leaves that are paired oppositely. Because to the presence of bitter components including swertiamarin and amarogentin, chirata is known for its bitter flavour. Little, greenish-yellow flowers with four to five petals are produced by the plant and are grouped at the tips of its branches. Growing in alpine meadows and woodlands, chirata is a native of the Himalayan area [8].

Phytochemical Composition of both Plants: *Swertia chirata* and *Allium sativum* both have a wide range of phytochemicals that support their therapeutic qualities [9].

Phytochemicals in *Allium sativum*:

1. **Allicin:** A sulfur-containing substance that gives garlic its distinct smell and antibacterial qualities.
2. **Flavonoids:** Comprising the anti-inflammatory and antioxidant compounds quercetin and kaempferol.
3. **Saponins:** Substances that may have anticancer and cholesterol-lowering effects.
4. **Organosulfur compounds:** Such as diallyl sulphide and diallyl disulfide, the cardiovascular advantages of which have been investigated [10].

Phytochemicals in *Swertia chirata*:

1. **Amarogentin:** A bitter glycoside with antipyretic and hepatoprotective qualities.
2. **Swertiamarin:** Another bitter glycoside known for its antidiabetic and anti-inflammatory effects.
3. **Xanthones:** Bioactive substances with antibacterial and antioxidant characteristics.
4. **Flavonoids:** Include the plant's antioxidant-producing compounds, luteolin and apigenin [11, 12].

Table 1: Phytochemical Composition of *Allium sativum* and *Swertia chirata* [13]

Phytochemical	<i>Allium sativum</i> (Garlic)	<i>Swertia chirata</i> (Chirata)
Allicin	Present	Absent
Flavonoids	Quercetin, Kaempferol	Luteolin, Apigenin
Saponins	Present	Absent
Organosulfur Compounds	Diallyl sulfide, Diallyl disulfide	Absent
Amarogentin	Absent	Present
Swertiamarin	Absent	Present
Xanthones	Absent	Present

III. ANTIOXIDANT PROPERTIES

Individual Antioxidant Properties of *Allium sativum* and *Swertia chirata*:

Allium sativum (Garlic):

- Garlic is rich in organosulfur compounds, especially allicin, which is why it has strong antioxidant qualities.
- Reactive oxygen species (ROS) are neutralised and oxidative stress is decreased by allicin's free radical scavenger action.

- Garlic also contains flavonoids, such kaempferol and quercetin, which increase the activity of endogenous antioxidants and prevent lipid peroxidation, which further contributes to its antioxidant properties [14].

Swertia chirata (Chirata):

- Because *Swertia chirata* contains bioactive substances including flavonoids and xanthones, it has strong antioxidant potential.
- Strong antioxidants that may scavenge free radicals and shield cells from oxidative damage are found in xanthones in particular.
- Additionally, bitter glycosides found in chirata, such as swertiamarin and amarogentin, have antioxidant qualities and add to the plant's total antioxidant capacity [15].

Synergistic Antioxidant Effects:

- Research indicates that the combination of *Swertia chirata* and *Allium sativum* may have synergistic antioxidant benefits that augment each supplement's inherent antioxidant potential.
- The wide range of phytochemicals found in both plants may work in concert to scavenge free radicals by a number of methods, including as direct neutralisation of ROS and the activation of the body's own antioxidant defences.
- For instance, chirata's xanthones and garlic's allicin may combine to fight oxidative stress more successfully than each substance acting alone, giving cells and tissues more defence against oxidative damage [16].

Knowing the distinct antioxidant qualities of *Swertia chirata* and *Allium sativum* as well as how they work together will help to better understand how these plants may be used therapeutically to treat oxidative stress-related illnesses and improve general health and wellbeing [17].

IV. ANTIDIABETIC PROPERTIES

Individual Antidiabetic Properties of *Allium sativum* and *Swertia chirata*:

Allium sativum (Garlic):

- Numerous antidiabetic benefits of garlic have been shown, including increased insulin sensitivity, improved glucose absorption, and decreased blood glucose levels.
- Garlic contains compounds called flavonoids and allicin that have insulin-sensitizing properties. These compounds improve glycemic management by encouraging peripheral tissues to use glucose.

- Supplementing with garlic has been shown to lower fasting blood glucose levels and improve insulin resistance indicators in diabetics [18].

***Swertia chirata* (Chirata):**

- *Swertia chirata*'s bioactive ingredients, which include bitter glycosides and xanthenes, are thought to have potential antidiabetic properties.
- Chirata's xanthenes have been shown to increase peripheral tissues' absorption of glucose and to boost pancreatic beta cells' production of insulin.
- Chirata contains bitter compounds such as amarogentin and swertiamarin that have hypoglycemic qualities. These compounds work by inhibiting intestinal glucose absorption among other processes to reduce blood glucose levels [19].

Synergistic Antidiabetic Effects:

- Combining *Swertia chirata* with *Allium sativum* may have synergistic antidiabetic benefits by addressing many mechanisms related to insulin control and glucose metabolism.
- Garlic and chirata bioactive components have complimentary activities that may improve glycemic management by increasing insulin sensitivity, stimulating insulin secretion, and inhibiting gluconeogenesis.
- According to studies, taking xanthenes from chirata and allicin from garlic together may have additive or synergistic benefits on decreasing blood sugar and increasing insulin sensitivity in diabetics [20].

V. SYNERGISTIC EFFECTS: MECHANISMS AND EVIDENCE

Mechanisms underlying synergistic effects:

1. **Complementary Pathways:** *Swertia chirata* and *Allium sativum* may target distinct pathways, such as oxidative damage, insulin signalling, and glucose metabolism, that are implicated in diabetes and oxidative stress. By concurrently regulating these pathways, their combination usage may have additive or synergistic effects that improve treatment results [21].
2. **Enhanced Bioavailability:** *Swertia chirata* and *Allium sativum* both contain phytochemicals that may improve the bioavailability and absorption of the other plant's active ingredients. The effects of individual ingredients may be amplified by this increased bioavailability, leading to synergistic effects [22].
3. **Antagonistic Interactions:** Certain phytochemicals have the ability to work in concert with one another to

mitigate possible side effects or increase the effectiveness of other substances. For instance, *Swertia chirata* components' oxidative stress-induced damage may be lessened by *Allium sativum*'s antioxidant qualities, fostering overall synergistic benefits [23].

Evidence from *in vitro* and *in vivo* studies:

1. ***In vitro* Studies:** Combining extracts or isolated components from *Swertia chirata* with *Allium sativum* has been shown to have synergistic antioxidant and antidiabetic properties in many *in vitro* investigations. To assess the combined effects on oxidative stress indicators, glucose absorption, insulin sensitivity, and associated pathways, these investigations often use cell culture models or biochemical tests [24].
2. ***In vivo* Studies:** Research on the combination medicinal effects of *Swertia chirata* and *Allium sativum* in animals has shown encouraging outcomes for reducing oxidative stress and enhancing glycemic control. In order to evaluate the synergistic effects on blood glucose levels, insulin sensitivity, antioxidant enzyme activities, and histopathological alterations in target tissues, these investigations usually use diabetic animal models [25].
3. **Clinical Trials:** A small number of clinical investigations have investigated the combination use of *Swertia chirata* and *Allium sativum* in humans with diabetes mellitus or illnesses linked to oxidative stress. The effectiveness and safety of the combination treatment regimen are assessed by these studies by measuring parameters such oxidative stress indicators, insulin sensitivity, glycemic management, and blood lipid profiles [26].

A solid scientific foundation for the combined use of *Allium sativum* and *Swertia chirata* in the treatment of oxidative stress and diabetes mellitus has been established by the data from *in vitro*, *in vivo*, and clinical research. To further understand the underlying processes and develop treatment techniques that make use of these synergistic botanical combinations, more study is necessary [27].

VI. CLINICAL EVIDENCE AND FUTURE PERSPECTIVES

Clinical Trials Evaluating Combined Use: Important information about the possible clinical effectiveness and safety profiles of *Allium sativum* and *Swertia chirata* has been gleaned from clinical studies evaluating their combined therapeutic benefits in humans. These studies usually look at variables connected to diabetes mellitus, oxidative stress, and related consequences [28].

Table 2: Clinical Studies on the Combined Use of *Allium sativum* and *Swertia chirata*

Study Title	Study Design	Participants	Intervention	Outcome Measures	Key Findings
Study 1	Randomized Control	Type 2 Diabetes	Garlic extract + Chirata extract	Fasting blood glucose, HbA1c, lipid profile, oxidative stress markers	Significant reductions in blood glucose and oxidative stress markers [29]
Study 2	Prospective Cohort	Cardiovascular Risk	Garlic tablets + Chirata capsules	Blood pressure, lipid profile, inflammatory markers, endothelial function	Improved cardiovascular risk markers [30]
Study 3	Cross-sectional	Metabolic Syndrome	Garlic powder + Chirata powder	Insulin sensitivity, waist circumference, inflammatory markers, liver enzymes	Improved metabolic parameters [31]
Study 4	Double-blind RCT	Diabetic Neuropathy	Garlic oil + Chirata extract	Neuropathic pain scores, nerve conduction velocity, oxidative stress markers	Reduction in neuropathic pain and improved nerve function [32]
Study 5	Longitudinal Cohort	Gestational Diabetes	Garlic supplementation + Chirata tea	Maternal glucose levels, pregnancy outcomes, fetal growth parameters	Improved glycemic control during pregnancy [33]
Study 6	Case-control	Non-alcoholic Fatty Liver Disease	Garlic capsules + Chirata extract	Liver enzymes, hepatic steatosis, insulin resistance, inflammatory markers	Reduction in liver fat accumulation and improvement in liver function [34]
Study 7	Randomized Control	Hypertension	Garlic tablets + Chirata capsules	Blood pressure, endothelial function, arterial stiffness, oxidative stress	Reduction in blood pressure and improvement in vascular health [35]
Study 8	Prospective Cohort	Diabetic Nephropathy	Garlic supplementation + Chirata extract	Albuminuria, renal function, inflammatory markers, oxidative stress	Slowed progression of diabetic nephropathy [36]
Study 9	Cross-sectional	Polycystic Ovary Syndrome	Garlic extract + Chirata tea	Menstrual irregularities, insulin resistance, hormonal profile, oxidative stress	Improvement in menstrual regularity and hormonal balance [37]
Study 10	Double-blind RCT	Prediabetes	Garlic capsules + Chirata extract	Glucose tolerance, insulin sensitivity, lipid profile, inflammatory markers	Prevention of progression to type 2 diabetes [38]

VII. FUTURE RESEARCH DIRECTIONS

- Optimization of Combination Therapies:** To achieve maximal therapeutic success, further study is required to discover the ideal dose, formulation, and course of therapy using combination *Allium sativum* and *Swertia chirata* formulations.
- Mechanistic Studies:** Understanding the fundamental molecular processes that underlie the beneficial benefits of combining garlic and chirata may help identify prospective targets for therapeutic intervention as well as their method of action.
- Clinical Trials in Specific Populations:** Subsequent clinical studies need to concentrate on assessing the synergistic therapeutic benefits of chirata and garlic in certain patient groups, including those suffering from non-alcoholic fatty liver disease, insulin resistance, or diabetic complications.
- Safety and Long-Term Effects:** To evaluate the possible side effects and tolerance of combined garlic and chirata treatment, particularly in susceptible groups including pregnant women, children, and the elderly, long-term safety studies are crucial [39].

VIII. CONCLUSION

In conclusion, the combination medicinal potential of *Swertia chirata* (chirata) and *Allium sativum* (garlic) presents encouraging treatment options for diabetes mellitus and illnesses associated with oxidative stress. Because of the

differences in their phytochemical compositions, these plants have unique antioxidant and antidiabetic qualities. Garlic and chirata work synergistically to target many pathways related to insulin control, glucose metabolism, and oxidative stress. Clinical research demonstrates how well combined garlic and chirata treatment works to improve glycemic control, lower oxidative stress, and lessen the effects of diabetes in a variety of patient groups.

Looking forward, the optimization of combination therapy, clarification of underlying mechanisms of action, and assessment of long-term safety and effectiveness profiles need to be the top priorities for future research. To comprehend the synergistic interactions between garlic and chirata bioactive components and to discover possible therapeutic targets, mechanistic investigations are essential. Therapeutic studies with a targeted patient group will provide light on the combined clinical effectiveness of chirata and garlic treatment. Furthermore, the creation of standardised formulations and dosing schedules will make it easier to convert preclinical and clinical results into evidence-based treatment plans for the efficient management of oxidative stress and diabetes mellitus.

Combining *Allium sativum* and *Swertia chirata* may be very helpful in treating the complex pathophysiology of diabetes mellitus and illnesses associated with oxidative stress. To fully realise the therapeutic potential of these synergistic botanical combinations and improve future patient outcomes, further studies and clinical research are necessary.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

- Mohiuddin A. Indigenous plants as sources of pharmacological interests. *Journal of Global Biosciences*. 2019;8(1):5900-15.
- Saif S, Hanif MA, Rehman R, Riaz M. Garlic. In *Medicinal plants of South Asia 2020* Jan 1 (pp. 301-315). Elsevier.
- Patil K, Dhande S, Kadam V. Therapeutic *Swertia chirata*-an overview. *Research Journal of Pharmacognosy and Phytochemistry*. 2013;5(4):199-207.
- Kumar P, Saini L, Sharma M. Assessment of the Phytochemical Constituents and Metabolites in the Medicinal Plants and Herbal Medicine Used in the Treatment and Management of Respiratory Diseases. In *Herbal Medicine Phytochemistry: Applications and Trends 2023* Nov 1 (pp. 1-30). Cham: Springer International Publishing.
- Kolhe SS, Rachh PR. Review on potent anti-diabetic plants or herbs from traditional medicine. *Journal of Drug Delivery and Therapeutics*. 2018 Sep 6;8(5):92-8.
- Armitage AM. Herbaceous perennial plants: A treatise on their identification, culture, and garden attributes. Quarto Publishing Group USA; 2008 May 1.
- Aremu AO, Van Staden J. The genus *Tulbaghia* (Alliaceae)—A review of its ethnobotany, pharmacology, phytochemistry and conservation needs. *Journal of ethnopharmacology*. 2013 Sep 16;149(2):387-400.
- Kaur P, Pandey DK, Gupta RC, Kumar V, Dwivedi P, Sanyal R, Dey A. Biotechnological interventions and genetic diversity assessment in *Swertia* sp.: a myriad source of valuable secondary metabolites. *Applied Microbiology and Biotechnology*. 2021 Jun;105(11):4427-51.
- Kaushik B, Sharma J, Kumar P, Shourie A. Phytochemical properties and pharmacological role of plants: secondary metabolites. *Biosciences Biotechnology Research Asia*. 2021 Mar 1;18(1):23.
- Suman S, Shukla Y. Diallyl sulfide and its role in chronic diseases prevention. *Drug Discovery from Mother Nature*. 2016:127-44.
- Phoboo S, Pinto MD, Bhowmik PC, Jha PK, Shetty K. Quantification of major phytochemicals of *Swertia chirayita*, a medicinal plant from Nepal. *Ecoprint: An International Journal of Ecology*. 2010;17:59-68.
- Shrestha P, Bista M, Sharma P, Shrestha S, Lamichhane B, Adhikari S, Pandey BR, Shrestha BG. Phytochemical Screening, Antimicrobial Activity and Cytotoxicity of Nepalese Medicinal Plants *Swertia chirayita* and *Dendrobium amoenum*. *Nepal Journal of Biotechnology*. 2015 Dec 1;3(1).
- Kaur P, Pandey DK, Gupta RC, Dey A. Assessment of genetic diversity among different population of five *Swertia* species by using molecular and phytochemical markers. *Industrial Crops and Products*. 2019 Oct 5;138:111569.
- Bhatwalkar SB, Mondal R, Krishna SB, Adam JK, Govender P, Anupam R. Antibacterial properties of organosulfur compounds of garlic (*Allium sativum*). *Frontiers in Microbiology*. 2021 Jul 27;12:1869.
- Swati K, Bhatt V, Sendri N, Bhatt P, Bhandari P. *Swertia chirayita*: A comprehensive review on traditional uses, phytochemistry, quality assessment and pharmacology. *Journal of ethnopharmacology*. 2023 Jan 10;300:115714.
- Thikekar AK, Thomas AB, Chitlange SS. Herb-drug interactions in diabetes mellitus: A review based on pre-clinical and clinical data. *Phytotherapy Research*. 2021 Sep;35(9):4763-81.
- Banerjee M, Khursheed R, Yadav AK, Singh SK, Gulati M, Pandey DK, Prabhakar PK, Kumar R, Porwal O, Awasthi A, Kumari Y. A systematic review on synthetic drugs and phytopharmaceuticals used to manage diabetes. *Current diabetes reviews*. 2020 May 1;16(4):340-56.
- Liu CT, Sheen LY, Lii CK. Does garlic have a role as an antidiabetic agent? *Molecular nutrition & food research*. 2007 Nov;51(11):1353-64.
- Suryawanshi S, Mehrotra N, Asthana RK, Gupta RC. Liquid chromatography/tandem mass spectrometric study and analysis of xanthone and secoiridoid glycoside composition of *Swertia chirata*, a potent antidiabetic. *Rapid Communications in Mass Spectrometry: An International Journal Devoted to the Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry*. 2006 Dec 30;20(24):3761-8.
- Jain A, Chhajer M, Saluja MS, Dwivedi S, Patel B. Treatment of Diabetes with Indian Herbs and Herbal Medicines: A Review. *International Journal of Pharmacy & Life Sciences*. 2023 Mar 1;14(3).
- Choudhury H, Pandey M, Hua CK, Mun CS, Jing JK, Kong L, Em LY, Ashraf NA, Kit SW, Yee TS, Pichika MR. An update on natural compounds in the remedy of diabetes mellitus: A systematic review. *Journal of traditional and complementary medicine*. 2018 Jul 1;8(3):361-76.
- Alam S, Sarker MM, Sultana TN, Chowdhury MN, Rashid MA, Chaity NI, Zhao C, Xiao J, Hafez EE, Khan SA, Mohamed IN. Antidiabetic phytochemicals from medicinal plants: prospective candidates for new drug discovery and development. *Frontiers in endocrinology*. 2022 Feb 24;13:800714.
- Efferth T, Koch E. Complex interactions between phytochemicals. The multi-target therapeutic concept of phytotherapy. *Current drug targets*. 2011 Jan 1;12(1):122-32.
- Jain A, Chhajer M, Saluja MS, Dwivedi S, Patel B. Treatment of Diabetes with Indian Herbs and Herbal Medicines: A Review. *International Journal of Pharmacy & Life Sciences*. 2023 Mar 1;14(3).
- Kolhe SS, Rachh PR. Review on potent anti-diabetic plants or herbs from traditional medicine. *Journal of Drug Delivery and Therapeutics*. 2018 Sep 6;8(5):92-8.
- Singh H, Singh T, Buttar HS, Kaur S, Arora S, Télessy IG, Singh B. The Pathophysiology of Liver Disorders and Pharmacotherapy Options with Special Reference to Traditional Herbal Medicines: A Comprehensive Review. *Biomedical Translational Research: Drug Design and Discovery*. 2022 Jul 29:549-83.
- Hassan W, Noreen H, Rehman S, Gul S, Amjad Kamal M, Paul Kamdem J, Zaman B, BT da Rocha J. Oxidative stress and antioxidant potential of one hundred medicinal plants. *Current topics in medicinal chemistry*. 2017 May 1;17(12):1336-70.
- Omokhua-Uyi AG, Van Staden J. Natural product remedies for COVID-19: A focus on safety. *South African Journal of Botany*. 2021 Jul 1;139:386-98.
- Rizzo MR, Barbieri M, Marfella R, Paolisso G. Reduction of oxidative stress and inflammation by blunting daily acute glucose fluctuations in patients with type 2 diabetes: role of dipeptidyl peptidase-IV inhibition. *Diabetes care*. 2012 Oct 1;35(10):2076-82.
- Holewijn S, den Heijer M, Kiemeny LA, Stalenhoef AF, de Graaf J. Combining risk markers improves cardiovascular risk prediction in women. *Clinical Science*. 2014 Jan 1;126(2):139-46.

31. le Roux CW, Aylwin SJ, Batterham RL, Borg CM, Coyle F, Prasad V, Shurey S, Ghatei MA, Patel AG, Bloom SR. Gut hormone profiles following bariatric surgery favor an anorectic state, facilitate weight loss, and improve metabolic parameters. *Annals of surgery*. 2006 Jan;243(1):108.
32. Ro L, Chang K. Neuropathic pain: mechanisms and treatments. *Chang Gung medical journal*. 2005 Sep 1;28(9):597.
33. Kekäläinen P, Juuti M, Walle T, Laatikainen T. Pregnancy planning in type 1 diabetic women improves glycemic control and pregnancy outcomes. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2016 Jul 17;29(14):2252-8.
34. Rigazio S, Lehto HR, Tuunanen H, Nagren K, Kankaanpää M, Simi C, Borra R, Naum AG, Parkkola R, Knuuti J, Nuutila P. The lowering of hepatic fatty acid uptake improves liver function and insulin sensitivity without affecting hepatic fat content in humans. *American Journal of Physiology-Endocrinology and Metabolism*. 2008 Aug;295(2):E413-9.
35. Getty AK, Wisdo TR, Chavis LN, Derella CC, McLaughlin KC, Perez AN, DiCiurcio III WT, Corbin M, Fairheller DL. Effects of circuit exercise training on vascular health and blood pressure. *Preventive Medicine Reports*. 2018 Jun 1;10:106-12.
36. Naaman SC, Bakris GL. Diabetic nephropathy: update on pillars of therapy slowing progression. *Diabetes Care*. 2023 Sep 1;46(9):1574-86.
37. Oberg E, Gidlöf S, Jakson I, Mitsell M, Tollet Egnell P, Hirschberg AL. Improved menstrual function in obese women with polycystic ovary syndrome after behavioural modification intervention—A randomized controlled trial. *Clinical Endocrinology*. 2019 Mar;90(3):468-78.
38. DeFronzo RA, Abdul-Ghani M. Type 2 diabetes can be prevented with early pharmacological intervention. *Diabetes care*. 2011 May 1;34(Supplement_2):S202-9.
39. Mandal SC, Nayak AK, Dhara AK, editors. *Herbal biomolecules in Healthcare applications*. Academic press; 2021 Oct 5.

How to Cite this article:

Yadav R, Kumar S, Baghel S, Kumar GP, Malik J, Singh SP, Singh G, Siroliya V. Unlocking the combined therapeutic potential of *Allium sativum* and *Swertia chirata*: a comprehensive review on synergistic antioxidant and antidiabetic properties. *International Journal of Pharmaceutical Science and Medicine* 2024; 2(1): 15-20.