

FINAL JEE–MAIN EXAMINATION – MARCH, 2021

(Held On Thursday 18th March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

PHYSICS

TEST PAPER WITH ANSWER & SOLUTION

SECTION-A

1. Which of the following statements are correct?
- (A) Electric monopoles do not exist whereas magnetic monopoles exist.
- (B) Magnetic field lines due to a solenoid at its ends and outside cannot be completely straight and confined.
- (C) Magnetic field lines are completely confined within a toroid.
- (D) Magnetic field lines inside a bar magnet are not parallel.
- (E) $\chi = -1$ is the condition for a perfect diamagnetic material, where χ is its magnetic susceptibility.

Choose the correct answer from the options given below :

- (1) (C) and (E) only
- (2) (B) and (D) only
- (3) (A) and (B) only
- (4) (B) and (C) only

2. An object of mass m_1 collides with another object of mass m_2 , which is at rest. After the collision the objects move with equal speeds in opposite direction. The ratio of the masses $m_2 : m_1$ is :
- (1) 3 : 1 (2) 2 : 1
- (3) 1 : 2 (4) 1 : 1

3. For an adiabatic expansion of an ideal gas, the fractional change in its pressure is equal to (where γ is the ratio of specific heats):

- (1) $-\gamma \frac{dV}{V}$ (2) $-\gamma \frac{V}{dV}$
- (3) $-\frac{1}{\gamma} \frac{dV}{V}$ (4) $\frac{dV}{V}$

4. A proton and an α -particle, having kinetic energies K_p and K_α , respectively, enter into a magnetic field at right angles.

The ratio of the radii of trajectory of proton to that of α -particle is 2 : 1. The ratio of $K_p : K_\alpha$ is :

- (1) 1 : 8 (2) 8 : 1
(3) 1 : 4 (4) 4 : 1


5. A plane electromagnetic wave propagating along y-direction can have the following pair of electric field (\vec{E}) and magnetic field (\vec{B}) components.

- (1) E_y, B_y or E_z, B_z
(2) E_y, B_x or E_x, B_y
(3) E_x, B_z or E_z, B_x
(4) E_x, B_y or E_y, B_x


6. Consider a uniform wire of mass M and length L . It is bent into a semicircle. Its moment of inertia about a line perpendicular to the plane of the wire passing through the centre is :

- (1) $\frac{1}{4} \frac{ML^2}{\pi^2}$ (2) $\frac{2}{5} \frac{ML^2}{\pi^2}$
(3) $\frac{ML^2}{\pi^2}$ (4) $\frac{1}{2} \frac{ML^2}{\pi^2}$

7. The velocity-displacement graph of a particle is shown in the figure.



The acceleration-displacement graph of the same particle is represented by :



10. If the angular velocity of earth's spin is increased such that the bodies at the equator start floating, the duration of the day would be approximately :
(Take : $g = 10 \text{ ms}^{-2}$, the radius of earth, $R = 6400 \times 10^3 \text{ m}$, Take $\pi = 3.14$)
- (1) 60 minutes
 - (2) does not change
 - (3) 1200 minutes
 - (4) 84 minutes

11. The decay of a proton to neutron is :
(1) not possible as proton mass is less than the neutron mass
(2) possible only inside the nucleus
(3) not possible but neutron to proton conversion is possible
(4) always possible as it is associated only with β^+ decay
12. In a series LCR circuit, the inductive reactance (X_L) is 10Ω and the capacitive reactance (X_C) is 4Ω . The resistance (R) in the circuit is 6Ω . The power factor of the circuit is :
- (1) $\frac{1}{2}$
 - (2) $\frac{1}{2\sqrt{2}}$
 - (3) $\frac{1}{\sqrt{2}}$
 - (4) $\frac{\sqrt{3}}{2}$


13. The angular momentum of a planet of mass M moving around the sun in an elliptical orbit is \vec{L} . The magnitude of the areal velocity of the planet is :

- (1) $\frac{4L}{M}$ (2) $\frac{L}{M}$
(3) $\frac{2L}{M}$ (4) $\frac{L}{2M}$

14. The function of time representing a simple harmonic motion with a period of $\frac{\pi}{\omega}$ is :

- (1) $\sin(\omega t) + \cos(\omega t)$
(2) $\cos(\omega t) + \cos(2\omega t) + \cos(3\omega t)$
(3) $\sin^2(\omega t)$
(4) $3\cos\left(\frac{\pi}{4} - 2\omega t\right)$


15. A solid cylinder of mass m is wrapped with an inextensible light string and, is placed on a rough inclined plane as shown in the figure. The frictional force acting between the cylinder and the inclined plane is :



[The coefficient of static friction, μ_s , is 0.4]

- (1) $\frac{7}{2}mg$ (2) $5mg$
(3) $\frac{mg}{5}$ (4) 0

7. A particle of mass m moves in a circular orbit under the central potential field, $U(r) = \frac{-C}{r}$, where C is a positive constant. The correct radius – velocity graph of the particle's motion is :



16. The time taken for the magnetic energy to reach 25% of its maximum value, when a solenoid of resistance R , inductance L is connected to a battery, is :

- (1) $\frac{L}{R} \ln 5$ (2) infinite
(3) $\frac{L}{R} \ln 2$ (4) $\frac{L}{R} \ln 10$

18. An ideal gas in a cylinder is separated by a piston in such a way that the entropy of one part is S_1 and that of the other part is S_2 . Given that $S_1 > S_2$. If the piston is removed then the total entropy of the system will be :

- (1) $S_1 \times S_2$ (2) $S_1 - S_2$
(3) $\frac{S_1}{S_2}$ (4) $S_1 + S_2$

19. Consider a sample of oxygen behaving like an ideal gas. At 300 K, the ratio of root mean square (rms) velocity to the average velocity of gas molecule would be :
(Molecular weight of oxygen is 32 g/mol; $R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$)


- (1) $\sqrt{\frac{3}{8}}$ (2) $\sqrt{\frac{8}{3}}$
(3) $\sqrt{\frac{3\pi}{8}}$ (4) $\sqrt{\frac{8\pi}{3}}$

20. The speed of electrons in a scanning electron microscope is $1 \times 10^7 \text{ ms}^{-1}$. If the protons having the same speed are used instead of electrons, then the resolving power of scanning proton microscope will be changed by a factor of:

- (1) 1837 (2) $\frac{1}{1837}$
(3) $\sqrt{1837}$ (4) $\frac{1}{\sqrt{1837}}$

SECTION-B


1. The projectile motion of a particle of mass 5 g is shown in the figure.



The initial velocity of the particle is $5\sqrt{2} \text{ ms}^{-1}$ and the air resistance is assumed to be negligible. The magnitude of the change in momentum between the points A and B is $x \times 10^{-2} \text{ kgms}^{-1}$. The value of x, to the nearest integer, is _____.


2. A ball of mass 4 kg, moving with a velocity of 10 ms^{-1} , collides with a spring of length 8 m and force constant 100 Nm^{-1} . The length of the compressed spring is x m. The value of x , to the nearest integer, is_____.

3. The typical output characteristics curve for a transistor working in the common-emitter configuration is shown in the figure.



The estimated current gain from the figure is

4. Consider a water tank as shown in the figure. Its cross-sectional area is 0.4 m^2 . The tank has an opening B near the bottom whose cross-section area is 1 cm^2 . A load of 24 kg is applied on the water at the top when the height of the water level is 40 cm above the bottom, the velocity of water coming out the opening B is $v \text{ ms}^{-1}$. The value of v , to the nearest integer, is____. [Take value of g to be 10 ms^{-2}]



5. A TV transmission tower antenna is at a height of 20 m. Suppose that the receiving antenna is at.

- (i) ground level
(ii) a height of 5 m.

The increase in antenna range in case (ii) relative to case (i) is $n\%$.


The value of n , to the nearest integer, is .

6. The radius of a sphere is measured to be (7.50 ± 0.85) cm. Suppose the percentage error in its volume is x . The value of x , to the nearest x , is_____.

7. An infinite number of point charges, each carrying $1 \mu\text{C}$ charge, are placed along the y -axis at $y = 1 \text{ m}, 2 \text{ m}, 4 \text{ m}, 8 \text{ m}, \dots$. The total force on a 1 C point charge, placed at the origin, is $x \times 10^3 \text{ N}$. The value of x , to the nearest integer, is_____.

[Take $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2/\text{C}^2$]

8. Consider a 72 cm long wire AB as shown in the figure. The galvanometer jockey is placed at P on AB at a distance x cm from A. The galvanometer shows zero deflection.



The value of x , to the nearest integer, is

9. Two wires of same length and thickness having specific resistances $6\Omega \text{ cm}$ and $3\Omega \text{ cm}$ respectively are connected in parallel. The effective resistivity is $\rho \Omega \text{ cm}$. The value of ρ , to the nearest integer, is_____.
10. A galaxy is moving away from the earth at a speed of 286 kms^{-1} . The shift in the wavelength of a red line at 630 nm is $x \times 10^{-10} \text{ m}$. The value of x , to the nearest integer, is_____.
[Take the value of speed of light c , as $3 \times 10^8 \text{ ms}^{-1}$]

FINAL JEE-MAIN EXAMINATION – MARCH, 2021

(Held On Thursday 18th March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

CHEMISTRY

TEST PAPER WITH ANSWER & SOLUTION

SECTION-A

1. The oxidation states of nitrogen in NO, NO₂, N₂O and NO₃⁻ are in the order of :

- (1) NO₃⁻ > NO₂ > NO > N₂O
- (2) NO₂ > NO₃⁻ > NO > N₂O
- (3) N₂O > NO₂ > NO > NO₃⁻
- (4) NO > NO₂ > N₂O > NO₃⁻

2. In basic medium, H₂O₂ exhibits which of the following reactions ?

- (A) Mn²⁺ → Mn⁴⁺
- (B) I₂ → I⁻
- (C) PbS → PbSO₄

Choose the most appropriate answer from the options given below :

- (1) (A), (C) only
- (2) (A) only
- (3) (B) only
- (4) (A), (B) only





3. In the reaction of hypobromite with amide, the carbonyl carbon is lost as :

- (1) CO₃²⁻
- (2) HCO₃⁻
- (3) CO₂
- (4) CO

4. The oxide that shows magnetic property is :

- (1) SiO₂
- (2) Mn₃O₄
- (3) Na₂O
- (4) MgO

5. Main Products formed during a reaction of 1-methoxy naphthalene with hydroiodic acid are:




6. Deficiency of vitamin K causes :
 (1) Increase in blood clotting time
 (2) Increase in fragility of RBC's
 (3) Cheilosis
 (4) Decrease in blood clotting time

7. An organic compound "A" on treatment with benzene sulphonyl chloride gives compound B. B is soluble in dil. NaOH solution. Compound A is :
 (1) $\text{C}_6\text{H}_5\text{-N-(CH}_3)_2$ (2) $\text{C}_6\text{H}_5\text{-NHCH}_2\text{CH}_3$
 (3) $\text{C}_6\text{H}_5\text{-CH}_2\text{NHCH}_3$ (4) $\text{C}_6\text{H}_5\text{-CH(NH}_2\text{)-CH}_3$

8. The first ionization energy of magnesium is smaller as compared to that of elements X and Y, but higher than that of Z. the elements X, Y and Z, respectively, are :
 (1) chlorine, lithium and sodium
 (2) argon, lithium and sodium
 (3) argon, chlorine and sodium
 (4) neon, sodium and chlorine

9. The secondary valency and the number of hydrogen bonded water molecule(s) in $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, respectively, are :
 (1) 6 and 4 (2) 4 and 1
 (3) 6 and 5 (4) 5 and 1

10. Given below are two statements :
 Statement I : Bohr's theory accounts for the stability and line spectrum of Li^+ ion.
 Statement II : Bohr's theory was unable to explain the splitting of spectral lines in the presence of a magnetic field.
 In the light of the above statements, choose the most appropriate answer from the options given below :
 (1) Both statement I and statement II are true.
 (2) Statement I is false but statement II is true.
 (3) Both statement I and statement II are false.
 (4) Statement I is true but statement II is false.




Consider the given reaction, percentage yield of :

- (1) $C > A > B$ (2) $B > C > A$
 (3) $A > C > B$ (4) $C > B > A$


12. The charges on the colloidal CdS sol and TiO_2 sol are, respectively :
- (1) positive and positive
 (2) positive and negative
 (3) negative and negative
 (4) negative and positive

13. Match List - I with List - II :

List - I	List - II
(Class of Chemicals)	(Example)
(a) Antifertility drug	(i) Meprobamate
(b) Antibiotic	(ii) Alitame
(c) Tranquilizer	(iii) Norethindrone
(d) Artificial Sweetener	(iv) Salvarsan
(1) (a)-(ii), (b)-(iii), (c)-(iv), (d)-(i)	
(2) (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)	
(3) (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii)	
(4) (a)-(ii), (b)-(iv), (c)-(i), (d)-(iii)	



Consider the above reaction, the product 'X' and 'Y' respectively are :



15. Match list-I with list-II :

List-I	List-II
(a) Be	(i) Treatment of cancer
(b) Mg	(ii) Extraction of metals
(c) Ca	(iii) Incendiary bombs and signals
(d) Ra	(iv) Windows of X-ray tubes
	(v) Bearings for motor engines.

Choose the most appropriate answer the option given below :

- (1) a-iv, b-iii, c-i, d-ii
- (2) a-iv, b-iii, c-ii, d-i
- (3) a-iii, b-iv, c-v, d-ii
- (4) a-iii, b-iv, c-ii, d-v

16. Given below are two statements :

Statement I : C_2H_5OH and $AgCN$ both can generate nucleophile.

Statement II : KCN and $AgCN$ both will generate nitrile nucleophile with all reaction conditions.

Choose the most appropriate option :

- (1) Statement I is true but statement II is false
- (2) Both statement I and statement II are true
- (3) Statement I is false but statement II is true
- (4) Both statement I and statement II are false

17. Given below are two statements :

Statement I : Non-biodegradable wastes are generated by the thermal power plants.

Statement II : Bio-degradable detergents leads to eutrophication.

In the light of the above statements, choose the most appropriate answer from the option given below :

- (1) Both statement I and statement II are false
- (2) Statement I is true but statement II is false
- (3) Statement I is false but statement II is true
- (4) Both statement I and statement II are true.


18. Match list-I with list-II :

List-I	List-II
(a) Mercury	(i) Vapour phase refining
(b) Copper	(ii) Distillation refining
(c) Silicon	(iii) Electrolytic refining
(d) Nickel	(iv) Zone refining

Choose the most appropriate answer from the option given below :

- (1) a-i, b-iv, c-ii, d-iii
- (2) a-ii, b-iii, c-i, d-iv
- (3) a-ii, b-iii, c-iv, d-i
- (4) a-ii, b-iv, c-iii, d-i

19. In the following molecules,



Hybridisation of carbon a, b and c respectively are :

- (1) sp^3 , sp , sp
- (2) sp^3 , sp^2 , sp
- (3) sp^3 , sp^2 , sp^2
- (4) sp^3 , sp , sp^2

20. A hard substance melts at high temperature and is an insulator in both solid and in molten state. This solid is most likely to be a / an :

- (1) Ionic solid
- (2) Molecular solid
- (3) Metallic solid
- (4) Covalent solid

SECTION-B

1. A reaction has a half life of 1 min. The time required for 99.9% completion of the reaction is _____ min. (Round off to the Nearest integer)
[Use : $\ln 2 = 0.69$, $\ln 10 = 2.3$]

2. The molar conductivities at infinite dilution of barium chloride, sulphuric acid and hydrochloric acid are 280, 860 and 426 $\text{Scm}^2 \text{mol}^{-1}$ respectively. The molar conductivity at infinite dilution of barium sulphate is _____ $\text{S cm}^2 \text{mol}^{-1}$ (Round off to the Nearest Integer).

3. The number of species below that have two lone pairs of electrons in their central atom is _____ (Round off to the Nearest integer)
 SF_4 , BF_4^- , ClF_3 , AsF_3 , PCl_5 , BrF_5 , XeF_4 , SF_6

4. A xenon compound 'A' upon partial hydrolysis gives XeO_2F_2 . The number of lone pair of electrons present in compound A is _____ (Round off to the Nearest integer)

5. The gas phase reaction

$$2\text{A}(\text{g}) \rightleftharpoons \text{A}_2(\text{g})$$
 at 400 K has $\Delta G^\circ = + 25.2 \text{ kJ mol}^{-1}$.
 The equilibrium constant K_C for this reaction is _____ $\times 10^{-2}$. (Round off to the Nearest integer)
 [Use : $R = 8.3 \text{ J mol}^{-1}\text{K}^{-1}$, $\ln 10 = 2.3$
 $\log_{10} 2 = 0.30$, 1 atm = 1 bar]
 [antilog $(-0.3) = 0.501$]


6. In Tollen's test for aldehyde, the overall number of electron(s) transferred to the Tollen's reagent formula $[\text{Ag}(\text{NH}_3)_2]^+$ per aldehyde group to form silver mirror is _____. (Round off to the Nearest integer)
7. The solubility of CdSO_4 in water is $8.0 \times 10^{-4} \text{ mol L}^{-1}$. Its solubility in 0.01 M H_2SO_4 solution is _____ $\times 10^{-6} \text{ mol L}^{-1}$. (Round off to the Nearest integer) (Assume that solubility is much less than 0.01 M)
8. A solute A dimerizes in water. The boiling point of a 2 molar solution of A is 100.52°C . The percentage association of A is _____. (Round off to the Nearest integer)
[Use : K_b for water = $0.52 \text{ K kg mol}^{-1}$
Boiling point of water = 100°C]

9. 10.0 ml of Na_2CO_3 solution is titrated against 0.2 M HCl solution. The following titre values were obtained in 5 readings.

4.8 ml, 4.9 ml, 5.0 ml, 5.0 ml and 5.0 ml

Based on these readings, and convention of titrimetric estimation of concentration of Na_2CO_3 solution is _____ mM.

(Round off to the Nearest integer)



Consider the above reaction where 6.1 g of benzoic acid is used to get 7.8 g of m-bromo benzoic acid. The percentage yield of the product is_____.

(Round off to the Nearest integer)

[Given : Atomic masses : C = 12.0u, H : 1.0u, O : 16.0u, Br = 80.0 u]

Official Ans. by NTA (78)

FINAL JEE–MAIN EXAMINATION – MARCH, 2021

(Held On Thursday 18th March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

MATHEMATICS

SECTION-A

1. Let $y = y(x)$ be the solution of the differential equation $\frac{dy}{dx} = (y+1)((y+1)e^{x^{2/2}} - x)$, $0 < x < 2.1$,

with $y(2) = 0$. Then the value of $\frac{dy}{dx}$ at $x = 1$ is equal to :

- (1) $\frac{-e^{3/2}}{(e^2 + 1)^2}$ (2) $-\frac{2e^2}{(1 + e^2)^2}$
(3) $\frac{e^{5/2}}{(1 + e^2)^2}$ (4) $\frac{5e^{1/2}}{(e^2 + 1)^2}$

TEST PAPER WITH SOLUTION

2. In a triangle ABC, if $|\overline{BC}| = 8$, $|\overline{CA}| = 7$, $|\overline{AB}| = 10$, then the projection of the vector \overline{AB} on \overline{AC} is equal to :

- (1) $\frac{25}{4}$ (2) $\frac{85}{14}$ (3) $\frac{127}{20}$ (4) $\frac{115}{16}$

3. Let the system of linear equations

$$4x + \lambda y + 2z = 0$$

$$2x - y + z = 0$$

$$\mu x + 2y + 3z = 0, \lambda, \mu \in \mathbb{R}.$$

has a non-trivial solution. Then which of the following is true ?

- (1) $\mu = 6, \lambda \in \mathbb{R}$ (2) $\lambda = 2, \mu \in \mathbb{R}$
(3) $\lambda = 3, \mu \in \mathbb{R}$ (4) $\mu = -6, \lambda \in \mathbb{R}$

4. Let $f : \mathbb{R} - \{3\} \rightarrow \mathbb{R} - \{1\}$ be defined by

$$f(x) = \frac{x-2}{x-3}. \text{ Let } g : \mathbb{R} \rightarrow \mathbb{R} \text{ be given as}$$

$$g(x) = 2x - 3. \text{ Then, the sum of all the values}$$

of x for which $f^{-1}(x) + g^{-1}(x) = \frac{13}{2}$ is equal to

- (1) 7 (2) 2 (3) 5 (4) 3

5. Let the centroid of an equilateral triangle ABC be at the origin. Let one of the sides of the equilateral triangle be along the straight line $x + y = 3$. If R and r be the radius of circumcircle and incircle respectively of $\triangle ABC$, then $(R + r)$ is equal to :

- (1) $\frac{9}{\sqrt{2}}$ (2) $7\sqrt{2}$ (3) $2\sqrt{2}$ (4) $3\sqrt{2}$

6. Consider a hyperbola $H : x^2 - 2y^2 = 4$. Let the tangent at a point $P(4, \sqrt{6})$ meet the x -axis at Q and latus rectum at $R(x_1, y_1)$, $x_1 > 0$. If F is a focus of H which is nearer to the point P , then the area of $\triangle QFR$ is equal to

- (1) $4\sqrt{6}$ (2) $\sqrt{6} - 1$
(3) $\frac{7}{\sqrt{6}} - 2$ (4) $4\sqrt{6} - 1$

7. If P and Q are two statements, then which of the following compound statement is a tautology ?

- (1) $((P \Rightarrow Q) \wedge \sim Q) \Rightarrow Q$
- (2) $((P \Rightarrow Q) \wedge \sim Q) \Rightarrow \sim P$
- (3) $((P \Rightarrow Q) \wedge \sim Q) \Rightarrow P$
- (4) $((P \Rightarrow Q) \wedge \sim Q) \Rightarrow (P \wedge Q)$

8. Let $g(x) = \int_0^x f(t) dt$, where f is continuous

function in $[0, 3]$ such that $\frac{1}{3} \leq f(t) \leq 1$ for all

$t \in [0, 1]$ and $0 \leq f(t) \leq \frac{1}{2}$ for all $t \in (1, 3]$.

The largest possible interval in which $g(3)$ lies is :

- (1) $\left[-1, -\frac{1}{2}\right]$
- (2) $\left[-\frac{3}{2}, -1\right]$
- (3) $\left[\frac{1}{3}, 2\right]$
- (4) $[1, 3]$

9. Let S_1 be the sum of first $2n$ terms of an arithmetic progression. Let S_2 be the sum of first $4n$ terms of the same arithmetic progression. If $(S_2 - S_1)$ is 1000, then the sum of the first $6n$ terms of the arithmetic progression is equal to:
(1) 1000 (2) 7000 (3) 5000 (4) 3000
10. Let a complex number be $w = 1 - \sqrt{3}i$. Let another complex number z be such that $|zw| = 1$ and $\arg(z) - \arg(w) = \frac{\pi}{2}$. Then the area of the triangle with vertices origin, z and w is equal to :
(1) 4 (2) $\frac{1}{2}$ (3) $\frac{1}{4}$ (4) 2
11. Let in a series of $2n$ observations, half of them are equal to a and remaining half are equal to $-a$. Also by adding a constant b in each of these observations, the mean and standard deviation of new set become 5 and 20, respectively. Then the value of $a^2 + b^2$ is equal to :
(1) 425 (2) 650 (3) 250 (4) 925

12. Let $S_1 : x^2 + y^2 = 9$ and $S_2 : (x - 2)^2 + y^2 = 1$. Then the locus of center of a variable circle S which touches S_1 internally and S_2 externally always passes through the points :
- (1) $(0, \pm\sqrt{3})$ (2) $\left(\frac{1}{2}, \pm\frac{\sqrt{5}}{2}\right)$
(3) $\left(2, \pm\frac{3}{2}\right)$ (4) $(1, \pm 2)$
13. Let \vec{a} and \vec{b} be two non-zero vectors perpendicular to each other and $|\vec{a}| = |\vec{b}|$. If $|\vec{a} \times \vec{b}| = |\vec{a}|$, then the angle between the vectors $(\vec{a} + \vec{b} + (\vec{a} \times \vec{b}))$ and \vec{a} is equal to :
- (1) $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (2) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$
(3) $\cos^{-1}\left(\frac{1}{\sqrt{2}}\right)$ (4) $\sin^{-1}\left(\frac{1}{\sqrt{6}}\right)$
14. Let in a Binomial distribution, consisting of 5 independent trials, probabilities of exactly 1 and 2 successes be 0.4096 and 0.2048 respectively. Then the probability of getting exactly 3 successes is equal to :
- (1) $\frac{32}{625}$ (2) $\frac{80}{243}$ (3) $\frac{40}{243}$ (4) $\frac{128}{625}$

15. Let a tangent be drawn to the ellipse $\frac{x^2}{27} + y^2 = 1$

at $(3\sqrt{3}\cos\theta, \sin\theta)$ where $\theta \in \left(0, \frac{\pi}{2}\right)$. Then the

value of θ such that the sum of intercepts on axes made by this tangent is minimum is equal to :

- (1) $\frac{\pi}{8}$ (2) $\frac{\pi}{4}$ (3) $\frac{\pi}{6}$ (4) $\frac{\pi}{3}$

16. Define a relation R over a class of $n \times n$ real matrices A and B as "ARB iff there exists a non-singular matrix P such that $PAP^{-1} = B$ ".

Then which of the following is true ?

- (1) R is symmetric, transitive but not reflexive,
(2) R is reflexive, symmetric but not transitive
(3) R is an equivalence relation
(4) R is reflexive, transitive but not symmetric

17. A pole stands vertically inside a triangular park ABC. Let the angle of elevation of the top of

the pole from each corner of the park be $\frac{\pi}{3}$.

If the radius of the circumcircle of ΔABC is 2, then the height of the pole is equal to :

- (1) $\frac{2\sqrt{3}}{3}$ (2) $2\sqrt{3}$ (3) $\sqrt{3}$ (4) $\frac{1}{\sqrt{3}}$

18. If $15\sin^4\alpha + 10\cos^4\alpha = 6$, for some $\alpha \in \mathbb{R}$, then the value of $27\sec^6\alpha + 8\operatorname{cosec}^6\alpha$ is equal to :
(1) 350 (2) 500 (3) 400 (4) 250

19. The area bounded by the curve $4y^2 = x^2(4 - x)(x - 2)$ is equal to :

- (1) $\frac{\pi}{8}$ (2) $\frac{3\pi}{8}$ (3) $\frac{3\pi}{2}$ (4) $\frac{\pi}{16}$

20. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined as

$$f(x) = \begin{cases} \frac{\sin(a+1)x + \sin 2x}{2x}, & \text{if } x < 0 \\ b, & \text{if } x = 0 \\ \frac{\sqrt{x + bx^3} - \sqrt{x}}{bx^{5/2}}, & \text{if } x > 0 \end{cases}$$

If f is continuous at $x = 0$, then the value of $a + b$ is equal to :

- (1) $-\frac{5}{2}$ (2) -2 (3) -3 (4) $-\frac{3}{2}$

SECTION-B

1. If $f(x)$ and $g(x)$ are two polynomials such that the polynomial $P(x) = f(x^3) + xg(x^3)$ is divisible by $x^2 + x + 1$, then $P(1)$ is equal to_____.

2. Let I be an identity matrix of order 2×2 and $P = \begin{bmatrix} 2 & -1 \\ 5 & -3 \end{bmatrix}$. Then the value of $n \in \mathbb{N}$ for which $P^n = 5I - 8P$ is equal to _____.

3. If $\sum_{r=1}^{10} r!(r^3 + 6r^2 + 2r + 5) = \alpha(11!)$, then the value of α is equal to _____.

4. The term independent of x in the expansion of $\left[\frac{x+1}{x^{2/3} - x^{1/3} + 1} - \frac{x-1}{x - x^{1/2}} \right]^{10}$, $x \neq 1$, is equal to _____.

5. Let $P(x)$ be a real polynomial of degree 3 which vanishes at $x = -3$. Let $P(x)$ have local minima at $x = 1$, local maxima at $x = -1$ and $\int_{-1}^1 P(x) dx = 18$, then the sum of all the coefficients of the polynomial $P(x)$ is equal to _____.

6. Let the mirror image of the point $(1, 3, a)$ with respect to the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) - b = 0$ be $(-3, 5, 2)$. Then the value of $|a + b|$ is equal to _____.

7. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfy the equation $f(x + y) = f(x) \cdot f(y)$ for all $x, y \in \mathbb{R}$ and $f(x) \neq 0$ for any $x \in \mathbb{R}$. If the function f is differentiable at $x = 0$ and $f'(0) = 3$, then

$$\lim_{h \rightarrow 0} \frac{1}{h} (f(h) - 1) \text{ is equal to } \underline{\hspace{2cm}}.$$

8. Let nC_r denote the binomial coefficient of x^r in the expansion of $(1 + x)^n$.

If $\sum_{k=0}^{10} (2^2 + 3k) {}^nC_k = \alpha \cdot 3^{10} + \beta \cdot 2^{10}$, $\alpha, \beta \in \mathbb{R}$, then $\alpha + \beta$ is equal to _____.

9. Let P be a plane containing the line

$$\frac{x-1}{3} = \frac{y+6}{4} = \frac{z+5}{2} \text{ and parallel to the line}$$

$$\frac{x-3}{4} = \frac{y-2}{-3} = \frac{z+5}{7}. \text{ If the point } (1, -1, \alpha) \text{ lies}$$

on the plane P, then the value of 15α is equal to _____.

10. Let $y = y(x)$ be the solution of the differential

equation $x dy - y dx = \sqrt{x^2 - y^2} dx$, $x \geq 1$, with

$y(1) = 0$. If the area bounded by the line $x = 1$, $x = e^\pi$, $y = 0$ and $y = y(x)$ is $\alpha e^{2\pi} + \beta$, then the value of $10(\alpha + \beta)$ is equal to _____.