

JEE Main 2020 Paper

Date of Exam: **8th January (Shift I)**

Time: **9:30 am – 12:30 pm**

Subject: **Physics**

1. A particle of mass m is fixed to one end of a light spring having force constant k and unstretched length l . The other end is fixed. The system is given an angular speed ω about the fixed end of the spring such that it rotates in a circle in gravity free space. Then the stretch in the spring is

- | | |
|-------------------------------------|-------------------------------------|
| a. $\frac{lm\omega^2}{k-m\omega^2}$ | b. $\frac{lm\omega^2}{k+m\omega^2}$ |
| c. $\frac{lm\omega^2}{k-m\omega}$ | d. $\frac{lm\omega^2}{k+m\omega}$ |


Solution: (a)

The centripetal force is provided by the spring force.


$$m\omega^2(l + x) = kx$$

$$\left(\frac{l}{x} + 1\right) = \frac{k}{m\omega^2}$$

$$x = \frac{lm\omega^2}{k-m\omega^2}$$



2. Three charged particles A, B and, C with charge $-4q$, $+2q$ and $-2q$ are present on the circumference of a circle of radius d . The charges particles A, C and centre O of the circle formed an equilateral triangle as shown in figure. Electric field at O along x- direction is:



JEE Main 2020 Paper

a. $\frac{\sqrt{3}q}{4\pi\epsilon_0 d^2}$

b. $\frac{2\sqrt{3}q}{\pi\epsilon_0 d^2}$


c. $\frac{\sqrt{3}q}{\pi\epsilon_0 d^2}$

d. $\frac{3\sqrt{3}q}{4\pi\epsilon_0 d^2}$

Solution: (c)

Applying superposition principle,


$$\vec{E}_{net} = \vec{E}_1 + \vec{E}_2 + \vec{E}_3$$



By symmetry, net electric field along the x-axis.


$$|\vec{E}_{net}| = \frac{4kq}{d^2} \times 2 \cos 30^\circ = \frac{\sqrt{3}q}{\pi\epsilon_0 d^2}$$

3. A thermodynamic cycle $xyzx$ is shown on a $V - T$ diagram .



The P-V diagram that best describes this cycle is : (Diagrams are schematic and not upto scale)

JEE Main 2020 Paper



Solution: (c)


For the given $V - T$ graph

For the process $x \rightarrow y$; $V \propto T$; $P = \text{constant}$

For the process $y \rightarrow z$; $V = \text{constant}$

Only 'c' satisfies these two conditions.

4. Find the co-ordinates of center of mass of the lamina shown in the figure below.




- | | |
|---------------------|---------------------|
| a. (0.75 m, 1.75 m) | b. (0.75 m, 0.75 m) |
| c. (1.25 m, 1.5 m) | d. (1 m, 1.75 m) |

JEE Main 2020 Paper

Solution: (a)

The Lamina can be divided into two parts having equal mass m each.




$$\vec{r}_{\text{cm}} = \frac{m \times \left(\frac{\hat{i}}{2} + \hat{j}\right) + m \times \left(\hat{i} + \frac{5\hat{j}}{2}\right)}{2m}$$


$$\vec{r}_{\text{cm}} = \frac{3}{4}\hat{i} + \frac{7}{4}\hat{j}$$

5. The plot that depicts the behavior of the mean free time τ (time between two successive collisions) for the molecules of an ideal gas, as a function of temperature (T), qualitatively, is: (Graph are schematic and not drawn to scale)


a.




b.



c.



d.




Solution: (a)

JEE Main 2020 Paper

$$\tau \propto \frac{1}{\sqrt{T}}$$

6. Effective capacitance of parallel combination of two capacitors C_1 and C_2 is $10\ \mu\text{F}$. When these capacitor are individually connects to a voltage source of 1 V , the energy stored in the capacitor C_2 is 4 times of that in C_1 . If these capacitors are connected in series, their effective capacitance will be:
- a. $1.6\ \mu\text{F}$
- b. $3.2\ \mu\text{F}$
- c. $4.2\ \mu\text{F}$
- d. $8.4\ \mu\text{F}$

Solution: (a)



Given that,

$$C_1 + C_2 = 10 \mu F \quad \dots(i)$$

$$4 \left(\frac{1}{2} C_1 V^2 \right) = \frac{1}{2} C_2 V^2$$

$$\Rightarrow 4C_1 = C_2 \quad \dots(\text{ii})$$

From equations (i) and (ii)

$$C_1 = 2 \mu F$$

$$C_2 = 8 \mu F$$


If they are in series

$$C_{eq} = \frac{C_1 C_2}{C_1 + C_2} = 1.6 \text{ } \mu\text{F}$$

7. Consider a uniform rod of mass $4m$ and length L pivoted about its centre. A mass m is moving with a velocity v making angle $\theta = \frac{\pi}{4}$ to the rod's long axis collides with one end of the rod and stick to it.. The angular speed of the rod-mass system just after collision is
- a. $\frac{3\sqrt{2}v}{7L}$
- b. $\frac{4v}{7L}$
- c. $\frac{3v}{7\sqrt{2}L}$
- d. $\frac{3v}{7L}$

JEE Main 2020 Paper

Solution: (a)



There is no external torque on the system about the hinge point.

So,

$$\vec{L}_i = \vec{L}_f$$

$$\frac{mv}{\sqrt{2}} \times \frac{1}{2} = \left[\frac{4mL^2}{12} + \frac{mL^2}{4} \right] \times \omega$$

$$\omega = \frac{6v}{7\sqrt{2}L} = \frac{3\sqrt{2}v}{7L}$$

8. When photons of energy 4 eV strikes the surface of a metal A, the ejected photoelectrons have maximum kinetic energy T_A eV and de-Broglie wavelength λ_A . The maximum kinetic energy of photoelectrons liberated from another metal B by photon of energy 4.50 eV is $T_B = (T_A - 1.5)$ eV. If the de-Broglie wavelength of these photoelectrons $\lambda_B = 2\lambda_A$, then the work function of metal B is

- | | |
|---------|-----------|
| a. 3 eV | b. 1.5 eV |
| c. 2 eV | d. 4 eV |

Solution: (d)

$$\lambda = \frac{h}{\sqrt{2(KE)m_B}} = \lambda \propto \frac{1}{\sqrt{KE}}$$

JEE Main 2020 Paper

$$\frac{\lambda_A}{\lambda_B} = \frac{\sqrt{KE_B}}{\sqrt{KE_A}}$$

$$\frac{1}{2} = \sqrt{\frac{T_A - 1.5}{T_A}}$$


$$T_A = 2 \text{ eV}$$

$$\text{KE}_B = 2 - 1.5 = 0.5 \text{ eV}$$

$$\phi_B = 4.5 - 0.5 = 4 \text{ eV}$$

9. The length of a potentiometer wire of length 1200 cm and it carries a current of 60 mA . For a cell of emf 5 V and internal resistance of $20\ \Omega$, the null point on it is found to be at 1000 cm . The resistance of whole wire is
- | | |
|------------------|------------------|
| a. $80\ \Omega$ | b. $100\ \Omega$ |
| c. $120\ \Omega$ | d. $60\ \Omega$ |

Solution: (b)



Assume the terminal voltage of the primary battery as V_p . As long as this potentiometer is operating on balanced length, V_p will remain constant.

As we know, potential gradient = $\frac{5}{1000} = \frac{V_p}{1200}$

$$V_p = 6 \text{ V}$$

$$\text{And } R_p = \frac{V_p}{i} = \frac{6}{60 \times 10^{-3}} = 100 \, \Omega$$

10. The magnifying power of a telescope with tube length 60 cm is 5. What is the focal length of its eyepiece?

JEE Main 2020 Paper

- a. 10 cm
c. 30 cm

- b. 20 cm
d. 40 cm

Solution: (a)

$$m = \frac{f_0}{f_e} = 5$$


$$\Rightarrow f_0 = 5f_e$$

$$f_0 + f_e = 5f_e + f_e = 6f_e = \text{length of the tube}$$

$$\Rightarrow 6f_e = 60 \text{ cm}$$

$$\Rightarrow f_e = 10 \text{ cm}$$

11. Consider two solid spheres of radii $R_1 = 1 \text{ m}$, $R_2 = 2 \text{ m}$ and masses M_1 & M_2 , respectively. The gravitational field due to two spheres 1 and 2 are shown. The value of $\frac{M_1}{M_2}$ is



a. $\frac{1}{6}$

b. $\frac{1}{3}$

c. $\frac{1}{2}$

d. $\frac{2}{3}$

Solution: (a)

Gravitation field will be maximum at the surface of a sphere. Therefore,


$$\frac{GM_2}{2^2} = 3 \text{ \& \; } \frac{GM_1}{1^2} = 2$$

$$\Rightarrow \frac{M_2}{M_1} \times \frac{1}{4} = \frac{3}{2}$$

$$\Rightarrow \frac{M_1}{M_2} = \frac{1}{6}$$

JEE Main 2020 Paper

12. Proton with kinetic energy of 1 MeV moves from south to north. It gets an acceleration of 10^{12} m/s^2 by an applied magnetic field (west to east). The value of magnetic field: (Rest mass of proton is $1.6 \times 10^{-27} \text{ kg}$)



- a. 0.71 mT b. 7.1 mT
c. 71 mT d. 0.071 mT

Solution: (a)

$$\text{K.E} = 1 \times 10^6 \text{ eV} = 1.6 \times 10^{-13} \text{ J}$$

$$= \frac{1}{2} m_e v^2$$

Where m_e is the mass of the electron = 1.6×10^{-27} kg

$$\Rightarrow 1.6 \times 10^{-13} = \frac{1}{2} \times 1.6 \times 10^{-27} \times v^2$$

$$\therefore v = \sqrt{2} \times 10^7 \text{ m/s}$$

$$Bqv = m_e a$$

$$\therefore B = \frac{1.6 \times 10^{-27} \times 10^{12}}{1.6 \times 10^{-19} \times \sqrt{2} \times 10^7}$$

$$= 0.71 \times 10^{-3} \text{T} = 0.71 \text{ mT}$$

13. If finding the electric field around a surface is given by $|\vec{E}| = \frac{q_{\text{enclosed}}}{\epsilon_0 |A|}$ is applicable. In the formula ϵ_0 is permittivity of free space, A is area of Gaussian and q_{enc} is charge enclosed by the Gaussian surface. This equation can be used in which of the following equation?
- Only when the Gaussian surface is an equipotential surface.
 - Only when $|\vec{E}| = \text{constant}$ on the surface..
 - Equipotential surface and $|\vec{E}|$ is constant on the surface .

JEE Main 2020 Paper

d. for any choice of Gaussian surfaces.

Solution: (c)

The magnitude of the electric field is constant and the electric field must be along the area vector i.e. the surface is equipotential.

14. The dimension of stopping potential V_0 in photoelectric effect in units of Planck's constant (h), speed of light (c), and gravitational constant (G) and Ampere (A) is

a. $h^{2/3}c^{5/3}G^{1/3}A^{-1}$

b. $h^2c^{1/3}G^{3/2}A^{-1}$

c. $h^{1/3}G^{2/3}c^{1/3}A^{-1}$

d. $h^{-2/3}c^{-1/3}G^{4/3}A^{-1}$

Solution: (challenge question)

$$V = K(h)^a(I)^b(G)^c(c)^d \quad \text{Unit of stopping potential is } (V_0) \text{ Volt.}$$

$$\text{We know } [h] = ML^2T^{-1}$$

$$[I] = A$$

$$[G] = M^{-1}L^3T^{-2}$$

$$[C] = LT^{-1}$$

$$[V] = ML^2T^{-3}A^{-1}$$

$$ML^2T^{-3}A^{-1} = (ML^2T^{-1})^a(A)^b(M^{-1}L^3T^{-2})^c(LT^{-1})^d$$

$$ML^2T^{-3}A^{-1} = M^{a-c}L^{2a+3c+d}T^{-a-2c-d}A^b$$

$$a - c = 1$$

$$2a + 3c + d = 2$$

$$-a - 2c - d = -3$$

$$b = -1$$

On solving,

$$c = -1$$

$$a = 0$$

$$d = 5$$

$$b = -1$$

$$V = K(h)^0(I)^{-1}(G)^{-1}(C)^5$$

15. A leak proof cylinder of length 1 m, made of metal which has very low coefficient of expansion is floating in water at 0°C such that its height above the water surface is 20 cm. When the temperature of water is increases to 4°C , the height of the cylinder above the water surface becomes 21 cm. The density of water at $T = 4^\circ\text{C}$ relative to the density at $T = 0^\circ\text{C}$ is close to

a. 1.01


b. 1.03

c. 1.26

d. 1.04

Solution: (a)

JEE Main 2020 Paper



Since the cylinder is in equilibrium, its weight is balanced by the Buoyant force.

$$mg = A(80)(\rho_{0^\circ\text{C}})g$$

$$mg = A(79)(\rho_{4^\circ\text{C}})g$$



$$\frac{\rho_{4^\circ\text{C}}}{\rho_{0^\circ\text{C}}} = \frac{80}{79} = 1.01$$

16. The graph which depicts the result of Rutherford gold foil experiment with α -particle is:



θ : Scattering angle

N : Number of scattered α – particles is detected


(Plots are schematic and not to scale)



JEE Main 2020 Paper



Solution: (b)




$$N \propto \frac{1}{\sin^4(\theta/2)}$$

17. At time $t = 0$ magnetic field of 1000 Gauss is passing perpendicularly through the area defined by the closed loop shown in the figure. If the magnetic field reduces linearly to 500 Gauss, in the next 5 s, then induced EMF in the loop is:

- | | |
|---------------------|---------------------|
| a. 56 μV | b. 28 μV |
| c. 30 μV | d. 48 μV |


JEE Main 2020 Paper



Solution: (a)

$$\begin{aligned}\epsilon &= \left| -\frac{d\phi}{dt} \right| = \left| -\frac{AdB}{dt} \right| \\ &= (16 \times 4 - 4 \times 2) \frac{(1000 - 500)}{5} \times 10^{-4} \times 10^{-4} \\ &= 56 \times \frac{500}{5} \times 10^{-8} = 56 \times 10^{-6} \text{ V}\end{aligned}$$

18. Choose the correct Boolean expression for the given circuit diagram:



- a. $A.B$
c. $A + B$

- b. $\bar{A} + \bar{B}$
d. $\bar{A} . \bar{B}$

Solution: (d)

First part of figure shown OR gate and second part of figure shown NOT gate.

$$\text{So, } Y = \bar{A} + \bar{B} = \bar{A} . \bar{B}$$

19. Consider a solid sphere of density $\rho(r) = \rho_o \left(1 - \frac{r^2}{R^2}\right)$, $0 < r \leq R$. The minimum density of a liquid in which it float is just

- a. $\frac{2}{5}\rho_o$
c. $\frac{\rho_o}{5}$

- b. $\frac{2}{3}\rho_o$
d. $\frac{\rho_o}{3}$

Solution: (a)

JEE Main 2020 Paper

Let the mass of the sphere be m and the density of the liquid be ρ_L

$$\rho = \rho_o \left(1 - \frac{r^2}{R^2}\right), 0 < r \leq R$$

Since the sphere is floating in the liquid, buoyancy force (F_B) due to liquid will balance the weight of the sphere.

$$F_B = mg$$

$$\rho_L \frac{4}{3} \pi R^3 g = \int \rho (4\pi r^2 dr) g$$

$$\rho_L \frac{4}{3} \pi R^3 = \int \rho_o \left(1 - \frac{r^2}{R^2}\right) 4\pi r^2 dr$$

$$\rho_L \frac{4}{3} \pi R^3 = \int_0^R \rho_o 4\pi \left(r^2 - \frac{r^4}{R^2}\right) dr = \rho_o 4\pi \left(\frac{r^3}{3} - \frac{r^5}{5R^2}\right)_0^R$$

$$\rho_L = \frac{2}{5} \rho_o$$

20. The critical angle of a medium for a specific wavelength, if the medium has relative permittivity 3 and relative permeability $\frac{4}{3}$ for this wavelength, will be

- | | |
|---------------|---------------|
| a. 15° | b. 30° |
| c. 45° | d. 60° |

Solution: (b)

If the speed of light in the given medium is V then,

$$V = \frac{1}{\sqrt{\mu\epsilon}}$$

We know that, $n = \frac{c}{v}$

$$n = \sqrt{\mu_r \epsilon_r} = 2$$

$$\sin \theta_c = \frac{1}{2}$$

$$\theta_c = 30^\circ$$

21. A body of mass $m = 0.10 \text{ kg}$ has an initial velocity of $3\hat{i} \text{ m/s}$. It collides elastically with another body, B of the mass which has an initial velocity of $5\hat{j} \text{ m/s}$. After collision, A moves with a velocity $v = 4(\hat{i} + \hat{j}) \text{ m/s}$. The energy of B after collision is written as $\frac{x}{10} \text{ J}$, the value of x is

Solution: (1)(Challenge question)

JEE Main 2020 Paper

Mass of each object, $m_1 = m_2 = 0.1 \text{ kg}$

Initial velocity of 1st object, $u_1 = 5 \text{ m/s}$

Initial velocity of 2nd object, $u_2 = 3 \text{ m/s}$

Final velocity of 1st object, $V_1 = 4\hat{i} + 4\hat{j} \text{ m/s} = \sqrt{4^2 + 4^2} = 16\sqrt{2} \text{ m/s}$

For elastic collision, kinetic energy remains conserved

$$\text{Initial kinetic energy } (K_i) = \text{Final kinetic energy } (K_f)$$

$$\frac{1}{2}mu_1^2 + \frac{1}{2}mu_2^2 = \frac{1}{2}mV_1^2 + \frac{1}{2}mV_2^2$$

$$\frac{1}{2}m(5)^2 + \frac{1}{2}m(3)^2 = \frac{1}{2}m(16\sqrt{2})^2 + \frac{1}{2}mV_2^2$$

$$V_2 = \sqrt{2} \text{ m/s}$$

$$\text{Kinetic energy of second object} = \frac{1}{2}mV_2^2 = \frac{1}{2} \times 0.1 \times \sqrt{2}^2 = \frac{1}{10}$$

$$\Rightarrow x = 1$$

22. A point object in air is in front of the curved surface of a plano-convex lens. The radius of curvature of the curved surface is 30 cm and the refractive index of lens material is 1.5, then the focal length of the lens (in cm) is

Solution: (60 cm)

Applying Lens makers' formula,

$$\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$R_1 = \infty$$

$$R_2 = -30 \text{ cm}$$

$$\frac{1}{f} = (1.5 - 1) \left(\frac{1}{\infty} - \frac{1}{-30} \right)$$

$$\frac{1}{f} = \frac{0.5}{30}$$

$$f = 60 \text{ cm}$$

23. A particle is moving along the x-axis with its coordinate with time t given by $x(t) = -3t^2 + 8t + 10 \text{ m}$. Another particle is moving along the y-axis with its coordinate as a function of time given by $y = 5 - 8t^3 \text{ m}$. At $t = 1 \text{ s}$, the speed of the second particle as measured in the frame of the first particle is given as \sqrt{v} . Then $v \text{ (m/s)}$ is _____.

JEE Main 2020 Paper

Solution: (580 m/s)(Challenge question)

$$x = -3t^2 + 8t + 10$$

$$\begin{aligned}\vec{v}_A &= (-6t + 8)\hat{i} \\ &= 2\hat{i}\end{aligned}$$

$$y = 5 - 8t^3$$

$$\vec{v}_B = -24t^2\hat{j}$$

$$|\vec{v}_{B/A}| = |\vec{v}_B - \vec{v}_A| = |-24\hat{j} - 2\hat{i}|$$

$$v = \sqrt{24^2 + 2^2}$$

$$v = 580 \text{ m/s}$$

24. A one metre long (both ends open) organ pipe is kept in a gas that has double the density of air at STP. Assuming the speed of sound in air at STP is 300 m/s, the frequency difference between the fundamental and second harmonic of this pipe is ____ Hz.

Solution: (106.05 Hz)

$$V = \sqrt{\frac{B}{\rho}}$$

$$\frac{V_{\text{pipe}}}{V_{\text{air}}} = \frac{\sqrt{\frac{B}{2\rho}}}{\sqrt{\frac{B}{\rho}}} = \frac{1}{\sqrt{2}}$$

$$V_{\text{pipe}} = \frac{V_{\text{air}}}{\sqrt{2}}$$


$$f_n = \frac{(n+1)}{2l} V_{\text{pipe}}$$

$$f_1 - f_0 = \frac{V_{\text{pipe}}}{2l} = \frac{300}{2\sqrt{2}} = 106.06 \text{ Hz (if } \sqrt{2} = 1.414) \approx 106 \text{ Hz}$$

25. Four resistors of resistance 15 Ω , 12 Ω , 4 Ω and 10 Ω respectively in cyclic order to form a wheatstone's network. The resistance that is to be connected in parallel with the resistance of 10 Ω to balance the network is _____ Ω .

Solution: (10 Ω)

JEE Main 2020 Paper



$$\frac{10R}{10+R} \times 12 = 15 \times 4 \Rightarrow R = 10\ \Omega$$

JEE Main 2020 Paper

Date: 8th January 2020



Time: 09:30 AM – 12:30 PM

Subject: Chemistry

1. The number of bonds between sulphur and oxygen atoms in $\text{S}_2\text{O}_8^{2-}$ and number of bonds between sulphur and sulphur atoms in rhombic sulphur, respectively, are:
- 8 and 6
 - 4 and 6
 - 8 and 8
 - 4 and 8

Answer: c

Solution: Here, we have to count S – O single bonds as well as S = O in $\text{S}_2\text{O}_8^{2-}$, as each double bond also has one sigma bond. The structure of $\text{S}_2\text{O}_8^{2-}$ and S_8 is shown below:



2. The predominant intermolecular forces present in ethyl acetate, a liquid, are:
- London dispersion, dipole-dipole and hydrogen bonding
 - hydrogen bonding and London dispersion
 - dipole-dipole and hydrogen bonding
 - London dispersion and dipole-dipole

Answer: d

Solution: London dispersion forces (also called as induced dipole - induced dipole interactions), exist because of the generation of temporary polarity due to collision of particles and for this very reason, they are present in all molecules and inert gases as well.

Because of the presence of a permanent dipole, there will be dipole-dipole interactions present here.

There is no H that is directly attached to an oxygen atom, so H-bonding cannot be present.

JEE Main 2020 Paper

3. For the Balmer series in the spectrum of H-atom,

$$\bar{\nu} = R_H \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

The correct statements among (A) to (D) are:

- A) The integer $n_1 = 2$.
 - B) The ionization energy of hydrogen can be calculated from the wave number of these lines.
 - C) The lines of longest wavelength corresponds to $n_2 = 3$.
 - D) As wavelength decreases, the lines of the series converge.
- a. B, C, D
 - b. A, B, D
 - c. A, C, D
 - d. A, B, C

Answer: c

Solution:

(A) is correct since the series studied in H-spectrum, including Balmer series, are de-excitation series or emission series. So, electrons get de-excited to $n = 2$ which means that $n_{\text{lower}} = 2$.

(B) It is possible to obtain I.E. from the formula above, but since the question has stated the formula for the Balmer series, n_{lower} has been fixed as 2. So, it is not possible to calculate I.E. from it. To calculate I.E., we'll have to put $n_{\text{lower}} = 1$, which isn't possible here.

(C) $\Delta E = hc/\lambda$

With n_{lower} fixed as 2, ΔE increases as n_{higher} is increased. So, the last line of the Balmer series, i.e. from infinity to $n = 2$, will have the maximum energy in the series and thus, the lowest wavelength. Similarly, the first line in the series, i.e. from $n = 3$ to $n = 2$ will have the lowest energy in the series and thus, the highest wavelength. Which makes this statement correct.

(D) As orbits with higher orbit number or those that are further away from the nucleus are considered, the energy gap in-between subsequent orbits decreases. Now, consider the following for example and with n_{lower} fixed as 2.

Energy of a photon released on transition from $n = 100$ to $n = 2$ will have similar energy to that of the photon that gets released on transition from $n = 101$ to $n = 2$, because energy of the 100th and the 101th orbit will be very close in value. That means they will also have very close values of wavelengths, which further implies that these two lines will be situated quite close to each other on the photographic plate.

In a similar fashion, we can see that as the n_{higher} increases, the lines start to converge together. And since, increasing the n_{higher} will indeed lead to an increase in the energy of the photon released, it will end up releasing photons of shorter wavelengths. Combining these two statements we can easily see that as the wavelength decreases, the spectral lines start to converge.

JEE Main 2020 Paper

4. The first ionization energy (in kJ/mol) of Na, Mg, Al and Si, respectively, are:
- 496, 737, 577, 786
 - 496, 577, 737, 786
 - 496, 577, 786, 737
 - 786, 737, 577, 496


Answer: a

Solution:

The expected order is $\text{Na} < \text{Mg} < \text{Al} < \text{Si}$.

But the actual/experimental order turns out to be $\text{Na} < \text{Al} < \text{Mg} < \text{Si}$, because of the fully filled s-subshell of magnesium and the s^2p^1 configuration of Al which makes it relatively easy for Al to lose its outermost electron.


5. The stoichiometry and solubility product of a salt with the solubility curve given below is, respectively:




- X_2Y , $2 \times 10^{-9}\text{M}^3$
- XY_2 , $1 \times 10^{-9}\text{M}^3$
- XY_2 , $4 \times 10^{-9}\text{M}^3$
- XY , $2 \times 10^{-6}\text{M}^3$

Answer: c


Solution:



$$K_{\text{sp}} = [\text{X}^+]^2[\text{Y}^{2-}] = 4 \times 10^{-6} \times 10^{-3} = 4 \times 10^{-9}$$




$$K_{\text{sp}} = [\text{X}^{2+}][\text{Y}^-]^2 = 10^{-3} \times 4 \times 10^{-6} = 4 \times 10^{-9}$$



JEE Main 2020 Paper

$$K_{sp} = [X^{2+}][Y^{-}]^2 = 10^{-3} \times (2 \times 10^{-3})^2 = 4 \times 10^{-9}$$



$$K_{sp} = [X^{+}][Y^{-}] = 10^{-3} \times 10^{-3} = 10^{-6}$$


6. The complex that can show fac- and mer-isomers is:

- a. $[\text{Co}(\text{NO}_2)_3(\text{NH}_3)_3]$
- b. $[\text{PtCl}_2(\text{NH}_3)_2]$
- c. $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^{+}$
- d. $[\text{CoCl}_2(\text{en})_2]$

Answer: a

Solution: Facial and meridional geometrical isomerism is observed only in $[\text{MA}_3\text{B}_3]$ type complexes which is given in option a.

7. A graph of vapour pressure and temperature for three different liquids X, Y and Z is shown below:



The following inferences are made:

- A) X has higher intermolecular interactions compared to Y
- B) X has lower intermolecular interactions compared to Y
- C) Z has lower intermolecular interactions compared to Y

The correct inference(s) is/are:


- a. C
- b. A
- c. B
- d. A and C

Answer: c

JEE Main 2020 Paper

Solution:

As shown in the plot below, for the same T, the vapour pressure of X is the highest and of Z is the lowest. Now, that means with the same average K.E. of X, Y and Z molecules, the X molecules are able to compensate their respective intermolecular forces better. So, X molecules have the highest vapour pressure. Which implies that the intermolecular forces in X are the weakest among the three. The opposite could be said for Z as well.



8. As per Hardy-Schulze formulation, the flocculation values of the following for ferric hydroxide sol are in the order:
- $\text{AlCl}_3 > \text{K}_3[\text{Fe}(\text{CN})_6] > \text{K}_2\text{CrO}_4 > \text{KBr} = \text{KNO}_3$
 - $\text{K}_3[\text{Fe}(\text{CN})_6] < \text{K}_2\text{CrO}_4 < \text{AlCl}_3 < \text{KBr} < \text{KNO}_3$
 - $\text{K}_3[\text{Fe}(\text{CN})_6] < \text{K}_2\text{CrO}_4 < \text{KBr} = \text{KNO}_3 = \text{AlCl}_3$
 - $\text{K}_3[\text{Fe}(\text{CN})_6] > \text{AlCl}_3 > \text{K}_2\text{CrO}_4 > \text{KBr} > \text{KNO}_3$

Answer: c

Solution:

The minimum concentration of an electrolyte which is required to cause the coagulation or flocculation of a sol is known as flocculation value.

Flocculation value is inversely proportional to coagulation power (coagulation power is directly proportional to the valency of the ions causing coagulation). $\text{Fe}(\text{OH})_3$ sol is a positive sol and thus we have to consider the valency of the anions in the given electrolytes.

9. The rate of a certain biochemical reaction at physiological temperature (T) occurs 10^6 times faster with enzyme than without. The change in activation energy upon adding enzyme is:
- $-6RT$
 - $-6 \times 2.303 RT$
 - $+6RT$
 - $+6 \times 2.303 RT$

Answer: b

JEE Main 2020 Paper

Solution:

$$K_1 = Ae^{-E_{a1}/RT} \text{----(1)}$$

$$K_2 = Ae^{-E_{a2}/RT} \text{----(2)}$$

Dividing equation 1 with equation 2, we get

$$\frac{K_1}{K_2} = e^{(E_{a2}-E_{a1})/RT}$$

$$10^{-6} = e^{(E_{a2}-E_{a1})/RT}$$


Taking \log_e on both sides, we get

$$\Delta E = E_{a2} - E_{a1} = -6 \times 2.303 RT$$

10. When gypsum is heated to 393K, it forms:

- a. $\text{CaSO}_4 \cdot \frac{1}{2} \text{H}_2\text{O}$
- b. Dead burnt plaster
- c. $\text{CaSO}_4 \cdot 5\text{H}_2\text{O}$
- d. Anhydrous CaSO_4

Answer: a



11. The third ionization enthalpy is minimum for:

- a. Mn
- b. Co
- c. Ni
- d. Fe

Answer: d

Solution:

Consider an element E

$E^{2+} \rightarrow E^{3+}$ would be the 3rd I.E. of the element E.

Electronic configuration of Mn is $[\text{Ar}]4s^23d^5$, Co is $[\text{Ar}]4s^23d^7$, Fe is $[\text{Ar}]4s^23d^6$, Ni is $[\text{Ar}]4s^23d^8$

Electronic configuration of Mn^{2+} is $[\text{Ar}]3d^5$, Co^{2+} is $[\text{Ar}]3d^7$, Fe^{2+} is $[\text{Ar}]3d^6$, Ni^{2+} is $[\text{Ar}]3d^8$

As it is evident from the above configurations of the E^{2+} for the given elements, Fe^{2+} would require the least amount of energy for removal of electron as it has the configuration $3d^6 4s^0$. That means that its E^{3+} form is the most stable among the four elements provided in their respective E^{3+} states, i.e., when compared, the next electron removal will require least amount of energy.



JEE Main 2020 Paper

12. The strength of an aqueous NaOH solution is most accurately determined by titrating:
(Note: consider that an appropriate indicator is used)
- Aq. NaOH in a pipette and aqueous oxalic acid in a burette
 - Aq. NaOH in a volumetric flask and concentrated H_2SO_4 in a conical flask
 - Aq. NaOH in a burette and concentrated H_2SO_4 in a conical flask
 - Aq. NaOH in a burette and aqueous oxalic acid in a conical flask

Answer: d

Solution: The standard solution is usually kept in burette. The oxalic acid is a primary standard solution while H_2SO_4 is a secondary standard solution.

13. The decreasing order of reactivity towards dehydrohalogenation (E_1) reaction of the following compounds is:



- $B > A > D > C$
- $B > D > C > A$
- $B > D > A > C$
- $D > B > C > A$

Answer: d

Solution: In E_1 mechanism, the rate determining step is formation of carbocation. So, stability of carbocation formed decides the rate.

In option D, the cation formed is resonance stabilised.






In option C, the cation formed is a 2° carbocation.

In option A and B, the carbocations formed are 1° but there is a chance of rearrangement in option b and after the rearrangement, the carbocation formed in option b will be allylic. So, the order of reaction is as follows:

$D > B > C > A$.


JEE Main 2020 Paper

14. Major product in the following reaction is:



Answer: c

Solution:



15. Arrange the following compounds in increasing order of C—OH bond length: methanol, phenol, p-ethoxyphenol

- Phenol < methanol < p-ethoxyphenol
- methanol < p-ethoxyphenol < phenol
- Phenol < p-ethoxyphenol < methanol
- methanol < phenol < p-ethoxyphenol

Answer: c

JEE Main 2020 Paper

Solution: In methanol, there is no resonance. In phenol, there is resonance. In p-Ethoxyphenol, there is resonance involved but the involvement of lone pair of oxygen in OH group is poor as compared with phenol due to the presence of lone pair oxygen in OCH_3 group which are also involved in resonance.

So, partial double bond character develops in C—OH bond of phenol and p-ethoxyphenol but in case of p-ethoxyphenol, resonance is poor as compared to phenol. So, bond length follows the order: methanol > p-ethoxyphenol > phenol

16. Among the gases (i) – (v), the gases that cause greenhouse effect are:



- i. CO_2
 - ii. H_2O
 - iii. CFC
 - iv. O_2
 - v. O_3
- a. i, ii iii and iv
 - b. i, iii, iv and v
 - c. i and iv
 - d. i, ii, iii and v

Answer: d


Solution: CO_2 , O_3 , H_2O vapours and $\text{CFC}'\text{s}$ are green house gases.

JEE Main 2020 Paper


17. The major products A and B in the following reactions are:




a.




b.



c.



d.




Answer: a


JEE Main 2020 Paper

Solution:

First reaction will generate a free radical which is stabilized by cyanide group.



The alkene undergo homolytic cleavage at the double bond and the terminal radical couple with [A] (secondary radical being more stable will not couple with [A]) forming product B (the secondary radical combines with hydrogen radical generated in the first step).



18. A flask contains a mixture of isohexane and 3-methylpentane. One of the liquids boils at 63 °C while the other boils at 60 °C. What is the best way to separate the two liquids and which one will be distilled out first?
- Fractional distillation, isohexane
 - Simple distillation, 3-methylpentane
 - Fractional distillation, 3-methylpentane
 - Simple distillation, isohexane

Answer: a

Solution: When the difference between the B.P. of the two liquids is less than around 40 °C, fractional distillation is more efficient. The difference between the boiling points of isohexane and 3-methylpentane is only 3 degrees. So, fractional distillation is the best suitable method. Since, isohexane has a lower boiling point, it comes out first.

19. Which of the given statement is not true for glucose?
- The pentacetate glucose does not react with hydroxylamine to give oxime.
 - Glucose reacts with hydroxylamine to form oxime.
 - Glucose gives Schiff's test for aldehyde.
 - Glucose exists in two crystalline forms α and β .

Answer: c


JEE Main 2020 Paper

Solution: Glucose exists in two crystalline forms α and β which are anomers of each other.

Glucose does not react with Schiff's reagent because after the internal cyclisation, it forms either α -anomer or β -anomer. In these forms, free aldehydic group is not present.

Glucose forms open chain structure in aqueous solution which contains aldehyde at chain end. This aldehydic group reacts with NH_4OH to form oxime. On the other hand, glucose penta acetate being a cyclic structure even in aqueous form does not have terminal carbonyl group. Therefore it will not react with NH_4OH .

20. The reagent used for the given conversion is:



- B_2H_6
- LiAlH_4
- NaBH_4
- H_2, Pd

Answer: a

Solution: B_2H_6 does not reduce amide, carbonyl group and cyanide. It selectively reduces carboxylic acid to alcohol. So, for this conversion, it is the best suitable reagent.

21. The volume (in mL) of 0.125 M AgNO_3 required to quantitatively precipitate chloride ions in 0.3 g of $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ is ____.

$$M_{[\text{Co}(\text{NH}_3)_6]\text{Cl}_3} = 267.46 \text{ g/mol}$$

$$M_{\text{AgNO}_3} = 169.87 \text{ g/mol}$$

Answer: 26.92

Solution: To react completely with one mole of $[\text{ML}_6]\text{Cl}_3$, 3 moles of AgNO_3 is required.

$$0.3 \text{ g } [\text{ML}_6]\text{Cl}_3 \text{ means } \frac{0.3}{267.46} \text{ moles of } [\text{ML}_6]\text{Cl}_3.$$

$$\text{So, moles of } \text{AgNO}_3 \text{ required will be } \frac{0.3 \times 3}{267.46} \text{ moles}$$

To find the volume,

$$\frac{0.3 \times 3}{267.46} = 0.125 \times V(L)$$

$$V(L) = 0.02692$$

$$V(\text{mL}) = 26.92$$


JEE Main 2020 Paper

22. What will be the electrode potential for the given half cell reaction at pH= 5?
 $2H_2O \rightarrow O_2 + 4H^+ + 4e^-$; $E^\circ = -1.23 V$

($R=8.314 \text{ J mol}^{-1}\text{K}^{-1}$; temp.=298 K; oxygen under std. atm. Pressure of 1bar.)

Answer: 1.52

Solution:



$$E = +1.23 - \frac{0.0591}{4} \log [H^+]^4$$

$$= +1.23 + (0.0591 \times \text{pH}) = +1.23 + 0.0591 \times 5$$

$$= +1.23 + 0.2955 = +1.5255 V$$


23. Ferrous sulphate heptahydrate is used to fortify foods with iron. The amount (in grams) of the salt required to achieve 10 ppm of iron in 100 kg of wheat is _____.
 Atomic weight: Fe=55.85; S=32.00; O=16.00)

Answer: 4.96

Solution: 10 ppm of Fe means 10 g of Fe in 10^6 g of wheat. So, for 100 kg i.e., 10^5 g

of wheat. Fe needed is 1 g. So, for 1 g of Fe, the mass of $FeSO_4 \cdot 7H_2O$ required is $\frac{278}{56} = 4.96$ g.

24. The magnitude of work done by gas that undergoes a reversible expansion along the path ABC shown in figure is _____.



Answer: 48

Solution:

Work done by the gas

= The area under the curve

= (Area of the square) + (Area of the triangle)


= 48 J

JEE Main 2020 Paper

25. The number of chiral centres in Penicillin is ____.

Answer: 3

Solution: The structure of penicillin is shown below:



So, the number of chiral centers= 3

JEE Main 2020 Paper

Date of Exam: 8th January 2020 (Shift 1)

Time: 9:30 A.M. to 12:30 P.M.

Subject: Mathematics

1. For which of the following ordered pairs (μ, δ) , the system of linear equations

$$x + 2y + 3z = 1$$

$$3x + 4y + 5z = \mu$$

$$4x + 4y + 4z = \delta$$

is inconsistent?

a. $(4, 6)$

b. $(3, 4)$

c. $(1, 0)$

d. $(4, 3)$

Answer: (d)

Solution:

$$D = \begin{vmatrix} 3 & 4 & 5 \\ 1 & 2 & 3 \\ 4 & 4 & 4 \end{vmatrix}$$

$$R_3 \rightarrow R_3 - 2R_1 + 2R_2$$

$$D = \begin{vmatrix} 3 & 4 & 5 \\ 1 & 2 & 3 \\ 0 & 0 & 0 \end{vmatrix} = 0$$

For inconsistent system, one of D_x , D_y , D_z should not be equal to 0

$$D_x = \begin{vmatrix} \mu & 4 & 5 \\ 1 & 2 & 3 \\ \delta & 4 & 4 \end{vmatrix} \quad D_y = \begin{vmatrix} 3 & \mu & 5 \\ 1 & 1 & 3 \\ 4 & \delta & 4 \end{vmatrix}$$

$$D_z = \begin{vmatrix} 3 & 4 & \mu \\ 1 & 2 & 1 \\ 4 & 4 & \delta \end{vmatrix}$$

For inconsistent system, $2\mu \neq \delta + 2$

\therefore The system will be inconsistent for $\mu = 4$, $\delta = 3$.

2. Let $y = y(x)$ be a solution of the differential equation, $\sqrt{1-x^2} \frac{dy}{dx} + \sqrt{1-y^2} = 0$, $|x| < 1$. If

$y\left(\frac{1}{2}\right) = \frac{\sqrt{3}}{2}$, then $y\left(\frac{-1}{\sqrt{2}}\right)$ is equal to :

a. $-\frac{1}{\sqrt{2}}$

b. $-\frac{\sqrt{3}}{2}$

c. $\frac{1}{\sqrt{2}}$

d. $\frac{\sqrt{3}}{2}$

Answer: (c)

JEE Main 2020 Paper

Solution:

$$\sqrt{1-x^2} \frac{dy}{dx} + \sqrt{1-y^2} = 0$$

$$\Rightarrow \frac{dy}{\sqrt{1-y^2}} + \frac{dx}{\sqrt{1-x^2}} = 0$$

$$\Rightarrow \sin^{-1} y + \sin^{-1} x = c$$

$$\text{If } x = \frac{1}{2}, y = \frac{\sqrt{3}}{2} \text{ then,}$$

$$\sin^{-1} \frac{\sqrt{3}}{2} + \sin^{-1} \frac{1}{2} = c$$

$$\therefore \frac{\pi}{3} + \frac{\pi}{6} = c \Rightarrow c = \frac{\pi}{2}$$

$$\Rightarrow \sin^{-1} y = \frac{\pi}{2} - \sin^{-1} x = \cos^{-1} x$$

$$\therefore \sin^{-1} y = \cos^{-1} \frac{1}{\sqrt{2}}$$

$$\Rightarrow \sin^{-1} y = \frac{\pi}{4}$$

$$\Rightarrow y\left(\frac{1}{\sqrt{2}}\right) = \frac{1}{\sqrt{2}}$$

3. If a, b and c are the greatest values of ${}^{19}C_p, {}^{20}C_q, {}^{21}C_r$ respectively, then :

a. $\frac{a}{11} = \frac{b}{22} = \frac{c}{42}$

b. $\frac{a}{10} = \frac{b}{11} = \frac{c}{42}$

c. $\frac{a}{11} = \frac{b}{22} = \frac{c}{21}$

d. $\frac{a}{10} = \frac{b}{11} = \frac{c}{21}$

Answer: (a)

Solution:

We know that, nC_r is maximum when $r = \begin{cases} \frac{n}{2}, & n \text{ is even} \\ \frac{n+1}{2} \text{ or } \frac{n-1}{2}, & n \text{ is odd} \end{cases}$

$$\text{Therefore, } \max({}^{19}C_p) = {}^{19}C_9 = a$$

$$\max({}^{20}C_q) = {}^{20}C_{10} = b$$

$$\max({}^{21}C_r) = {}^{21}C_{11} = c$$

JEE Main 2020 Paper

$$\therefore \frac{a}{{}^{19}C_9} = \frac{b}{\frac{20}{10} \times {}^{19}C_9} = \frac{c}{\frac{21}{11} \times \frac{20}{10} \times {}^{19}C_9}$$

$$\Rightarrow \frac{a}{1} = \frac{b}{2} = \frac{c}{\frac{42}{11}}$$

$$\Rightarrow \frac{a}{11} = \frac{b}{22} = \frac{c}{42}$$

4. Which of the following is a tautology?

a. $(P \wedge (P \rightarrow Q)) \rightarrow Q$

b. $P \wedge (P \vee Q)$

c. $(Q \rightarrow (P \wedge (P \rightarrow Q)))$

d. $P \vee (P \wedge Q)$

Answer: (a)

Solution:

$$(P \wedge (P \rightarrow Q)) \rightarrow Q$$

$$= (P \wedge (\sim P \vee Q)) \rightarrow Q$$

$$= [(P \wedge \sim P) \vee (P \wedge Q)] \rightarrow Q$$

$$= P \wedge Q \rightarrow Q$$

$$= \sim (P \wedge Q) \vee Q$$

$$= \sim P \vee \sim Q \vee Q$$

$$= T$$

5. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be such that for all $x \in \mathbf{R}$, $(2^{1+x} + 2^{1-x}), f(x)$ and $(3^x + 3^{-x})$ are in A.P., then the minimum value of $f(x)$ is :

a. 0

b. 4

c. 3

d. 2

Answer: (c)

Solution:

$$2^{1-x} + 2^{1+x}, f(x), 3^x + 3^{-x} \text{ are in A.P.}$$

$$\therefore f(x) = \frac{3^x + 3^{-x} + 2^{1+x} + 2^{1-x}}{2} = \frac{(3^x + 3^{-x})}{2} + \frac{2^{1+x} + 2^{1-x}}{2}$$

Applying A.M. \geq G.M. inequality, we get

$$\frac{(3^x + 3^{-x})}{2} \geq \sqrt{3^x \cdot 3^{-x}}$$

09/09/2020 (Shift 1) - Mathematics

Page 14

Downloaded from @JEEPrep

$$\Rightarrow \frac{2^{1+x} + 2^{1-x}}{2} \geq 2 \quad \dots (2)$$

$$f(x) \geq 1 + 2 = 3$$

6. The locus of a point which divides the line segment joining the point $(0, -1)$ and a point on the parabola, $x^2 = 4y$, internally in the ratio $1:2$, is :

b. $4x^2 - 3y = 2$

d. $9x^2 - 3y = 2$

Solution:

$$h = \frac{2t}{3}, k = \frac{-2 + t^2}{3}$$
$$3k + 2 = \left(\frac{3h}{2}\right)^2$$

7. For $a > 0$, let the curves $C_1: y^2 = ax$ and $C_2: x^2 = ay$ intersect at origin O and a point P . Let the line $x = b$ ($0 < b < a$) intersect the chord OP and the x -axis at points Q and R , respectively. If the line $x = b$ bisects the area bounded by the curves, C_1 and C_2 , and the area of $\Delta OQR = \frac{1}{2}$, then ' a ' satisfies the equation :

b. $x^6 - 12x^3 - 4 = 0$

d. $x^6 - 6x^3 + 4 = 0$


Solution:

JEE Main 2020 Paper

Given, $ar(\Delta APQ) = \frac{1}{2}$

$$\Rightarrow \frac{1}{2} \times b \times b = \frac{1}{2}$$

$$\Rightarrow b = 1$$



As per the question

$$\Rightarrow \int_0^1 \left(\sqrt{ax} - \frac{x^2}{a} \right) dx = \frac{1}{2} \int_0^a \left(\sqrt{ax} - \frac{x^2}{a} \right) dx$$

$$\Rightarrow \frac{2}{3} \sqrt{a} - \frac{1}{3a} = \frac{a^2}{6}$$

$$\Rightarrow 2a\sqrt{a} - 1 = \frac{a^3}{2}$$

$$\Rightarrow 4a\sqrt{a} = 2 + a^3$$

$$\Rightarrow 16a^3 = 4 + a^6 + 4a^3$$

$$\Rightarrow a^6 - 12a^3 + 4 = 0.$$

8. The inverse function of $f(x) = \frac{8^{2x} - 8^{-2x}}{8^{2x} + 8^{-2x}}$, $x \in (-1, 1)$, is

a. $\frac{1}{4} (\log_8 e) \log_e \left(\frac{1-x}{1+x} \right)$

b. $\frac{1}{4} (\log_8 e) \log_e \left(\frac{1+x}{1-x} \right)$

c. $\frac{1}{4} \log_e \left(\frac{1+x}{1-x} \right)$

d. $\frac{1}{4} \log_e \left(\frac{1-x}{1+x} \right)$

Answer: (b)

Solution:

$$f(x) = \frac{8^{2x} - 8^{-2x}}{8^{2x} + 8^{-2x}} = \frac{8^{4x} - 1}{8^{4x} + 1}$$

JEE Main 2020 Paper

$$\text{Put } y = \frac{8^{4x}-1}{8^{4x}+1}$$

Applying componendo-dividendo on both sides

$$\frac{y+1}{y-1} = \frac{2 \times 8^{4x}}{-2}$$

$$\frac{y+1}{y-1} = -8^{4x} \Rightarrow 8^{4x} = \frac{1+y}{1-y}$$

$$\Rightarrow x = \frac{1}{4} \log_8 \left(\frac{1+y}{1-y} \right)$$

$$f^{-1}(x) = \frac{1}{4} \log_8 \left(\frac{1+x}{1-x} \right) = \frac{1}{4} \log_8 e \left(\log_e \frac{1+x}{1-x} \right)$$

9. $\lim_{x \rightarrow 0} \left(\frac{3x^2+2}{7x^2+2} \right)^{\frac{1}{x^2}}$ is equal to :

a. e

b. $\frac{1}{e^2}$

c. $\frac{1}{e}$

d. e^2

Answer: (b)

Solution:

$$\text{Let } L = \lim_{x \rightarrow 0} \left(\frac{3x^2+2}{7x^2+2} \right)^{\frac{1}{x^2}} \quad [\text{Intermediate form } 1^\infty]$$

$$\therefore L = e^{\lim_{x \rightarrow 0} \frac{1}{x^2} \left(\frac{3x^2+2}{7x^2+2} - 1 \right)}$$

$$= e^{\lim_{x \rightarrow 0} \frac{1}{x^2} \left(-\frac{4x^2}{7x^2+2} \right)}$$

$$= \frac{1}{e^2}$$

10. Let $f(x) = (\sin(\tan^{-1} x) + \sin(\cot^{-1} x))^2 - 1$, where $|x| > 1$.

If $\frac{dy}{dx} = \frac{1}{2} \frac{d}{dx} (\sin^{-1}(f(x)))$ and $y(\sqrt{3}) = \frac{\pi}{6}$, then $y(-\sqrt{3})$ is equal to :

a. $\frac{\pi}{3}$

b. $\frac{2\pi}{3}$

c. $-\frac{\pi}{6}$

d. $\frac{5\pi}{6}$

JEE Main 2020 Paper

Answer: (Bonus)

Solution:

$$f(x) = [\sin(\tan^{-1} x) + \sin(\cot^{-1} x)]^2 - 1$$

$$\text{Put } \tan^{-1} x = \phi, \text{ where } \phi \in \left(-\frac{\pi}{2}, -\frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$$

$$= [\sin(\tan^{-1} x) + \sin(\cot^{-1} x)]^2 - 1 = [\sin \phi + \cos \phi]^2 - 1$$

$$= 1 + 2 \sin \phi \cos \phi - 1 = \sin 2\phi = \frac{2x}{1+x^2}$$

$$\text{It is given that } \frac{dy}{dx} = \frac{1}{2} \frac{d}{dx} (\sin^{-1}(f(x)))$$

$$\Rightarrow \frac{dy}{dx} = -\frac{1}{1+x^2}, \text{ for } |x| > 1$$

$$|x| > 1 \Rightarrow x > 1 \text{ and } x < -1$$

To get the value of $y(-\sqrt{3})$, we have to integrate the value of $\frac{dy}{dx}$. To integrate the expression, the interval should be continuous. Therefore, we have to integrate the expression in both the intervals.

$$\Rightarrow y = -\tan^{-1} x + C_1, \text{ for } x > 1 \text{ and } y = -\tan^{-1} x + C_2, \text{ for } x < -1$$

$$\text{For } x > 1, C_1 = \frac{\pi}{2} \because y(\sqrt{3}) = \frac{\pi}{6} \text{ is given.}$$

But C_2 can't be determined as no other information is given for $x < -1$. Therefore, all the options can be true as C_2 can't be determined.

11. If the equation, $x^2 + bx + 45 = 0$ ($b \in \mathbf{R}$) has conjugate complex roots and they satisfy $|z + 1| = 2\sqrt{10}$, then :

a. $b^2 + b = 12$

b. $b^2 - b = 42$

c. $b^2 - b = 30$

d. $b^2 + b = 72$

Answer: (c)

Solution:

$$\text{Given } x^2 + bx + 45 = 0, b \in \mathbf{R}, \text{ let roots of the equation be } p \pm iq$$

$$\text{Then, sum of roots} = 2p = -b$$

$$\text{Product of roots} = p^2 + q^2 = 45$$

$$\text{As } p \pm iq \text{ lies on } |z + 1| = 2\sqrt{10}, \text{ we get}$$

$$(p + 1)^2 + q^2 = 40$$

JEE Main 2020 Paper

$$\Rightarrow p^2 + q^2 + 2p + 1 = 40$$

$$\Rightarrow 45 - b + 1 = 40$$

$$\Rightarrow b = 6$$

$$\Rightarrow b^2 - b = 30.$$

12. The mean and standard deviation (s.d.) of 10 observations are 20 and 2 respectively. Each of these 10 observations is multiplied by p and then reduced by q , where $p \neq 0$ and $q \neq 0$. If the new mean and standard deviation become half of their original values, then q is equal to:

- [illegible]

Answer: (a)

Solution:

If mean \bar{x} is multiplied by p and then q is subtracted from it,

then new mean $\bar{x}' = p\bar{x} - q$

$$\therefore \bar{x}' = \frac{1}{2}\bar{x} \text{ and } \bar{x} = 10$$

$$\Rightarrow 10 = 20p - q \quad \dots (1)$$

If standard deviation is multiplied by p , new standard deviation (σ') is $|p|$ times of the initial standard deviation (σ).

$$\sigma' = |p|\sigma$$

$$\Rightarrow \frac{1}{2}\sigma = |p|\sigma \Rightarrow |p| = \frac{1}{2}$$

If $p = \frac{1}{2}$, $q = 0$

If $p = -\frac{1}{2}$, $q = -20$.

13. If $\int \frac{\cos x}{\sin^3 x (1 + \sin^6 x)^{\frac{2}{3}}} dx = f(x)(1 + \sin^6 x)^{\frac{1}{\lambda}} + c$, where c is a constant of integration, then $\lambda f\left(\frac{\pi}{3}\right)$ is equal to :

- a. $-\frac{9}{8}$ b. $\frac{9}{8}$
c. 2 d. -2

Answer: (d)

Solution:

Let $\sin x = t \Rightarrow \cos x \, dx = dt$

JEE Main 2020 Paper

$$\therefore \int \frac{dt}{t^3 (1+t^6)^{\frac{2}{3}}} = \int \frac{dt}{t^7 \left(1+\frac{1}{t^6}\right)^{\frac{2}{3}}}$$

$$\text{Let } 1 + \frac{1}{t^6} = u \Rightarrow -6t^{-7} dt = du$$

$$\Rightarrow \int \frac{dt}{t^7 \left(1+\frac{1}{t^6}\right)^{\frac{2}{3}}} = -\frac{1}{6} \int \frac{du}{u^{\frac{2}{3}}} = -\frac{3}{6} u^{\frac{1}{3}} + c = -\frac{1}{2} \left(1 + \frac{1}{t^6}\right)^{\frac{1}{3}} + c$$

$$= -\frac{(1+\sin^6 x)^{\frac{1}{3}}}{2 \sin^2 x} + c = f(x)(1 + \sin^6 x)^{\frac{1}{\lambda}}$$

$$\therefore \lambda = 3 \text{ and } f(x) = -\frac{1}{2 \sin^2 x}$$

$$\Rightarrow \lambda f\left(\frac{\pi}{3}\right) = -2.$$

14. Let A and B be two independent events such that $P(A) = \frac{1}{3}$ and $P(B) = \frac{1}{6}$. Then, which of the following is TRUE ?

a. $P(A/(A \cup B)) = \frac{1}{4}$

b. $P(A/B') = \frac{1}{3}$

c. $P(A/B) = \frac{2}{3}$

d. $P(A'/B') = \frac{1}{3}$

Answer: (b)

Solution:

If X and Y are independent events, then

$$P\left(\frac{X}{Y}\right) = \frac{P(X \cap Y)}{P(Y)} = \frac{P(X)P(Y)}{P(Y)} = P(X)$$

$$\text{Therefore, } P\left(\frac{A}{B}\right) = P(A) = \frac{1}{3} \Rightarrow P\left(\frac{A}{B'}\right) = P(A) = \frac{1}{3}$$

15. If volume of parallelopiped whose coterminous edges are given by $\vec{u} = \hat{i} + \hat{j} + \lambda \hat{k}$,

$\vec{v} = \hat{i} + \hat{j} + 3\hat{k}$ and $\vec{w} = 2\hat{i} + \hat{j} + \hat{k}$ be 1 cu. unit. If θ be the angle between the edges \vec{u} and \vec{w} , then, $\cos \theta$ can be :

a. $\frac{7}{6\sqrt{6}}$

b. $\frac{5}{7}$

c. $\frac{7}{6\sqrt{3}}$

d. $\frac{5}{3\sqrt{3}}$

Answer: (c)

Solution:

$$\text{Volume of parallelepiped} = [\vec{u} \ \vec{v} \ \vec{w}]$$

JEE Main 2020 Paper

$$\Rightarrow \begin{vmatrix} 1 & 1 & \lambda \\ 1 & 1 & 3 \\ 2 & 1 & 1 \end{vmatrix} = \pm 1$$

$$\Rightarrow \lambda = 2 \text{ or } 4$$

For $\lambda = 4$,

$$\cos \theta = \frac{2+1+4}{\sqrt{6}\sqrt{18}} = \frac{7}{6\sqrt{3}}$$

16. Let two points be $A(1, -1)$ and $B(0, 2)$. If a point $P(x', y')$ be such that the area of $\Delta PAB = 5$ sq. units and it lies on the line, $3x + y - 4\lambda = 0$, then the value of λ is :

- a. 4
b. 1
c. -3
d. 3

Answer: (d)

Solution:

Area of triangle is

$$A = \frac{1}{2} \begin{vmatrix} 1 & -1 & 1 \\ 0 & 2 & 1 \\ x' & y' & 1 \end{vmatrix} = \pm 5$$

$$\Rightarrow (2 - y') - x' - 2x' = \pm 10$$

$$\Rightarrow -3x' - y' + 2 = \pm 10$$

$$3x' + y' = 12 \text{ or } 3x' + y' = -8$$

$$\Rightarrow \lambda = 3 \text{ or } -2$$

17. The shortest distance between the lines

$$\frac{x-3}{3} = \frac{y-8}{-1} = \frac{z-3}{1} \text{ and}$$

$$\frac{x+3}{-3} = \frac{y+7}{2} = \frac{z-6}{4} \text{ is :}$$

- a. $2\sqrt{30}$
b. $\frac{7}{2}\sqrt{30}$
c. 3
d. $3\sqrt{30}$

Answer: (d)

Solution:

$$\vec{AB} = -3\hat{i} - 7\hat{j} + 6\hat{k} - (3\hat{i} + 8\hat{j} + 3\hat{k}) = -6\hat{i} - 15\hat{j} + 3\hat{k}$$

$$\vec{p} = 3\hat{i} - \hat{j} + \hat{k}$$

$$\vec{q} = -3\hat{i} + 2\hat{j} + 4\hat{k}$$

JEE Main 2020 Paper

$$\vec{p} \times \vec{q} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 1 \\ -3 & 2 & 4 \end{vmatrix} = -6\hat{i} - 15\hat{j} + 9\hat{k}$$

$$\text{Shortest distance} = \frac{|\vec{AB} \cdot (\vec{p} \times \vec{q})|}{|\vec{p} \times \vec{q}|} = \frac{|36 + 225 + 9|}{\sqrt{36 + 225 + 9}} = 3\sqrt{30}.$$

18. Let the line $y = mx$ and the ellipse $2x^2 + y^2 = 1$ intersect a point P in the first quadrant. If the normal to this ellipse at P meets the co-ordinate axes at $(-\frac{1}{3\sqrt{2}}, 0)$ and $(0, \beta)$, then β is equal to :

a. $\frac{2}{\sqrt{3}}$


b. $\frac{2}{3}$

c. $\frac{2\sqrt{2}}{3}$

d. $\frac{\sqrt{2}}{3}$

Answer: (d)

Solution:



Let $P \equiv (x_1, y_1)$

$2x^2 + y^2 = 1$ is given equation of ellipse.

$$\Rightarrow 4x + 2yy' = 0$$

$$\Rightarrow y'|_{(x_1, y_1)} = -\frac{2x_1}{y_1}$$

Therefore, slope of normal at $P(x_1, y_1)$ is $\frac{y_1}{2x_1}$

Equation of normal at $P(x_1, y_1)$ is

$$(y - y_1) = \frac{y_1}{2x_1}(x - x_1)$$

It passes through $(-\frac{1}{3\sqrt{2}}, 0)$

JEE Main 2020 Paper

$$\Rightarrow -y_1 = \frac{y_1}{2x_1} \left(-\frac{1}{3\sqrt{2}} - x_1 \right)$$

$$\Rightarrow x_1 = \frac{1}{3\sqrt{2}}$$

$$\Rightarrow y_1 = \frac{2\sqrt{2}}{3} \text{ as } P \text{ lies in first quadrant}$$

Since $(0, \beta)$ lies on the normal of the ellipse at point P , hence we get

$$\beta = \frac{y_1}{2} = \frac{\sqrt{2}}{3}$$

19. If c is a point at which Rolle's theorem holds for the function, $f(x) = \log_e \left(\frac{x^2 + \alpha}{7x} \right)$ in the interval $[3, 4]$, where $\alpha \in \mathbf{R}$, then $f''(c)$ is equal to :

a. $-\frac{1}{24}$

b. $\frac{-1}{12}$

c. $\frac{\sqrt{3}}{7}$

d. $\frac{1}{12}$

Answer: (d)

Solution:

Rolle's theorem is applicable on $f(x)$ in $[3, 4]$

$$\Rightarrow f(3) = f(4)$$

$$\Rightarrow \ln \left(\frac{9 + \alpha}{21} \right) = \ln \left(\frac{16 + \alpha}{28} \right)$$

$$\Rightarrow \frac{9 + \alpha}{21} = \frac{16 + \alpha}{28}$$

$$\Rightarrow 36 + 4\alpha = 48 + 3\alpha \Rightarrow \alpha = 12$$

$$\text{Now, } f(x) = \ln \left(\frac{x^2 + 12}{7x} \right) \Rightarrow f'(x) = \frac{7x}{x^2 + 12} \times \frac{7x \times 2x - (x^2 + 12) \times 7}{(7x)^2}$$

$$f'(x) = \frac{x^2 - 12}{x(x^2 + 12)}$$

$$f'(c) = 0 \Rightarrow c = 2\sqrt{3}$$

$$f''(x) = \frac{-x^4 + 48x^2 + 144}{x^2(x^2 + 12)^2}$$

$$\therefore f''(c) = \frac{1}{12}$$

JEE Main 2020 Paper

20. Let $f(x) = x \cos^{-1}(\sin(-|x|))$, $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, then which of the following is true ?

- a. $f'(0) = -\frac{\pi}{2}$
- b. f' is decreasing in $\left(-\frac{\pi}{2}, 0\right)$ and increasing in $\left(0, \frac{\pi}{2}\right)$
- c. f is not differentiable at $x = 0$
- d. f' is increasing in $\left(-\frac{\pi}{2}, 0\right)$ and decreasing in $\left(0, \frac{\pi}{2}\right)$

Answer: (b)

Solution:

$$f(x) = x \cos^{-1}(\sin(-|x|))$$

$$\Rightarrow f(x) = x \cos^{-1}(-\sin |x|)$$

$$\Rightarrow f(x) = x[\pi - \cos^{-1}(\sin |x|)]$$

$$\Rightarrow f(x) = x \left[\pi - \left(\frac{\pi}{2} - \sin^{-1}(\sin |x|) \right) \right]$$

$$\Rightarrow f(x) = x \left(\frac{\pi}{2} + |x| \right)$$

$$\Rightarrow f(x) = \begin{cases} x \left(\frac{\pi}{2} + x \right), & x \geq 0 \\ x \left(\frac{\pi}{2} - x \right), & x < 0 \end{cases}$$

$$\Rightarrow f'(x) = \begin{cases} \left(\frac{\pi}{2} + 2x \right), & x \geq 0 \\ \left(\frac{\pi}{2} - 2x \right), & x < 0 \end{cases}$$

Therefore, $f'(x)$ is decreasing $\left(-\frac{\pi}{2}, 0\right)$ and increasing in $\left(0, \frac{\pi}{2}\right)$.

21. An urn contains 5 red marbles, 4 black marbles and 3 white marbles. Then the number of ways in which 4 marbles can be drawn so that at most three of them are red is _____.

Answer: (490)

Solution:

Number of ways to select at most 3 red balls = $P(0 \text{ red balls}) + P(1 \text{ red ball}) + P(2 \text{ red balls})$
 $+ P(3 \text{ red balls})$

$$= {}^7C_4 + {}^5C_1 \times {}^7C_3 + {}^5C_2 \times {}^7C_2 + {}^5C_3 \times {}^7C_1$$

$$= 35 + 175 + 210 + 70 = 490$$

JEE Main 2020 Paper

22. Let the normal at a P on the curve $y^2 - 3x^2 + y + 10 = 0$ intersect the y -axis at $\left(0, \frac{3}{2}\right)$. If m is the slope of the tangent at P to the curve, then $|m|$ is equal to_____.

Answer: (4)

Solution:


Let co-ordinate of P be (x_1, y_1)

Differentiating the curve w.r.t x

$$2yy' - 6x + y' = 0$$

Slope of tangent at P

$$\Rightarrow y' = \frac{6x_1}{1 + 2y_1}$$



$$\therefore m_{\text{normal}} = \left(\frac{y_1 - \frac{3}{2}}{x_1 - 0} \right)$$

$$\therefore m_{\text{normal}} \times m_{\text{tangent}} = -1$$

$$\Rightarrow \frac{\frac{3}{2} - y_1}{-x_1} \times \frac{6x_1}{1 + 2y_1} = -1$$

$$\Rightarrow y_1 = 1$$

$$\Rightarrow x_1 = \pm 2$$

$$\text{Slope of tangent} = \pm \frac{12}{3} = \pm 4$$

$$\Rightarrow |m| = 4$$

23. The least positive value of ' a ' for which the equation, $2x^2 + (a - 10)x + \frac{33}{2} = 2a$ has real roots is _____.

Answer: (8)

Solution:

JEE Main 2020 Paper

$$\therefore 2x^2 + (a - 10)x + \frac{33}{2} = 2a, a \in \mathbf{Z}^+ \text{ has real roots}$$

$$\Rightarrow D \geq 0 \Rightarrow (a - 10)^2 - 4 \times 2 \times \left(\frac{33}{2} - 2a\right) \geq 0$$

$$\Rightarrow (a - 10)^2 - 4(33 - 4a) \geq 0$$

$$\Rightarrow a^2 - 4a - 32 \geq 0 \Rightarrow a \in (-\infty, -4] \cup [8, \infty)$$

Thus, minimum value of ' a ' $\forall a \in \mathbf{Z}^+$ is 8.

24. The sum $\sum_{k=1}^{20}(1 + 2 + 3 + \dots + k)$ is _____.

Answer: (1540)

Solution:

$$\begin{aligned} &= \sum_{k=1}^{20} \frac{k(k+1)}{2} \\ &= \frac{1}{2} \sum_{k=1}^{20} k^2 + k \\ &= \frac{1}{2} \left[\frac{20(21)(41)}{6} + \frac{20(21)}{2} \right] = \frac{1}{2} [2870 + 210] = 1540 \end{aligned}$$

25. The number of all 3×3 matrices A, with entries from the set $\{-1, 0, 1\}$ such that the sum of the diagonal elements of (AA^T) is 3, is _____.

Answer: (672)

Solution:

$$\text{tr}(AA^T) = 3$$

$$\text{Let } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \text{ and } A^T = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{bmatrix}$$

$$\text{tr}(AA^T) = a_{11}^2 + a_{12}^2 + a_{13}^2 + a_{21}^2 + a_{22}^2 + a_{23}^2 + a_{31}^2 + a_{32}^2 + a_{33}^2 = 3$$

So out of 9 elements (a_{ij}) 's, 3 elements must be equal to 1 or -1 and rest elements must be 0.

So, the total possible cases will be

When there is 6(0's) and 3(1's) then the total possibilities is 9C_6

For 6(0's) and 3(-1 's) total possibilities is 9C_6

For 6(0's), 2(1's) and 1(-1 's) total possibilities is ${}^9C_6 \times 3$

For 6(0's), 1(1's) and 2(-1 's) total possibilities is ${}^9C_6 \times 3$

$$\therefore \text{Total number of cases} = {}^9C_6 \times 8 = 672$$