

Acemiz Plus Tablet

1. Drug Name and Strength

- **Generic Name:** Acemiz Plus
- **Brand Names:** (may vary by manufacturer)
- **Dosage Forms:** Tablets
- **Strengths:** 100 mg

2. Indications

- Acemiz Plus is prescribed for:
 - Pain relief (mild to moderate pain)
 - Inflammatory conditions (e.g., arthritis)
 - Fever reduction

3. Mechanism of Action

- **Aceclofenac:** Non-steroidal anti-inflammatory drug (NSAID) that reduces pain and inflammation by inhibiting cyclooxygenase (COX) enzymes.
- **Paracetamol (Acetaminophen):** Analgesic and antipyretic that works centrally in the brain to reduce pain and fever.

4. Dosage and Administration

- **Adults:**
 - Typical dosage: 1 tablet every 8-12 hours as needed, not exceeding the recommended daily limit.
- **Children:** Dosage should be determined by a healthcare provider based on age and weight.

5. Contraindications

- Known hypersensitivity to any ingredient in Acemiz Plus.
- Severe liver or kidney impairment.
- Active gastrointestinal bleeding or ulceration.

6. Warnings and Precautions

- Use with caution in patients with a history of gastrointestinal disorders.
- Monitor liver and kidney function during prolonged use.
- Pregnant or breastfeeding women should consult their healthcare provider.

7. Side Effects

- **Common:**
 - Nausea, vomiting
 - Stomach pain
 - Headache
 - Dizziness
- **Serious (Rare):**
 - Allergic reactions (anaphylaxis)
 - Liver damage
 - Kidney issues
 - Severe gastrointestinal complications

8. Drug Interactions

- Other NSAIDs: Increased risk of side effects.
- Anticoagulants (e.g., warfarin): May enhance bleeding risk.
- Alcohol: Increased risk of liver toxicity.

- Certain antihypertensives: Reduced efficacy.

9. Storage Conditions

- Store at room temperature, away from light and moisture.
- Keep out of reach of children.

10. Patient Counseling Information

- Take the medication as directed, not exceeding the recommended dosage.
- Inform healthcare provider of any other medications being taken.
- Report any signs of allergic reaction or severe gastrointestinal issues.
- Stay hydrated and avoid alcohol while taking this medication.

11. Overdosage

- Symptoms may include nausea, vomiting, and stomach pain. Management involves supportive care and may include gastric lavage in severe cases.