

1. Chapter 1: Introduction

Overview of number partitioning problem.

Application: randomized control trials.

Other applications.

- Circuit design, etc.

Importance as a basic NP-complete problem.

Two questions of interest:

1. What is optimal solution.
2. How to find optimal solution.

1.1. Physical Interpretations

1.2. Statistical-to-Computational Gap

2. Number Packing Problem

Let N be the dimensionality, and $\Sigma_N := \{\pm 1\}$ be the binary cube. Suppose we're given $g \sim \mathcal{N}(0, I_N)$. We want to find $x \in \Sigma_N$ such that we minimize $|\langle x, g \rangle|$.

Definition 2.1. Let $\delta > 0$. The *solution set* $S(\delta; g)$ is the set of all $x \in \Sigma_N$ that are δ -optimal, satisfying

$$\frac{1}{\sqrt{N}} |\langle g, x \rangle| \leq 2^{-\delta N}. \quad (2.1)$$

(2.1) Hi

2.1. Existing Results

1. $X_i, 1 \leq i \leq n$ i.i.d. uniform from $\{1, 2, \dots, M := 2^m\}$, with $\kappa := \frac{m}{n}$, then phase transition going from $\kappa < 1$ to $\kappa > 1$.
2. Average case, X_i i.i.d. standard Normal.
3. Karmarkar [KKLO86] - NPP value is $\Theta(\sqrt{N}2^{-N})$ whp as $N \rightarrow \infty$ (doesn't need Normality).
4. Best polynomial-time algorithm: Karmarkar-Karp [KK82] - Discrepancy $O(N^{-\alpha \log N}) = 2^{-\Theta(\log^2 N)}$ whp as $N \rightarrow \infty$
5. PDM (paired differencing) heuristic - fails for i.i.d. uniform inputs with objective $\Theta(n^{-1})$ (Lueker).
6. LDM (largest differencing) heuristic - works for i.i.d. Uniforms, with $n^{-\Theta(\log n)}$ (Yakir, with constant $\alpha = \frac{1}{2 \ln 2}$ calculated non-rigorously by Boettcher and Mertens).
7. Krieger - $O(n^{-2})$ for balanced partition.
8. Hoberg [HHRY17] - computational hardness for worst-case discrepancy, as poly-time oracle that can get discrepancy to within $O(2^{\sqrt{n}})$ would be oracle for Minkowski problem.
9. Gamarnik-Kizildag: Information-theoretic guarantee $E_n = n$, best computational guarantee $E_n = \Theta(\log^2 n)$.

10. Existence of m -OGP for $m = O(1)$ and $E_n = \Theta(n)$.
11. Absence for $\omega(1) \leq E_n = o(n)$
12. Existence for $\omega(\sqrt{n \log_2 n}) \leq E_n \leq o(n)$ for $m = \omega_{n(1)}$ (with changing η, β)
 1. While OGP not ruled out for $E_n \leq \omega(\sqrt{n \log_2 n})$, argued that it is tight.
13. For $\varepsilon \in (0, \frac{1}{5})$, no stable algorithm can solve $\omega(n \log^{-\frac{1}{5}+\varepsilon} n) \leq E_n \leq o(n)$
14. Possible to strengthen to $E_n = \Theta(n)$ (as $2^{-\Theta(n)} \leq 2^{-o(n)}$)

3. Glossary and conventions

Conventions:

1. \log means \log in base 2, \exp is \exp base 2 - natural \log /exponent is \ln/e^x .
- 2.

Glossary:

1. “instance”/“disorder” - g , instance of the NPP problem
2. “discrepancy” - for a given g , value of $\min_{x \in \Sigma_N} |\langle g, x \rangle|$
3. “energy” - negative exponent of discrepancy, i.e. if discrepancy is 2^{-E} , then energy is E . Lower energy indicates “worse” discrepancy.
4. “near-ground state”/“approximate solution”

4. Literature Review

[1]

[2]

[3]

[4]

[5]

[6]

[7]

[8]

[9]

[10]

[11]

[12]

[13]

[14]

[15]

[16]

[17]

[18]

[19]

[20]

[21]

[22]

[23]

[24]

[25]

[26]

[27]

[28]

[29]

[30]

[31]

[32]

[33]

[34]

[35]

[36]

[37]

[38]

[39]

[40]

[41]

[42]

[43]

[44]

[45]

[46]

[47]

[48]

[49]

[50]

[51]

[52]

[53]

[54]

[55]

[56]

[57]

[58]

[59]

[60]

[61]

[62]

[63]

[64]

[65]

[66]

[67]

[68]

[69]

[70]

[71]

[72]

[73]

[74]

[75]

[76]

[77]

[78]

[79]

[80]

[81]

[82]

[83]

[84]

[85]

[86]

[87]

[88]

5. Low-Degree Algorithms

What are algorithms of interest

For our purposes, an *algorithm* is a function which takes as input a problem instance $g \sim \mathcal{N}(0, I_N)$ and outputs some $x \in \Sigma_N$. This definition can be extended to functions giving outputs on \mathbf{R}^N (and rounding to a vertex on the hypercube Σ_N), or to taking as additional input some randomness ω , allowing for randomized algorithms. However, most of our analysis will focus on the deterministic case.

To further restrict the category of algorithms considered, we specifically restrict to

Why study low-degree algorithms (poly time heuristic + simple)

Different notions of degree.

Overview of section

- Efron-Stein notion
- Hermite notion
- Algorithms and Stability Bounds

5.1. Efron-Stein Polynomials (TODO)

Definition 5.1. Let π be a probability distribution on \mathbf{R} . The L^2 space $L^2(\mathbf{R}^N, \pi^{\otimes N})$ is the space of functions $f : \mathbf{R}^N \rightarrow \mathbf{R}$ with finite L^2 norm.

$$\mathbf{E}[f^2] := \int_{x=(x_1, \dots, x_n) \in \mathbf{R}^N} f(x)^2 d\pi^{\otimes N}(x) < \infty.$$

Alternatively, this is the space of L^2 functions of N i.i.d. random variables x_i , distributed as π .

Motivation for studying decompositions of functions by projecting onto coordinates.

This section largely follows [76, § 8.3].

Want to decompose

$$f = \sum_{S \subseteq [n]} f^{\neg S} \quad (5.1)$$

Want $f^{\neg S}$ to only depend on the coordinates in S .

If $T \not\subseteq S$ and g depends only on the coordinates in T , then $\langle f^{\neg S}, g \rangle = 0$.

Definition 5.2. Let $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ and $J \subseteq [n]$, with $\bar{J} = [n] \setminus J$. The *projection of f onto J* is the function $f^{\subseteq J} : \mathbf{R}^N \rightarrow \mathbf{R}$ given by

$$f^{\subseteq J}(x) = \mathbf{E}[f(x_1, \dots, x_n) \mid x_i, i \in J].$$

This is f with the \bar{J} coordinates re-randomized, so $f^{\subseteq J}$ only depends on x_J .

In particular, we should have that

$$f^{\subseteq J} = \sum_{S \subseteq J} f^{\neg S} \quad (5.2)$$

First, we consider the case $J = \emptyset$. It is clear that $f^{\neg \emptyset} = f^{\subseteq \emptyset}$, which is the constant function $\mathbf{E}[f]$.

Next, if $J = \{j\}$ is a singleton, (5.2) gives

$$f^{\subseteq \{j\}} = f^{\neg \emptyset} + f^{\neg \{j\}},$$

and as $f^{\subseteq \{j\}}(x) = \mathbf{E}[f \mid x_j]$, we get

$$f^{\neg \{j\}} = \mathbf{E}[f \mid x_j] - \mathbf{E}[f].$$

This function only depends on x_j ; all other coordinates are averaged over. It measures what difference in expectation of f is given x_j .

Continuing on to sets of two coordinates, some brief manipulation gives, for $J = \{i, j\}$,

$$\begin{aligned} f^{\subseteq \{i, j\}} &= f^{\neg \emptyset} + f^{\neg \{i\}} + f^{\neg \{j\}} + f^{\neg \{i, j\}} \\ &= f^{\subseteq \emptyset} + (f^{\subseteq \{i\}} - f^{\subseteq \emptyset}) + (f^{\subseteq \{j\}} - f^{\subseteq \emptyset}) + f^{\neg \{i, j\}} \\ \therefore f^{\neg \{i, j\}} &= f^{\subseteq \{i, j\}} - f^{\subseteq \{i\}} - f^{\subseteq \{j\}} + f^{\subseteq \emptyset}. \end{aligned}$$

Inductively, all the $f^{\neg J}$ can be defined via the principle of inclusion-exclusion.

This construction, along with some direct calculations, leads to the following theorem on Efron-Stein decompositions:

Theorem 5.3 ([76, Thm 8.35]). *Let $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$. Then f has a unique decomposition as*

$$f = \sum_{S \subseteq [n]} f^{\neg S}$$

where the functions $f^{\neg S} \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ satisfy

1. $f^{=S}$ depends only on the coordinates in S ;
2. if $T \subsetneq S$ and $g \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ only depends on coordinates in T , then $\langle f^{=S}, g \rangle = 0$.

In addition, this decomposition has the following properties:

3. Condition 2. holds whenever $S \not\subseteq T$.
4. The decomposition is orthogonal: $\langle f^{=S}, f^{=T} \rangle = 0$ for $S \neq T$.
5. $\sum_{S \subseteq T} f^{=S} = f^{=T}$.
6. For each $S \subseteq [n]$, $f \mapsto f^{=S}$ is a linear operator.

Definition 5.4. The Efron-Stein degree of a function $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ is

$$\deg_{\text{ES}}(f) = \max_{S \subseteq [n] \text{ s.t. } f^{=S} \neq 0} |S|.$$

If $f = (f_1, \dots, f_M) : \mathbf{R}^N \rightarrow \mathbf{R}^M$ is a multivariate function, then the Efron-Stein degree of f is the maximum degree of the f_i .

Intuitively, the Efron-Stein degree is the maximum size of multivariate interaction that f accounts for. Of course, this degree is also bounded by N .

As we are interested in how a function behaves under small changes in its input, we are led to consider the following “noise operator.” First, we need the following notion of distance between problem instances:

Definition 5.5. For $p \in [0, 1]$, and $x \in \mathbf{R}^N$, we say $y \in \mathbf{R}^N$ is p -resampled from x if y is chosen as follows: for each $i \in [n]$, independently,

$$y_i = \begin{cases} x_i & \text{with probability } p \\ \text{drawn from } \pi & \text{with probability } 1 - p \end{cases}.$$

We say (x, y) is a p -resampled pair.

Note that being p -resampled and being p -correlated are rather different - for one, there is a nonzero probability that, for π a continuous probability distribution, $x = y$ when they are p -resampled, even though this a.s. never occurs.

Definition 5.6. For $p \in [0, 1]$, the noise operator is the linear operator T_p on $L^2(\mathbf{R}^N, \pi^{\otimes N})$, defined by, for y p -resampled from x

$$T_p f(x) = \mathbf{E}_{y \text{ } p\text{-resampled from } x} [f(y)]$$

In particular, $\langle f, T_p f \rangle = \mathbf{E}_{(x,y) \text{ } p\text{-resampled}} [f(x) \cdot f(y)]$.

Lemma 5.7. Let $p \in [0, 1]$ and $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ have Efron-Stein decomposition $f = \sum_{S \subseteq [n]} f^{=S}$. Then

$$T_p f(x) = \sum_{S \subseteq [n]} p^{|S|} f^{=S}.$$

Proof: Let J denote a p -random subset of $[n]$, i.e. with J formed by including each $i \in [n]$ independently with probability p . By definition, $T_p f(x) = \mathbf{E}_J[f^{\subseteq J}(x)]$ (i.e. pick a random subset of coordinates to fix, and re-randomize the rest). We know by Theorem 5.3 that $f^{\subseteq J} = \sum_{S \subseteq J} f^{\subseteq S}$, so

$$T_p f(x) = \mathbf{E}_J \left[\sum_{S \subseteq J} f^{\subseteq S} \right] = \sum_{S \subseteq [n]} \mathbf{E}_J[I(S \subseteq J)] \cdot f^{\subseteq S} = \sum_{S \subseteq [n]} p^{|S|} f^{\subseteq S},$$

since for a fixed $S \subseteq [n]$, the probability that $S \subseteq J$ is $p^{|S|}$. \square

Putting these facts together, we can derive the following stability bound on functions of bounded Efron-Stein degree.

Theorem 5.8. *Let $p \in [0, 1]$ and let $f = (f_1, \dots, f_M) : \mathbf{R}^N \rightarrow \mathbf{R}^M$ be a multivariate function with Efron-Stein degree D and each $f_i \in L^2(\mathbf{R}^N, \pi^{\otimes N})$. Suppose that (x, y) are a p -resampled pair under $\pi^{\otimes N}$, and $\mathbf{E}\|f(x)\|^2 = 1$. Then*

$$\mathbf{E}\|f(x) - f(y)\|^2 \leq 2(1 - p^D) \leq 2(1 - p)D. \quad (5.3)$$

Proof: Observe that

$$\begin{aligned} \mathbf{E}\|f(x) - f(y)\|^2 &= \mathbf{E}\|f(x)\|^2 + \mathbf{E}\|f(y)\|^2 - 2\mathbf{E}\langle f(x), f(y) \rangle \\ &= 2 - 2 \left(\sum_i \mathbf{E}[f_i(x)f_i(y)] \right) \\ &= 2 - 2 \left(\sum_i \langle f_i, T_p f_i \rangle \right). \end{aligned} \quad (5.4)$$

Here, we have for each $i \in [n]$ that

$$\langle f_i, T_p f_i \rangle = \left\langle \sum_{S \subseteq [n]} f_i^{\subseteq S}, \sum_{S \subseteq [n]} p^{|S|} f_i^{\subseteq S} \right\rangle = \sum_{S \subseteq [n]} p^{|S|} \|f_i^{\subseteq S}\|^2,$$

by Lemma 5.7 and orthogonality. Now, as each f_i has Efron-Stein degree at most D , the sum above can be taken only over $S \subseteq [n]$ with $0 \leq |S| \leq D$, giving the bound

$$p^D \mathbf{E}[f_i(x)^2] \leq \langle f_i, T_p f_i \rangle = \mathbf{E}[f_i(x) \cdot T_p f_i(x)] \leq \mathbf{E}[f_i(x)^2].$$

Summing up over i , and using that $\mathbf{E}\|f(x)\|^2 = 1$, gives

$$p^D \leq \sum_i \langle f_i, T_p f_i \rangle = \mathbf{E}\langle f(x), f(y) \rangle \leq 1.$$

Finally, we can substitute into (5.4) to get

$$\mathbf{E}\|f(x) - f(y)\|^2 \leq 2 - 2p^D = 2(1 - p^D) \leq 2(1 - p)D,$$

¹This follows from the identity $(1 - p^D) = (1 - p)(1 + p + p^2 + \dots + p^{D-1})$; the bound is tight for $p \approx 1$.

as desired. □

5.2. Hermite Polynomials (TODO)

Alternatively, we can consider the much more restrictive (but more concrete) class of honest polynomials. When considered as functions of independent Normal variables, such functions admit a simple description in terms of *Hermite polynomials*, which enables us to prove similar bounds as Theorem 5.8. This theory is much more classical, so we encourage the interested reader to see [76, § 11] for details.

To start, we consider the following space of L^2 functions:

Definition 5.9. Let γ_N be the N -dimensional standard Normal measure on \mathbf{R}^N . Then the N -dimensional Gaussian space is the space $L^2(\mathbf{R}^N, \gamma^N)$ of L^2 functions of N i.i.d. standard Normal random variables.

Note that under the usual L^2 inner product, $\langle f, g \rangle = \mathbf{E}[f \cdot g]$, this is a separable Hilbert space.

It is a well-known fact that the monomials $1, z, z^2, \dots$ form a complete basis for $L^2(\mathbf{R}, \gamma)$ [76, Thm 11.22]. However, these are far from an orthonormal “Fourier” basis; for instance, we know $\mathbf{E}[z^2] = 1$ for $z \sim \mathcal{N}(0, 1)$. By the Gram-Schmidt process, these monomials can be converted into the (normalized) *Hermite polynomials* h_j for $j \geq 0$, given as

$$h_0(z) = 1, \quad h_1(z) = z, \quad h_2(z) = \frac{z^2 - 1}{\sqrt{2}}, \quad h_3(z) = \frac{z^3 - 3z}{\sqrt{6}}, \quad \dots \quad (5.5)$$

Note here that each h_j is a degree j polynomial.

It is then straightforward to show the following:

Theorem 5.10 ([76, Prop 11.30]). *The Hermite polynomials $(h_j)_{j \geq 0}$ form a complete orthonormal basis for $L^2(\mathbf{R}, \gamma)$.*

To extend this to $L^2(\mathbf{R}^N, \gamma^N)$, we can take products. For a multi-index $\alpha \in \mathbb{N}^N$, we define the multivariate Hermite polynomial $h_\alpha : \mathbf{R}^N \rightarrow \mathbf{R}$ as

$$h_\alpha(z) := \prod_{j=1}^N h_{\alpha_j}(z_j).$$

The degree of h_α is clearly $|\alpha| = \sum_j \alpha_j$.

Theorem 5.11. *The Hermite polynomials $(h_\alpha)_{\alpha \in \mathbb{N}^N}$ form a complete orthonormal basis for $L^2(\mathbf{R}^N, \gamma^N)$. In particular, every $f \in L^2(\mathbf{R}^N, \gamma^N)$ has a unique expansion in L^2 norm as*

$$f(z) = \sum_{\alpha \in \mathbb{N}^N} \hat{f}(\alpha) h_\alpha(z).$$

As a consequence of the uniqueness of the expansion in , we see that polynomials are their own Hermite expansion. Namely, let $H^{\leq k} \subseteq L^2(\mathbf{R}^N, \gamma^N)$ be the subset of multivariate polynomials of degree at most k . Then, any $f \in H^{\leq k}$ can be Hermite expanded as

$$f(z) = \sum_{\alpha \in \mathbb{N}^N} \hat{f}(\alpha) h_{\alpha}(z) = \sum_{|\alpha| \leq k} \hat{f}(\alpha) h_{\alpha}(z).$$

Thus, $H^{\leq k}$ is the closed linear span of the set $\{h_{\alpha} : |\alpha| \leq k\}$.

Def. noise operator/Ornstein-Uhlenbeck operator

Compute effect of noise operator on Hermite polys

Thrm. Hermite polys form basis for 1D Gaussian space

Thrm. Products of Hermite polys form basis for N-dim Gaussian space

Noise operator on arbitrary function with given Hermite expansion

Definition 5.12. Let (g, g') be N -dimensional standard Normal vectors. We say (g, g') are p -correlated if (g_i, g'_i) are p -correlated for each $i \in [n]$, and these pairs are mutually independent.

In a similar way to the Efron-Stein case, we can consider the resulting “noise operator,” as a way of measuring the effect on a function of a small change in the input.

Definition 5.13. For $p \in [0, 1]$, the *Gaussian noise operator* T_p is the linear operator on $L^2(\mathbf{R}^N, \gamma^N)$, given by

$$T_p f(x) = \mathbf{E}_{y \text{ } p\text{-correlated to } x} [f(y)] = \mathbf{E}_{y \sim \mathcal{N}(0, I_N)} [f(px + \sqrt{1-p^2}y)]$$

In particular, a straightforward computation with the Normal moment generating function gives

Remark that degree D function can be expressed in terms of degree D and lower Hermite polynomials - gives a Hilbert basis which reflects the natural algebraic grading.

Thrm. Function stability for degree D polynomials.

5.3. Algorithms

Def. Randomized algorithm

Def. degree of algorithm is degree as multivariate function.

Discussion of how low-degree algs are approximate for class of Lipschitz algorithms?

Need for rounding function to land on Σ_N

Construction of randomized rounding function.

Constr. rounded algorithm.

Lemma. stability of rounding

Thrm. Stability of randomized algorithms (part 1 of Prop 1.9)

Show that Markov gives a useful bound on

Lemma 5.14. Let $f : \mathbf{R}^N \rightarrow \mathbf{R}^N$, $p \in [0, 1]$, and X, Y marginally N -dimensional standard Normal vectors. Suppose that $\mathbf{E} \|f(X)\|_2^2 = 1$ and either of the following cases hold:

I. (X, Y) are a p -resampled pair, and f is a degree- D function.

II. (X, Y) are p -correlated, and f is a degree- D polynomial.

Then

$$\mathbf{E}\|f(X) - f(Y)\|_2^2 \leq 2(1 - p^D).$$

5.4. Algorithms

Definition 5.15. A *randomized algorithm* is a measurable function $\mathcal{A}^\circ : (g, \omega) \mapsto \mathbf{x} \in \mathbf{R}^N$, where $\omega \in \Omega_N$ is an independent random variable in some Polish space. Such an \mathcal{A}° is *deterministic* if it does not depend on ω .

Example. Let $\mathbf{U} = (U_1, \dots, U_N)$ be i.i.d. $\text{Unif}([-1, 1])$. Then, we define the random coordinate-wise function

$$\text{round}_{\mathbf{U}}(\mathbf{x}) = (\text{round}_{U_1}(x_1), \dots, \text{round}_{U_N}(x_N)),$$

where we define

$$\text{round}_U(x) = \begin{cases} 1 & x \geq U \\ -1 & x < U \end{cases}.$$

Example. Given a real-valued algorithm \mathcal{A}° , we can convert it into a Σ_N -valued algorithm \mathcal{A} via

$$\mathcal{A}(g, \omega, \mathbf{U}) := \text{round}_{\mathbf{U}}(\mathcal{A}^\circ(g, \omega)).$$

Definition 5.16. Algorithm \mathcal{A} is $(\varepsilon, \eta, p_{\text{unstable}})$ -stable if, for g, g' $(1 - \varepsilon)$ -correlated/resampled, we have

$$\mathbf{P}\left(\|\mathcal{A}(g) - \mathcal{A}(g')\| \leq \eta\sqrt{N}\right) \geq 1 - p_{\text{unstable}}.$$

By the will of God (i.e. writeup pending), we have the following:

Lemma 5.17. Algorithm \mathcal{A} with degree $\leq D$ and norm $\mathbf{E}\|\mathcal{A}(g)\|^2 \leq CN$ has

$$\mathbf{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^2 \leq 2CN\varepsilon D,$$

and (because of randomized rounding)

$$\mathbf{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^4 \leq 16CN^2\varepsilon D.$$

Thus,

$$\mathbf{P}\left(\|\mathcal{A}(g) - \mathcal{A}(g')\| \geq \eta\sqrt{N}\right) \leq \frac{16CN^2\varepsilon D}{\eta^4 N^2} \asymp \frac{\varepsilon D}{\eta^4}.$$

As a consequence, a degree D algorithm \mathcal{A} has $p_{\text{unstable}} = o_{N(1)} \text{ for } \eta^4 \gg \varepsilon D$.

[76]

[60]

6. Summary of Parameters

Parameter	Meaning	Desired Direction	Intuition
N	Dimension	Large	Showing hardness <i>asymptotically</i> , want “bad behavior” to pop up in low dimensions.
δ	Solution tightness; want to find x such that $ \langle g, x \rangle \leq 2^{-\delta N}$	Small	Smaller δ implies weaker solutions, e.g. $\delta = 0$ is just finding solutions ≤ 1 . Want to show even weak solutions are hard to find.
E	Solution tightness; “energy level”; want to find x such that $ \langle g, x \rangle \leq 2^{-E}$	Small	Smaller E implies weaker solutions, and can consider full range of $1 \ll E \ll N$. Know that $E > (\log^2 N)$ by Karmarkar-Karp
D	Algorithm degree (in either Efron-Stein sense or usual polynomial sense.)	Large	Higher degree means more complexity. Want to show even complex algorithms fail.
ε	Complement of correlation/resample probability; (g, g') are $(1 - \varepsilon)$ -correlated.	Small	ε is “distance” between g, g' . Want to show that small changes in disorder lead to “breaking” of landscape.
η	Algorithm instability; \mathcal{A} is stable if $\ \mathcal{A}(g) - \mathcal{A}(g')\ \leq \eta\sqrt{N}$, for (g, g') $(1 - \varepsilon)$ -correlated.	Large	Large η indicates a more unstable algorithm; want to show that even weakly stable algorithms fail.

Table 1: Explanation of Parameters

7. Conditional Landscape Obstruction

Explain what the obstruction is.

We start with a bound on the geometry of the binary hypercube.

Lemma 7.1. *Suppose that $K \leq N/2$, and let $h(x) = -x \log(x) - (1 - x) \log(x)$ be the binary entropy function. Then, for $p := K/N$,*

$$\sum_{k \leq K} \binom{N}{k} \leq \exp(Nh(p)) \leq \exp\left(2Np \log\left(\frac{1}{p}\right)\right).$$

Proof: Consider a $\text{Bin}(N, p)$ random variable S . Summing its PMF from 0 to K , we have

$$1 \geq \mathbf{P}(S \leq K) = \sum_{k \leq K} \binom{N}{k} p^k (1 - p)^{N-k} \geq \sum_{k \leq K} \binom{N}{k} p^K (1 - p)^{N-K}.$$

Here, the last inequality follows from the fact that $p \leq (1 - p)$, and we multiply each term by $\left(\frac{p}{1-p}\right)^{K-k} < 1$. Now rearrange to get

$$\begin{aligned} \sum_{k \leq K} \binom{N}{k} &\leq p^{-K} (1-p)^{-(N-K)} \\ &= \exp(-K \log(p) - (N-K) \log(1-p)) \\ &= \exp\left(N \cdot \left(-\frac{K}{N} \log(p) - \left(\frac{N-K}{N}\right) \log(1-p)\right)\right) \\ &= \exp(N \cdot (-p \log(p) - (1-p) \log(1-p))) = \exp(Nh(p)). \end{aligned}$$

The final equality then follows from the bound $h(p) \leq 2p \log(1/p)$ for $p \leq 1/2$. \square

Lemma 7.2 (Hypercube Neighborhood Size). *Fix $x \in \Sigma_N$, and let $\eta \leq \frac{1}{2}$. Then the number of x' within distance $2\sqrt{\eta N}$ of x is*

$$\left| \{x' \in \Sigma_N \mid \|x - x'\| \leq 2\eta\sqrt{N}\} \right| \leq \exp(2\eta \log(1/\eta)N)$$

Proof: Let k be the number of coordinates which differ between x and x' (i.e. the Hamming distance). We have $\|x - x'\|^2 = 4k$, so $\|x - x'\| \leq 2\eta\sqrt{N}$ iff $k \leq N\eta$. Moreover, for $\eta \leq \frac{1}{2}$, $k \leq \frac{N}{2}$. Thus, by Lemma 7.1, we get

$$\sum_{k \leq N\eta} \binom{N}{k} \leq \exp(Nh(\eta)) \leq \exp(2\eta \log(1/\eta)N). \quad \square$$

Next, we can consider what this probability is in the case of correlated Normals.

Lemma 7.3. *Suppose (g, g') are $(1 - \varepsilon)$ -correlated Normal vectors, and let $x \in \Sigma_N$. Then*

$$\mathbf{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g) \leq \exp\left(-E - \frac{1}{2} \log(\varepsilon) + O(\log N)\right).$$

Proof: Let \tilde{g} be an independent Normal vector to g , and observe that g' can be represented as $g' = pg + \sqrt{1 - p^2}\tilde{g}$, for $p = 1 - \varepsilon$. Thus, $\langle g', x \rangle = p\langle g, x \rangle + \sqrt{1 - p^2}\langle \tilde{g}, x \rangle$. Observe $\langle g, x \rangle$ is constant given g , and $\langle \tilde{g}, x \rangle$ is a Normal r.v. with mean 0 and variance N , so $\langle g', x \rangle \sim \mathcal{N}(p\langle g, x \rangle, (1 - p^2)N)$. This random variable is nondegenerate for $(1 - p^2)N > 0$, with density bounded above as

$$\begin{aligned} \varphi_g(z) &= \frac{1}{\sqrt{2\pi(1 - p^2)N}} e^{-\frac{(z - p\langle g, x \rangle)^2}{2(1 - p^2)N}} \leq \frac{1}{\sqrt{2\pi(1 - p^2)N}} \\ &\leq \frac{1}{\sqrt{2\pi\varepsilon N}} = \exp\left(-\frac{1}{2} \log(\varepsilon) + O(\log N)\right) \end{aligned}$$

Integrating this bound over the interval $|z| \leq 2^{-E}$, we conclude that

$$\mathbf{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g) = \int_{|z| \leq -2^{-E}} \varphi_{g, |S|}(z) dz \leq \exp\left(-E - \frac{1}{2} \log(\varepsilon) + O(\log N)\right). \quad \square$$

Note for instance that here ε can be exponentially small in E (e.g. $\varepsilon = \exp(-E/10)$), which for the case $E = \Theta(N)$ implies ε can be exponentially small in N .

First, we consider the probability of a solution being optimal for a resampled instance.

Lemma 7.4. *Suppose (g, g') are $(1 - \varepsilon)$ -resampled Normal vectors, and let $x \in \Sigma_N$. Then,*

$$\mathbf{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g, g \neq g') \leq 2^{-E+O(1)}.$$

Proof: Let $S = \{i \in [N] : g_i \neq g'_i\}$ be the set of indices where g and g' differ. We can express

$$\langle g', x' \rangle = \sum_{i \in [N]} g'_i x_i = \sum_{i \notin S} g_i x_i + \sum_{i \in S} g'_{i'} x_i \sim \mathcal{N}\left(\sum_{i \notin S} g_i x_i, |S|\right).$$

Let $\mu := \sum_{i \notin S} g_i x_i$. The conditional distribution of interest can now be expressed as $\mathbf{P}(|\langle g', x' \rangle| \leq 2^{-E} \mid g, |S| \geq 1)$. Given $|S| \geq 1$, the quantity $\langle g', x' \rangle$ is a nondegenerate random variable, with density bounded above as

$$\varphi_{g, |S|}(z) = \frac{1}{\sqrt{2\pi|S|}} e^{-\frac{(z-\mu)^2}{2|S|}} \leq \frac{1}{\sqrt{2\pi|S|}} \leq \frac{1}{\sqrt{2\pi}}.$$

Hence, the quantity of interest can be bounded as

$$\mathbf{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g, g \neq g') \leq \int_{|z| \leq -2^{-E}} \varphi_{g, |S|}(z) dz \leq \sqrt{\frac{2}{\pi}} 2^{-E} = 2^{-E+O(1)}. \quad \square$$

Note that in the resampled case, we can compute the probability that $g = g'$ as

$$\mathbf{P}(g = g') = \prod_{i=1}^N \mathbf{P}(g_i = g_{i'}) = (1 - \varepsilon)^N,$$

which for $\varepsilon \ll 1$ is approximately $1 - N\varepsilon$. Thus, for $\varepsilon \gg \omega\left(\frac{1}{N}\right)$, we have

$$\mathbf{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g) \leq 2^{-E+O(1)}$$

8. Proof of Low-Degree Hardness in Linear Energy Regime.

Throughout this section, we let $E = \delta N$ for some $\delta > 0$, and aim to rule out the existence of low-degree algorithms achieving these energy levels. This corresponds to the statistically optimal regime, as per [54].

For now, let \mathcal{A} denote a Σ_N -valued deterministic algorithm. We discuss the extension to random, \mathbf{R}^N -valued algorithms later on in (section ???).

8.1. Hermite Case

First, we consider

Assume for sake of contradiction that $p_{\text{solve}} \geq \Omega(1)$.

Let g, g' be $(1 - \varepsilon)$ -correlated instances. We define the following events:

$$\begin{aligned} S_{\text{solve}} &= \{\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g')\} \\ S_{\text{stable}} &= \{\|\mathcal{A}(g) - \mathcal{A}(g')\| \leq 2\sqrt{\eta N}\} \\ S_{\text{cond}} &= \left\{ \nexists x' \in S(E; g') \text{ such that } \right. \\ &\quad \left. \|x - x'\| \leq 2\sqrt{\eta N} \right\} \end{aligned}$$

In this case, set $\varepsilon = 2^{-\delta N} = o(1)$

Lemma 8.1. *There exists an $\eta > 0$ of constant order such that*

$$\mathbf{P}(S_{\text{cond}}) \geq 1 - e^{-cN}$$

for an arbitrary constant c .

$$D = o(2^N).$$

$$D\varepsilon = \frac{D}{2^N} * 2^{(1-\delta)N}$$

Lemma 8.2. *For any $\omega(\log^2 N) \leq E \leq \Theta(N)$, there exist choices of ε, η (depending on N, E) such that $\mathbf{P}(S_{\text{ogp}}) = o(1)$.*

Proof: Observe that

$$\mathbf{P}(S_{\text{ogp}}) = \mathbf{E}[\mathbf{P}(S_{\text{ogp}} | g)]. \quad (8.1)$$

Conditional on g , we can compute $\mathbf{P}(S_{\text{ogp}} | g) = \mathbf{P}(\exists x' \in S(E; g'), \|x - x'\| \leq 2\eta\sqrt{N})$ by setting $x = \mathcal{A}(g)$ (so x only depends on g), and union bounding Lemma 7.4 over the x' within $2\eta\sqrt{N}$ of x , as per Lemma 7.2:

$$\mathbf{P}(S_{\text{ogp}} | g) \leq \exp_2(-E + N\eta^2 \log_2(1/\eta^2) + O(1)).$$

We want to choose η such that

$$-E + N\eta^2 \log_2(1/\eta^2) = -\Omega(N)$$

$$\frac{E}{N} > \eta^2 \log(1/\eta^2)$$

Using the fact that $\sqrt{2x} \geq -x \log_2 x$, it suffices to pick η^2 with

$$\frac{E}{N} > 2\eta,$$

so $\eta^2 := \frac{E^2}{2N^2}$ is a valid choice.

By the choice of $\eta^2 \ll (h^{-1})(\delta) \asymp 1$, this bound gives $\mathbf{P}(S_{\text{ogp}}|g) \leq \exp_2(-O(N)) = o(1)$. Integrating over g gives the overall bound. \square

When $CD\epsilon N^2 = \omega_{N(1)}$ (i.e. goes to infinity),

$$\begin{aligned} \mathbf{P}(S_{\text{stable}}) &\leq \frac{16CD\epsilon N^2}{16\eta^4 N^2} \\ &= \frac{CD\epsilon}{\eta^4} = \frac{4CD\epsilon N^4}{E^4} \end{aligned}$$

$D\epsilon \rightarrow 0$ same as $D = o(\frac{1}{\epsilon}) = o(N)$.

Lemma 8.3. $\mathbf{P}(S_{\text{solve}}, S_{\text{stable}}) \leq \mathbf{P}(S_{\text{ogp}}) = o(1)$.

Proof: The first inequality follows from definition, with $x = \mathcal{A}(g)$ and $x' = \mathcal{A}(g')$. For the second, observe that

$$\mathbf{P}(S_{\text{ogp}}) = \mathbf{E}[\mathbf{P}(S_{\text{ogp}} | g)].$$

Now, let $M = \delta N$, we can compute $\mathbf{P}(S_{\text{ogp}} | g) = \mathbf{P}(\exists x' \in S(\delta; g'), \|x - x'\| \leq \eta\sqrt{N})$ by setting $x = \mathcal{A}(g)$ (so x only depends on g), and union bounding Lemma 7.4 over the x' within $\eta\sqrt{N}$ of x , as per Lemma 7.2:

$$\mathbf{P}(S_{\text{ogp}} | g) \leq \exp_2\left(-\delta N + Nh\left(\frac{\eta^2}{4}\right) + O(\log_2 N)\right).$$

By the choice of $\eta^2 \ll (h^{-1})(\delta) \asymp 1$, this bound gives $\mathbf{P}(S_{\text{ogp}}|g) \leq \exp_2(-O(N)) = o(1)$. Integrating over g gives the overall bound. \square

However, by the choice of parameters above, we also have

$$\begin{aligned} \mathbf{P}(S_{\text{solve}}, S_{\text{stable}}) &\geq \mathbf{P}(S_{\text{solve}}) + \mathbf{P}(S_{\text{stable}}) - 1 \\ &\geq p_{\text{solve}}^4 + p_{\text{unstable}} \geq \Omega(1) - o(1) = \Omega(1), \end{aligned} \tag{8.2}$$

which is a contradiction.

9. Randomized Rounding Things

Claim: no two adjacent points on Σ_N (or pairs within $k = O(1)$ distance) which are both good solutions to the same problem. The reason is that this would require a subset of k signed coordinates $\pm g_{\{i_1\}}, \dots, \pm g_{\{i_k\}}$ to have small sum, and there are only $2^k \text{binom}\{N\}{k} = O(N^k)$ possibilities, each of which is centered Gaussian with variance at least 1, so the smallest is typically of order $\Omega(N^{\{-k\}})$.

Thus, rounding would destroy the solution.

- Say we're in the case where rounding changes the solution. (i.e. rounding moves x to point that is not the closest point x_* , whp.)

- Let p_1, \dots, p_N be the probabilities of disagreeing with x_* on each coordinate.
 - We know that $\sum p_i$ diverges (this is equivalent to the statement that rounding will change the solution whp).
 - Reason: for each coord, chance of staying at that coordinate is $e^{-\Theta(p_i)}$.
- For each i , flip coin with heads prob $2p_i$, and keep all the heads.
 - By Borel-Cantelli type argument, typical number of heads will be $\omega(1)$.
- For every coin with a head, change coord with prob 50%, if tails, keep coord.
 - Randomized rounding in artificially difficult way. (I.e. this multistage procedure accomplishes the same thing as randomized rounding.)
- Now, randomized rounding is done by choosing a random set of $\omega(1)$ coordinates, and making those iid Uniform in $\{-1, 1\}$.
- Pick a large constant (e.g. 100), and only randomize the first 100 heads, and condition on the others (i.e. choose the others arbitrarily). Note that since $100l = \omega(1)$, there are at least 100 heads whp.
- Now rounded point is random point in 100 dimensional subcube, but at most one of them is a good solution by the claim at the top of the page.
- Combining, the probability for rounding to give a good solution is at most $o(1) + 2^{\{-100\}}$. Since 100 is arbitrary, this is $o(1)$ by sending parameters to 0 and/or infinity in the right order.

[76]

Bibliography

- [1] D. Achlioptas and A. Coja-Oghlan, “Algorithmic Barriers from Phase Transitions,” in *2008 49th Annual IEEE Symposium on Foundations of Computer Science*, Oct. 2008, pp. 793–802. doi: 10.1109/FOCS.2008.11.
- [2] D. Achlioptas and F. Ricci-Tersenghi, “On the Solution-Space Geometry of Random Constraint Satisfaction Problems.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/cs/0611052>
- [3] L. Addario-Berry, L. Devroye, G. Lugosi, and R. I. Oliveira, “Local Optima of the Sherrington-Kirkpatrick Hamiltonian.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1712.07775>
- [4] B. Alidaee, F. Glover, G. A. Kochenberger, and C. Rego, “A New Modeling and Solution Approach for the Number Partitioning Problem,” *Journal of Applied Mathematics and Decision Sciences*, vol. 2005, no. 2, pp. 113–121, Jan. 2005, doi: 10.1155/JAMDS.2005.113.
- [5] M. F. Argüello, T. A. Feo, and O. Goldschmidt, “Randomized Methods for the Number Partitioning Problem,” *Computers & Operations Research*, vol. 23, no. 2, pp. 103–111, Feb. 1996, doi: 10.1016/0305-0548(95)E0020-L.
- [6] L. Asproni, D. Caputo, B. Silva, G. Fazzi, and M. Magagnini, “Accuracy and Minor Embedding in Subqubo Decomposition with Fully Connected Large Problems: A Case Study about the Number Partitioning Problem,” *Quantum Machine Intelligence*, vol. 2, no. 1, p. 4, Jun. 2020, doi: 10.1007/s42484-020-00014-w.

- [7] B. Aubin, W. Perkins, and L. Zdeborová, “Storage Capacity in Symmetric Binary Perceptrons,” *Journal of Physics A: Mathematical and Theoretical*, vol. 52, no. 29, p. 294003, Jul. 2019, doi: 10.1088/1751-8121/ab227a.
- [8] A. S. Bandeira, A. Perry, and A. S. Wein, “Notes on Computational-to-Statistical Gaps: Predictions Using Statistical Physics.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1803.11132>
- [9] N. Bansal, “Constructive Algorithms for Discrepancy Minimization.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1002.2259>
- [10] B. Barak, S. B. Hopkins, J. Kelner, P. K. Kothari, A. Moitra, and A. Potechin, “A Nearly Tight Sum-of-Squares Lower Bound for the Planted Clique Problem.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1604.03084>
- [11] H. Bauke, S. Franz, and S. Mertens, “Number Partitioning as a Random Energy Model,” *Journal of Statistical Mechanics: Theory and Experiment*, vol. 2004, no. 4, p. P4003, Apr. 2004, doi: 10.1088/1742-5468/2004/04/P04003.
- [12] M. Bayati, D. Gamarnik, and P. Tetali, “Combinatorial Approach to the Interpolation Method and Scaling Limits in Sparse Random Graphs,” *The Annals of Probability*, vol. 41, no. 6, Nov. 2013, doi: 10.1214/12-AOP816.
- [13] Q. Berthet and P. Rigollet, “Computational Lower Bounds for Sparse PCA.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1304.0828>
- [14] S. Boettcher and S. Mertens, “Analysis of the Karmarkar-Karp Differencing Algorithm,” *The European Physical Journal B*, vol. 65, no. 1, pp. 131–140, Sep. 2008, doi: 10.1140/epjb/e2008-00320-9.
- [15] C. Borgs, J. Chayes, and B. Pittel, “Phase Transition and Finite-size Scaling for the Integer Partitioning Problem,” *Random Structures & Algorithms*, vol. 19, no. 3–4, pp. 247–288, Oct. 2001, doi: 10.1002/rsa.10004.
- [16] M. Brennan and G. Bresler, “Optimal Average-Case Reductions to Sparse PCA: From Weak Assumptions to Strong Hardness.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1902.07380>
- [17] M. Brennan, G. Bresler, and W. Huleihel, “Reducibility and Computational Lower Bounds for Problems with Planted Sparse Structure.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1806.07508>
- [18] K. Chandrasekaran and S. Vempala, “Integer Feasibility of Random Polytopes.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1111.4649>
- [19] W.-K. Chen, D. Gamarnik, D. Panchenko, and M. Rahman, “Suboptimality of Local Algorithms for a Class of Max-Cut Problems,” *The Annals of Probability*, vol. 47, no. 3, May 2019, doi: 10.1214/18-AOP1291.

- [20] E. G. Coffman Jr., M. R. Garey, and D. S. Johnson, “An Application of Bin-Packing to Multiprocessor Scheduling,” *SIAM Journal on Computing*, vol. 7, no. 1, pp. 1–17, Feb. 1978, doi: 10.1137/0207001.
- [21] E. G. Coffman and G. S. Lueker, *Probabilistic Analysis of Packing and Partitioning Algorithms*. in Wiley-Interscience Series in Discrete Mathematics and Optimization. New York: J. Wiley & sons, 1991.
- [22] A. Coja-Oghlan and C. Efthymiou, “On Independent Sets in Random Graphs,” *Random Structures & Algorithms*, vol. 47, no. 3, pp. 436–486, Oct. 2015, doi: 10.1002/rsa.20550.
- [23] D. Corus, P. S. Oliveto, and D. Yazdani, “Artificial Immune Systems Can Find Arbitrarily Good Approximations for the NP-hard Number Partitioning Problem,” *Artificial Intelligence*, vol. 274, pp. 180–196, Sep. 2019, doi: 10.1016/j.artint.2019.03.001.
- [24] Y. Deshpande and A. Montanari, “Improved Sum-of-Squares Lower Bounds for Hidden Clique and Hidden Submatrix Problems.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1502.06590>
- [25] I. Diakonikolas, D. M. Kane, and A. Stewart, “Statistical Query Lower Bounds for Robust Estimation of High-dimensional Gaussians and Gaussian Mixtures.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1611.03473>
- [26] V. Feldman, E. Grigorescu, L. Reyzin, S. Vempala, and Y. Xiao, “Statistical Algorithms and a Lower Bound for Detecting Planted Clique.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1201.1214>
- [27] F. F. Ferreira and J. F. Fontanari, “Probabilistic Analysis of the Number Partitioning Problem,” *Journal of Physics A: Mathematical and General*, vol. 31, no. 15, p. 3417, Apr. 1998, doi: 10.1088/0305-4470/31/15/007.
- [28] D. Gamarnik and E. C. Kızıldağ, “Algorithmic Obstructions in the Random Number Partitioning Problem.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2103.01369>
- [29] D. Gamarnik, E. C. Kızıldağ, W. Perkins, and C. Xu, “Algorithms and Barriers in the Symmetric Binary Perceptron Model.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2203.15667>
- [30] D. Gamarnik and E. Kizildag, “Computing the Partition Function of the Sherrington-Kirkpatrick Model Is Hard on Average,” *The Annals of Applied Probability*, vol. 31, no. 3, Jun. 2021, doi: 10.1214/20-AAP1625.
- [31] D. Gamarnik, A. Jagannath, and A. S. Wein, “Hardness of Random Optimization Problems for Boolean Circuits, Low-Degree Polynomials, and Langevin Dynamics.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2004.12063>
- [32] D. Gamarnik and I. Zadik, “High-Dimensional Regression with Binary Coefficients. Estimating Squared Error and a Phase Transition.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1701.04455>

- [33] D. Gamarnik and I. Zadik, “The Landscape of the Planted Clique Problem: Dense Subgraphs and the Overlap Gap Property.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1904.07174>
- [34] D. Gamarnik and M. Sudan, “Limits of Local Algorithms over Sparse Random Graphs.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/1304.1831>
- [35] D. Gamarnik and A. Jagannath, “The Overlap Gap Property and Approximate Message Passing Algorithms for \mathbb{Z}_p -Spin Models.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1911.06943>
- [36] D. Gamarnik, “The Overlap Gap Property: A Geometric Barrier to Optimizing over Random Structures,” *Proceedings of the National Academy of Sciences*, vol. 118, no. 41, p. e2108492118, Oct. 2021, doi: 10.1073/pnas.2108492118.
- [37] D. Gamarnik, A. Jagannath, and S. Sen, “The Overlap Gap Property in Principal Submatrix Recovery,” *Probability Theory and Related Fields*, vol. 181, no. 4, pp. 757–814, Dec. 2021, doi: 10.1007/s00440-021-01089-7.
- [38] D. Gamarnik and M. Sudan, “Performance of Sequential Local Algorithms for the Random NAE- \mathbb{Z}_K -SAT Problem,” *SIAM Journal on Computing*, vol. 46, no. 2, pp. 590–619, Jan. 2017, doi: 10.1137/140989728.
- [39] D. Gamarnik and I. Zadik, “Sparse High-Dimensional Linear Regression. Algorithmic Barriers and a Local Search Algorithm.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1711.04952>
- [40] M. R. Garey and D. S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*. in A Series of Books in the Mathematical Sciences. New York: W. H. Freeman, 1979.
- [41] I. P. Gent and T. Walsh, “Analysis of Heuristics for Number Partitioning,” *Computational Intelligence*, vol. 14, no. 3, pp. 430–451, 1998, doi: 10.1111/0824-7935.00069.
- [42] I. Gent and T. Walsh, “Phase Transitions and Annealed Theories: Number Partitioning as a Case Study,” *Instituto Cultura*, Jun. 2000.
- [43] C. Harshaw, F. Sävje, D. Spielman, and P. Zhang, “Balancing Covariates in Randomized Experiments with the Gram-Schmidt Walk Design.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/1911.03071>
- [44] T. Hastie, R. Tibshirani, and J. Friedman, *The Elements of Statistical Learning*. in Springer Series in Statistics. New York, NY: Springer New York, 2009. doi: 10.1007/978-0-387-84858-7.
- [45] H. Hatami, L. Lovász, and B. Szegedy, “Limits of Locally–Globally Convergent Graph Sequences,” *Geometric and Functional Analysis*, vol. 24, no. 1, pp. 269–296, Feb. 2014, doi: 10.1007/s00039-014-0258-7.
- [46] R. Hoberg, H. Ramadas, T. Rothvoss, and X. Yang, “Number Balancing Is as Hard as Minkowski’s Theorem and Shortest Vector.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/1611.08757>

- [47] S. B. Hopkins, P. K. Kothari, A. Potechin, P. Raghavendra, T. Schramm, and D. Steurer, “The Power of Sum-of-Squares for Detecting Hidden Structures.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1710.05017>
- [48] S. Hopkins, “Statistical Inference and the Sum of Squares Method,” 2018.
- [49] S. B. Hopkins, J. Shi, and D. Steurer, “Tensor Principal Component Analysis via Sum-of-Squares Proofs.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1507.03269>
- [50] B. Huang and M. Sellke, “Strong Low Degree Hardness for Stable Local Optima in Spin Glasses.” Accessed: Jan. 30, 2025. [Online]. Available: <http://arxiv.org/abs/2501.06427>
- [51] M. Jerrum, “Large Cliques Elude the Metropolis Process,” *Random Structures & Algorithms*, vol. 3, no. 4, pp. 347–359, Jan. 1992, doi: 10.1002/rsa.3240030402.
- [52] D. S. Johnson, C. R. Aragon, L. A. McGeoch, and C. Schevon, “Optimization by Simulated Annealing: An Experimental Evaluation; Part I, Graph Partitioning,” *Operations Research*, vol. 37, no. 6, pp. 865–892, 1989, Accessed: Mar. 15, 2025. [Online]. Available: <http://www.jstor.org/stable/171470>
- [53] D. S. Johnson, C. R. Aragon, L. A. McGeoch, and C. Schevon, “Optimization by Simulated Annealing: An Experimental Evaluation; Part II, Graph Coloring and Number Partitioning,” *Operations Research*, vol. 39, no. 3, pp. 378–406, 1991, Accessed: Mar. 15, 2025. [Online]. Available: <http://www.jstor.org/stable/171393>
- [54] N. Karmarkar, R. M. Karp, G. S. Lueker, and A. M. Odlyzko, “Probabilistic Analysis of Optimum Partitioning,” *Journal of Applied Probability*, vol. 23, no. 3, pp. 626–645, 1986, doi: 10.2307/3214002.
- [55] N. Karmarkar and R. M. Karp, “The Differencing Method of Set Partitioning,” 1983. Accessed: Mar. 15, 2025. [Online]. Available: <https://www2.eecs.berkeley.edu/Pubs/TechRpts/1983/6353.html>
- [56] M. Kearns, “Efficient Noise-Tolerant Learning from Statistical Queries,” *Journal of the ACM*, vol. 45, no. 6, pp. 983–1006, Nov. 1998, doi: 10.1145/293347.293351.
- [57] E. C. Kızıldağ, “Planted Random Number Partitioning Problem.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2309.15115>
- [58] J. Kojić, “Integer Linear Programming Model for Multidimensional Two-Way Number Partitioning Problem,” *Computers & Mathematics with Applications*, vol. 60, no. 8, pp. 2302–2308, Oct. 2010, doi: 10.1016/j.camwa.2010.08.024.
- [59] R. E. Korf, “From Approximate to Optimal Solutions: A Case Study of Number Partitioning,” in *Proceedings of the 14th International Joint Conference on Artificial Intelligence - Volume 1*, in IJCAI'95. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc., Aug. 1995, pp. 266–272.
- [60] R. E. Korf, “A Complete Anytime Algorithm for Number Partitioning,” *Artificial Intelligence*, vol. 106, no. 2, pp. 181–203, Dec. 1998, doi: 10.1016/S0004-3702(98)00086-1.

- [61] R. E. Korf, "Multi-Way Number Partitioning," in *Proceedings of the 21st International Joint Conference on Artificial Intelligence*, in IJCAI'09. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc., Jul. 2009, pp. 538–543.
- [62] P. K. Kothari, R. Mori, R. O'Donnell, and D. Witmer, "Sum of Squares Lower Bounds for Refuting Any CSP." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1701.04521>
- [63] J. Kratica, J. Kojić, and A. Savić, "Two Metaheuristic Approaches for Solving Multidimensional Two-Way Number Partitioning Problem," *Computers & Operations Research*, vol. 46, pp. 59–68, Jun. 2014, doi: 10.1016/j.cor.2014.01.003.
- [64] A. M. Krieger, D. Azriel, and A. Kapelner, "Nearly Random Designs with Greatly Improved Balance," *Biometrika*, vol. 106, no. 3, pp. 695–701, Sep. 2019, doi: 10.1093/biomet/asz026.
- [65] D. Kunisky, A. S. Wein, and A. S. Bandeira, "Notes on Computational Hardness of Hypothesis Testing: Predictions Using the Low-Degree Likelihood Ratio." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1907.11636>
- [66] J. Lauer and N. Wormald, "Large Independent Sets in Regular Graphs of Large Girth," *Journal of Combinatorial Theory, Series B*, vol. 97, no. 6, pp. 999–1009, Nov. 2007, doi: 10.1016/j.jctb.2007.02.006.
- [67] A. Levy, H. Ramadas, and T. Rothvoss, "Deterministic Discrepancy Minimization via the Multiplicative Weight Update Method." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1611.08752>
- [68] S. Lovett and R. Meka, "Constructive Discrepancy Minimization by Walking on The Edges." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1203.5747>
- [69] G. S. Lueker, "A Note on the Average-Case Behavior of a Simple Differencing Method for Partitioning," *Operations Research Letters*, vol. 6, no. 6, pp. 285–287, Dec. 1987, doi: 10.1016/0167-6377(87)90044-7.
- [70] R. Meka, A. Potechin, and A. Wigderson, "Sum-of-Squares Lower Bounds for Planted Clique," in *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing*, Portland Oregon USA: ACM, Jun. 2015, pp. 87–96. doi: 10.1145/2746539.2746600.
- [71] R. Merkle and M. Hellman, "Hiding Information and Signatures in Trapdoor Knapsacks," *IEEE Transactions on Information Theory*, vol. 24, no. 5, pp. 525–530, Sep. 1978, doi: 10.1109/TIT.1978.1055927.
- [72] S. Mertens, "The Easiest Hard Problem: Number Partitioning." Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/cond-mat/0310317>
- [73] S. Mertens, "A Physicist's Approach to Number Partitioning," *Theoretical Computer Science*, vol. 265, no. 1, pp. 79–108, Aug. 2001, doi: 10.1016/S0304-3975(01)00153-0.
- [74] M. Mézard, T. Mora, and R. Zecchina, "Clustering of Solutions in the Random Satisfiability Problem," *Physical Review Letters*, vol. 94, no. 19, p. 197205, May 2005, doi: 10.1103/PhysRevLett.94.197205.

- [75] W. Michiels, J. Korst, E. Aarts, and J. Van Leeuwen, "Performance Ratios for the Differencing Method Applied to the Balanced Number Partitioning Problem," *STACS 2003*, vol. 2607. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 583–595, 2003. doi: 10.1007/3-540-36494-3_51.
- [76] R. O'Donnell, "Analysis of Boolean Functions." Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2105.10386>
- [77] P. Raghavendra, T. Schramm, and D. Steurer, "High-Dimensional Estimation via Sum-of-Squares Proofs." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1807.11419>
- [78] M. Rahman and B. Virag, "Local Algorithms for Independent Sets Are Half-Optimal," *The Annals of Probability*, vol. 45, no. 3, May 2017, doi: 10.1214/16-AOP1094.
- [79] T. Rothvoss, "Constructive Discrepancy Minimization for Convex Sets." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1404.0339>
- [80] V. Santucci, M. Baiocchi, and G. Di Bari, "An Improved Memetic Algebraic Differential Evolution for Solving the Multidimensional Two-Way Number Partitioning Problem," *Expert Systems with Applications*, vol. 178, p. 114938, Sep. 2021, doi: 10.1016/j.eswa.2021.114938.
- [81] R. H. Storer, S. W. Flanders, and S. David Wu, "Problem Space Local Search for Number Partitioning," *Annals of Operations Research*, vol. 63, no. 4, pp. 463–487, Aug. 1996, doi: 10.1007/BF02156630.
- [82] L.-H. Tsai, "Asymptotic Analysis of an Algorithm for Balanced Parallel Processor Scheduling," *SIAM Journal on Computing*, vol. 21, no. 1, pp. 59–64, Feb. 1992, doi: 10.1137/0221007.
- [83] P. Turner, R. Meka, and P. Rigollet, "Balancing Gaussian Vectors in High Dimension." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1910.13972>
- [84] M. J. Wainwright, *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. in Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press, 2019. doi: 10.1017/9781108627771.
- [85] A. S. Wein, "Optimal Low-Degree Hardness of Maximum Independent Set." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/2010.06563>
- [86] J. Wen *et al.*, "Optical Experimental Solution for the Multiway Number Partitioning Problem and Its Application to Computing Power Scheduling," *Science China Physics, Mechanics & Astronomy*, vol. 66, no. 9, p. 290313, Sep. 2023, doi: 10.1007/s11433-023-2147-3.
- [87] B. Yakir, "The Differencing Algorithm LDM for Partitioning: A Proof of a Conjecture of Karmarkar and Karp," *Mathematics of Operations Research*, vol. 21, no. 1, pp. 85–99, Feb. 1996, doi: 10.1287/moor.21.1.85.
- [88] L. Zdeborová and F. Krzakala, "Statistical Physics of Inference: Thresholds and Algorithms," *Advances in Physics*, vol. 65, no. 5, pp. 453–552, Sep. 2016, doi: 10.1080/00018732.2016.1211393.