

# 1. Number Packing Problem

Let  $N$  be the dimensionality, and  $\Sigma_N := \{\pm 1\}$  be the binary cube. Suppose we're given  $g \sim \mathcal{N}(0, I_N)$ . We want to find  $x \in \Sigma_N$  such that we minimize  $|\langle x, g \rangle|$ .

Definition 1.1: Let  $\delta > 0$ . The *solution set*  $S(\delta; g)$  is the set of all  $x \in \Sigma_N$  that are  $\delta$ -optimal, satisfying

$$\frac{1}{\sqrt{N}} |\langle g, x \rangle| \leq 2^{-\delta N}. \quad (1.1)$$

(1.1) Hi

## 1.1. Existing Results

1.  $X_i, 1 \leq i \leq n$  i.i.d. uniform from  $\{1, 2, \dots, M := 2^m\}$ , with  $\kappa := \frac{m}{n}$ , then phase transition going from  $\kappa < 1$  to  $\kappa > 1$ .
2. Average case,  $X_i$  i.i.d. standard Normal.
3. Karmarkar [KKLO86] - NPP value is  $\Theta(\sqrt{N}2^{-N})$  whp as  $N \rightarrow \infty$  (doesn't need Normality).
4. Best polynomial-time algorithm: Karmarkar-Karp [KK82] - Discrepancy  $O(N^{-\alpha \log N}) = 2^{-\Theta(\log^2 N)}$  whp as  $N \rightarrow \infty$
5. PDM (paired differencing) heuristic - fails for i.i.d. uniform inputs with objective  $\Theta(n^{-1})$  (Lueker).
6. LDM (largest differencing) heuristic - works for i.i.d. Uniforms, with  $n^{-\Theta(\log n)}$  (Yakir, with constant  $\alpha = \frac{1}{2 \ln 2}$  calculated non-rigorously by Boettcher and Mertens).
7. Krieger -  $O(n^{-2})$  for balanced partition.
8. Hoberg [HHRY17] - computational hardness for worst-case discrepancy, as poly-time oracle that can get discrepancy to within  $O(2^{\sqrt{n}})$  would be oracle for Minkowski problem.
9. Gamarnik-Kizildag: Information-theoretic guarantee  $E_n = n$ , best computational guarantee  $E_n = \Theta(\log^2 n)$ .
10. Existence of  $m$ -OGP for  $m = O(1)$  and  $E_n = \Theta(n)$ .
11. Absence for  $\omega(1) \leq E_n = o(n)$
12. Existence for  $\omega(\sqrt{n \log_2 n}) \leq E_n \leq o(n)$  for  $m = \omega_{n(1)}$  (with changing  $\eta, \beta$ )
  1. While OGP not ruled out for  $E_n \leq \omega(\sqrt{n \log_2 n})$ , argued that it is tight.
13. For  $\varepsilon \in (0, \frac{1}{5})$ , no stable algorithm can solve  $\omega(n \log^{-\frac{1}{5} + \varepsilon} n) \leq E_n \leq o(n)$
14. Possible to strengthen to  $E_n = \Theta(n)$  (as  $2^{-\Theta(n)} \leq 2^{-o(n)}$ )

## 2. Glossary and conventions

Conventions:

1. log means log in base 2, exp is exp base 2 - natural log/exponent is  $\ln/e^x$ .
- 2.

Glossary:

1. "instance"/"disorder" -  $g$ , instance of the NPP problem

2. “discrepancy” - for a given  $g$ , value of  $\min_{x \in \Sigma_N} |\langle g, x \rangle|$
3. “energy” - negative exponent of discrepancy, i.e. if discrepancy is  $2^{-E}$ , then energy is  $E$ . Lower energy indicates “worse” discrepancy.
4. “near-ground state”/“approximate solution”

### 3. Low-Degree Algorithms

What are algorithms of interest

For our purposes, an *algorithm* is a function which takes as input

Why study low-degree algorithms (poly time heuristic + simple)

Different notions of degree.

Overview of section

- Efron-Stein notion
- Hermite notion
- Algorithms and Stability Bounds

#### 3.1. Efron-Stein Polynomials (TODO)

Definition 3.1.1: Let  $\pi$  be a probability distribution on  $\mathbb{R}$ . The  $L^2$  space  $L^2(\mathbb{R}^N, \pi^{\otimes N})$  is the space of functions  $f : \mathbb{R}^N \rightarrow \mathbb{R}$  with finite  $L^2$  norm.

$$\mathbb{E}[f^2] := \int_{x=(x_1, \dots, x_n) \in \mathbb{R}^N} f(x)^2 d\pi^{\otimes N}(x) < \infty.$$

Alternatively, this is the space of  $L^2$  functions of  $N$  i.i.d. random variables  $x_i$  distributed as  $\pi$ .

Motivation for studying decompositions of functions by projecting onto coordinates.

Want to decompose

$$f = \sum_{S \subseteq [n]} f^{\perp S} \tag{3.1}$$

Want  $f^{\perp S}$  to only depend on the coordinates in  $S$ .

If  $T \not\subseteq S$  and  $g$  depends only on the coordinates in  $T$ , then  $\langle f^{\perp S}, g \rangle = 0$ .

Definition 3.1.2: Let  $f \in L^2(\mathbb{R}^N, \pi^{\otimes N})$  and  $J \subseteq [n]$ , with  $\bar{J} = [n] \setminus J$ . The *projection of  $f$  onto  $J$*  is the function  $f^{\subseteq J} : \mathbb{R}^N \rightarrow \mathbb{R}$  given by

$$f^{\subseteq J}(x) = \mathbb{E}[f(x_1, \dots, x_n) \mid x_i, i \in J].$$

This is  $f$  with the  $\bar{J}$  coordinates re-randomized, so  $f^{\subseteq J}$  only depends on  $x_J$ .

In particular, we should have that

$$f^{\subseteq J} = \sum_{S \subseteq J} f^{=S} \quad (3.2)$$

First, we consider the case  $J = \emptyset$ . It is clear that  $f^{\subseteq \emptyset} = f^{=\emptyset}$ , which is the constant function  $\mathbb{E}[f]$ .

Next, if  $J = \{j\}$  is a singleton, (3.2) gives

$$f^{\subseteq \{j\}} = f^{=\emptyset} + f^{=\{j\}},$$

and as  $f^{\subseteq \{j\}}(x) = \mathbb{E}[f \mid x_j]$ , we get

$$f^{=\{j\}} = \mathbb{E}[f \mid x_j] - \mathbb{E}[f].$$

This function only depends on  $x_j$ ; all other coordinates are averaged over. It measures what difference in expectation of  $f$  is given  $x_j$ .

Continuing on to sets of two coordinates, some brief manipulation gives, for  $J = \{i, j\}$ ,

$$\begin{aligned} f^{\subseteq \{i, j\}} &= f^{=\emptyset} + f^{=\{i\}} + f^{=\{j\}} + f^{=\{i, j\}} \\ &= f^{\subseteq \emptyset} + (f^{\subseteq \{i\}} - f^{\subseteq \emptyset}) + (f^{\subseteq \{j\}} - f^{\subseteq \emptyset}) + f^{=\{i, j\}} \\ \therefore f^{=\{i, j\}} &= f^{\subseteq \{i, j\}} - f^{\subseteq \{i\}} - f^{\subseteq \{j\}} + f^{\subseteq \emptyset}. \end{aligned}$$

Inductively, all the  $f^{=J}$  can be defined via the principle of inclusion-exclusion.

To see that these functions are indeed orthogonal, we need the following computation:

**Lemma 3.1.3:** Let  $f, g \in L^2(\mathbb{R}^N, \pi^{\otimes N})$  and  $I, J \subseteq [n]$  be subsets of coordinates. Assume that  $f$  only depends on coordinates in  $I$  and likewise for  $g$  and  $J$ . Then  $\langle f, g \rangle = \langle f^{\subseteq I \cap J}, g^{\subseteq I \cap J} \rangle$ .

*Proof:* Assume without loss of generality that  $I \cup J = [n]$ . Then, given  $x \in \mathbb{R}^N$ , we can split it into  $(x_{I \cap J}, x_{I \setminus J}, x_{J \setminus I})$ . Abusing notation slightly to only include the coordinates a function actually depends on, we have

$$\begin{aligned} \langle f, g \rangle &= \mathbb{E}_{x_{I \cap J}, x_{I \setminus J}, x_{J \setminus I}} [f(x_{I \cap J}, x_{I \setminus J}) \cdot g(x_{I \cap J}, x_{J \setminus I})] \\ &= \mathbb{E}_{x_{I \cap J}} [\mathbb{E}_{x_{I \setminus J}} [f(x_{I \cap J}, x_{I \setminus J})] \cdot \mathbb{E}_{x_{J \setminus I}} [g(x_{I \cap J}, x_{J \setminus I})]] \\ &= \mathbb{E}_{x_{I \cap J}} [f^{\subseteq I \cap J}(x_{I \cap J}) \cdot g^{\subseteq I \cap J}(x_{I \cap J})] \\ &= \langle f^{\subseteq I \cap J}, g^{\subseteq I \cap J} \rangle. \end{aligned}$$

The first line follows from Adam's law and independence of  $x_{I \setminus J}$  and  $x_{J \setminus I}$ , while the second follows from definition of  $f^{\subseteq I \cap J}$  and  $g^{\subseteq I \cap J}$ .  $\square$

Theorem 3.1.4 (O'Donnell, Theorem 8.35): Let  $f \in L^2(\mathbb{R}^N, \pi^{\otimes N})$ . Then  $f$  has a unique decomposition as

$$f = \sum_{S \subseteq [n]} f^{\neg S}$$

where the functions  $f^{\neg S} \in L^2(\mathbb{R}^N, \pi^{\otimes N})$  satisfy

1.  $f^{\neg S}$  depends only on the coordinates in  $S$ ;
2. if  $T \subsetneq S$  and  $g \in L^2(\mathbb{R}^N, \pi^{\otimes N})$  only depends on coordinates in  $T$ , then  $\langle f^{\neg S}, g \rangle = 0$ .

In addition, this decomposition has the following properties:

3. Condition 2. holds whenever  $S \not\subseteq T$ .
4. The decomposition is orthogonal:  $\langle f^{\neg S}, f^{\neg T} \rangle = 0$  for  $S \neq T$ .
5.  $\sum_{S \subseteq T} f^{\neg S} = f^{\neg T}$ .
6. For each  $S \subseteq [n]$ ,  $f \mapsto f^{\neg S}$  is a linear operator.

Definition 3.1.5: The *Efron-Stein degree* of a function  $f \in L^2(\mathbb{R}^N, \pi^{\otimes N})$  is

$$\deg_{\text{ES}}(f) = \max_{S \subseteq [n] \text{ s.t. } f^{\neg S} \neq 0} |S|.$$

If  $f = (f_1, \dots, f_M) : \mathbb{R}^N \rightarrow \mathbb{R}^M$  is a multivariate function, then the Efron-Stein degree of  $f$  is the maximum degree of the  $f_i$ .

Intuitively, the Efron-Stein degree is the maximum size of multiway interactions that  $f$  accounts for.

Motivation for “noise operator” - see how function behaves for small change in input parameters.

Definition 3.1.6: For  $p \in [0, 1]$ , and  $x \in \mathbb{R}^N$ , we say  $y \in \mathbb{R}^N$  is *p-resampled from x* if  $y$  is chosen as follows: for each  $i \in [n]$ , independently,

$$y_i = \begin{cases} x_i & \text{with probability } p \\ \text{drawn from } \pi & \text{with probability } 1 - p \end{cases}$$

We say  $(x, y)$  is a *p-resampled pair*.

Def. noise operator.

Definition 3.1.7: For  $p \in [0, 1]$ , the *noise operator* is the linear operator  $T_p$  on  $L^2(\mathbb{R}^N, \pi^{\otimes N})$ , defined by, for  $y$   $p$ -resampled from  $x$

$$T_p f(x) = \mathbb{E}_{y \text{ } p\text{-resampled from } x} [f(y)]$$

In particular,  $\langle f, T_p f \rangle = \mathbb{E}_{(x,y) \text{ } p\text{-resampled}} [f(x) \cdot f(y)]$ .

Lemma 3.1.8: Let  $p \in [0, 1]$  and  $f \in L^2(\mathbb{R}^N, \pi^{\otimes N})$  have Efron-Stein decomposition  $f = \sum_{S \subseteq [n]} f^S$ . Then

$$T_p f(x) = \sum_{S \subseteq [n]} p^{|S|} f^S.$$

*Proof:* Let  $J$  denote a  $p$ -random subset of  $[n]$ , i.e. with  $J$  formed by including each  $i \in [n]$  independently with probability  $p$ . By definition,  $T_p f(x) = \mathbb{E}_J [f^{\subseteq J}(x)]$  (i.e. pick a random subset of coordinates to fix, and re-randomize the rest). We know by Theorem 3.1.4 that  $f^{\subseteq J} = \sum_{S \subseteq J} f^S$ , so

$$T_p f(x) = \mathbb{E}_J \left[ \sum_{S \subseteq J} f^S \right] = \sum_{S \subseteq [n]} \mathbb{E}_J [I(S \subseteq J)] \cdot f^S = \sum_{S \subseteq [n]} p^{|S|} f^S,$$

since for a fixed  $S \subseteq [n]$ , the probability that  $S \subseteq J$  is  $p^{|S|}$ . □

Lem. Noise operator formula in E-S decomposition. (Ex. 8.18)

Thrm. Function stability for degree  $D$  functions

Theorem 3.1.9: Let  $p \in [0, 1]$  and let  $f = (f_1, \dots, f_M) : \mathbb{R}^N \rightarrow \mathbb{R}^M$  be a multivariate function with Efron-Stein degree  $D$  and each  $f_i \in L^2(\mathbb{R}^N, \pi^{\otimes N})$ . Suppose that  $(x, y)$  are a  $p$ -resampled pair under  $\pi^{\otimes N}$ , and  $\mathbb{E} \|f(x)\|^2 = 1$ . Then

$$\mathbb{E} \|f(x) - f(y)\|^2 \leq 2(1 - p^D) \leq 2(1 - p)D \quad (3.3)$$

*Proof:* Observe that

$$\begin{aligned} \mathbb{E} \|f(x) - f(y)\|^2 &= \mathbb{E} \|f(x)\|^2 + \mathbb{E} \|f(y)\|^2 - 2\mathbb{E} \langle f(x), f(y) \rangle \\ &= 2 - 2 \left( \sum_i \mathbb{E} [f_i(x) f_i(y)] \right) \\ &= 2 - 2 \left( \sum_i \langle f_i, T_p f_i \rangle \right). \end{aligned} \quad (3.4)$$

Here, we have for each  $i \in [n]$  that

$$\langle f_i, T_p f_i \rangle = \left\langle \sum_{S \subseteq [n]} f_i^{\neg S}, \sum_{S \subseteq [n]} p^{|S|} f_i^{\neg S} \right\rangle = \sum_{S \subseteq [n]} p^{|S|} \|f_i^{\neg S}\|^2,$$

by Lemma 3.1.8 and orthogonality. Now, as each  $f_i$  has Efron-Stein degree at most  $D$ , the sum above can be taken only over  $S \subseteq [n]$  with  $0 \leq |S| \leq D$ , giving the bound

$$p^D \mathbb{E}[f_i(x)^2] \leq \langle f_i, T_p f_i \rangle = \mathbb{E}[f_i(x) \cdot T_p f_i(x)] \leq \mathbb{E}[f_i(x)^2].$$

Summing up over  $i$ , and using that  $\mathbb{E}\|f(x)\|^2 = 1$ , gives

$$p^D \leq \sum_i \langle f_i, T_p f_i \rangle = \mathbb{E}\langle f(x), f(y) \rangle \leq 1.$$

Finally, we can substitute into (3.4) to get

$$\mathbb{E}\|f(x) - f(y)\|^2 \leq 2 - 2p^D = 2(1 - p^D) \stackrel{1}{\leq} 2(1 - p)D,$$

as desired. □

### 3.2. Hermite Polynomials (TODO)

Disclaimer of “this theory is much more classical, see (ref) for details.”

Def. Gaussian space

**Definition 3.2.1:** Let  $\gamma_N$  be the  $N$ -dimensional standard Normal measure on  $\mathbb{R}^N$ . Then the  $N$ -dimensional Gaussian space is the space  $L^2(\mathbb{R}^N, \gamma_N)$  of  $L^2$  functions of  $N$  i.i.d. standard Normal random variables.

Note that under the usual  $L^2$  inner product,  $\langle f, g \rangle = \mathbb{E}[f \cdot g]$ , Gaussian space is a separable Hilbert space.

To us, the interesting functions in this space are those given by degree  $D$  multivariate polynomials (here “degree” is used in the traditional sense.)

Thrm. monomials form basis of 1D Gaussian space (cite)

It is a well-known fact that the monomials  $1, z, z^2, \dots$  form a complete basis for  $L^2(\mathbb{R}, \gamma)$  (O’Donnell 11.22). However, these are far from an orthonormal “Fourier” basis; for instance, we know  $\mathbb{E}[z^2] = 1$  for  $z \sim \mathcal{N}(0, 1)$ . By the Gram-Schmidt process, these monomials can be converted into the polynomials  $h_j$  for  $j \geq 0$ , given as

$$h_0(z) = 1, \quad h_1(z) = z, \quad h_2(z) = \frac{z^2 - 1}{\sqrt{2}}, \quad h_3(z) = \frac{z^3 - 3z}{\sqrt{6}}, \quad \dots \quad (3.5)$$

---

<sup>1</sup>This follows from the identity  $(1 - p^D) = (1 - p)(1 + p + p^2 + \dots + p^{D-1})$ ; the bound is tight for  $p \approx 1$ .

Note here that each  $h_j$  is a degree  $j$  polynomial.

and thus the collection of  $(h_j)_{0 \leq j \leq k}$

These formulas require knowledge of the moments of a standard Normal random variable, so a more convenient way to derive them is by analyzing the standard Normal moment generating function. Recall that for  $z \sim \mathcal{N}(0, 1)$ , we have

$$\mathbb{E}[\exp(tz)] = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{tz - \frac{1}{2}z^2} dz = e^{\frac{1}{2}t^2} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-\frac{1}{2}(z-t)^2} dz = \exp\left(\frac{1}{2}t^2\right).$$

**Theorem 3.2.2 (O'Donnell, meow):** The Hermite polynomials  $(h_j)_{j \geq 0}$  form a complete orthonormal basis for  $L^2(\mathbb{R}, \gamma)$ .

To extend this to  $L^2(\mathbb{R}^N, \gamma^N)$ , we can take products. For a multi-index  $\alpha \in \mathbb{N}^N$ , we define the multivariate Hermite polynomial  $h_\alpha : \mathbb{R}^N \rightarrow \mathbb{R}$  as

$$h_\alpha(z) := \prod_{j=1}^N h_{\alpha_j}(z_j).$$

The degree of  $h_\alpha$  is clearly  $|\alpha| = \sum_j \alpha_j$ .

**Theorem 3.2.3:** The Hermite polynomials  $(h_\alpha)_{\alpha \in \mathbb{N}^N}$  form a complete orthonormal basis for  $L^2(\mathbb{R}^N, \gamma^N)$ . In particular, every  $f \in L^2(\mathbb{R}^N, \gamma^N)$  has a unique expansion in  $L^2$  norm as

$$f(z) = \sum_{\alpha \in \mathbb{N}^N} \hat{f}(\alpha) h_\alpha(z).$$

As a consequence of the uniqueness of the expansion in , we see that polynomials are their own Hermite expansion. Namely, let  $H^{\leq k} \subseteq L^2(\mathbb{R}^N, \gamma^N)$  be the subset of multivariate polynomials of degree at most  $k$ . Then, any  $f \in H^{\leq k}$  can be Hermite expanded as

$$f(z) = \sum_{\alpha \in \mathbb{N}^N} \hat{f}(\alpha) h_\alpha(z) = \sum_{|\alpha| \leq k} \hat{f}(\alpha) h_\alpha(z).$$

Thus,  $H^{\leq k}$  is the closed linear span of the set  $\{h_\alpha : |\alpha| \leq k\}$ .

Def. noise operator/Ornstein-Uhlenbeck operator

Compute effect of noise operator on Hermite polys

Thrm. Hermite polys form basis for 1D Gaussian space

Thrm. Products of Hermite polys form basis for N-dim Gaussian space

## Noise operator on arbitrary function with given Hermite expansion

Definition 3.2.4: Let  $(g, g')$  be  $N$ -dimensional standard Normal vectors. We say  $(g, g')$  are  $p$ -correlated if  $(g_i, g'_i)$  are  $p$ -correlated for each  $i \in [n]$ , and these pairs are mutually independent.

In a similar way to the Efron-Stein case, we can consider the resulting “noise operator,” as a way of measuring the effect on a function of a small change in the input.

Definition 3.2.5: For  $p \in [0, 1]$ , the *Gaussian noise operator*  $T_p$  is the linear operator on  $L^2(\mathbb{R}^N, \gamma^N)$ , given by

$$T_p f(x) = \mathbb{E}_{y \text{ } p\text{-correlated to } x} [f(y)] = \mathbb{E}_{y \sim \mathcal{N}(0, I_N)} \left[ f\left(px + \sqrt{1-p^2}y\right) \right]$$

In particular, a straightforward computation with the Normal moment generating function gives

Remark that degree  $D$  function can be expressed in terms of degree  $D$  and lower Hermite polynomials - gives a Hilbert basis which reflects the natural algebraic grading.

Thrm. Function stability for degree  $D$  polynomials.

### 3.3. Algorithms

Def. Randomized algorithm

Def. degree of algorithm is degree as multivariate function.

Discussion of how low-degree algs are approximate for class of Lipschitz algorithms?

Need for rounding function to land on  $\Sigma_N$

Construction of randomized rounding function.

Constr. rounded algorithm.

Lemma. stability of rounding

Thrm. Stability of randomized algorithms (part 1 of Prop 1.9)

Show that Markov gives a useful bound on

Lemma 3.3.1: Let  $f : \mathbb{R}^N \rightarrow \mathbb{R}^N$ ,  $p \in [0, 1]$ , and  $X, Y$  marginally  $N$ -dimensional standard Normal vectors. Suppose that  $\mathbb{E}\|f(X)\|_2^2 = 1$  and either of the following cases hold:

- I.  $(X, Y)$  are a  $p$ -resampled pair, and  $f$  is a degree- $D$  function.
- II.  $(X, Y)$  are  $p$ -correlated, and  $f$  is a degree- $D$  polynomial.

Then

$$\mathbb{E}\|f(X) - f(Y)\|_2^2 \leq 2(1 - p^D).$$



### 3.4. Algorithms

**Definition 3.4.1:** A *randomized algorithm* is a measurable function  $\mathcal{A}^\circ : (g, \omega) \mapsto x \in \mathbb{R}^N$ , where  $\omega \in \Omega_N$  is an independent random variable in some Polish space. Such an  $\mathcal{A}^\circ$  is *deterministic* if it does not depend on  $\omega$ .

**Example 3.4.1:** Let  $\mathbf{U} = (U_1, \dots, U_N)$  be i.i.d.  $\text{Unif}([-1, 1])$ . Then, we define the random coordinate-wise function

$$\text{round}_{\mathbf{U}}(\mathbf{x}) = (\text{round}_{U_1}(x_1), \dots, \text{round}_{U_N}(x_N)),$$

where we define

$$\text{round}_U(x) = \begin{cases} 1 & x \geq U \\ -1 & x < U \end{cases}.$$

**Example 3.4.2:** Given a real-valued algorithm  $\mathcal{A}^\circ$ , we can convert it into a  $\Sigma_N$ -valued algorithm  $\mathcal{A}$  via

$$\mathcal{A}(g, \omega, \mathbf{U}) := \text{round}_{\mathbf{U}}(\mathcal{A}^\circ(g, \omega)).$$

**Definition 3.4.2:** Algorithm  $\mathcal{A}$  is  $(\varepsilon, \eta, p_{\text{unstable}})$ -stable if, for  $g, g'$   $(1 - \varepsilon)$ -correlated/resampled, we have

$$\mathbb{P}(\|\mathcal{A}(g) - \mathcal{A}(g')\| \leq \eta\sqrt{N}) \geq 1 - p_{\text{unstable}}.$$

By the will of God (i.e. writeup pending), we have the following:

**Lemma 3.4.3:** Algorithm  $\mathcal{A}$  with degree  $\leq D$  and norm  $\mathbb{E}\|\mathcal{A}(g)\|^2 \leq CN$  has

$$\mathbb{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^2 \leq 2CN\varepsilon D,$$

and (because of randomized rounding)

$$\mathbb{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^4 \leq 16CN^2\varepsilon D.$$

Thus,

$$\mathbb{P}(\|\mathcal{A}(g) - \mathcal{A}(g')\| \geq \eta\sqrt{N}) \leq \frac{16CN^2\varepsilon D}{\eta^4 N^2} \asymp \frac{\varepsilon D}{\eta^4}.$$

As a consequence, a degree  $D$  algorithm  $\mathcal{A}$  has  $p_{\text{unstable}} = o_{N(1)} \text{ for } \eta^4 \gg \varepsilon D$ .

## 4. Summary of Parameters

Parameter	Meaning	Desired Direction	Intuition
$N$	Dimension	Large	Showing hardness <i>asymptotically</i> , want “bad behavior” to pop up in low dimensions.
$\delta$	Solution tightness; want to find $x$ such that $ \langle g, x \rangle  \leq 2^{-\delta N}$	Small	Smaller $\delta$ implies weaker solutions, e.g. $\delta = 0$ is just finding solutions $\leq 1$ . Want to show even weak solutions are hard to find.
$E$	Solution tightness; “energy level”; want to find $x$ such that $ \langle g, x \rangle  \leq 2^{-E}$	Small	Smaller $E$ implies weaker solutions, and can consider full range of $1 \ll E \ll N$ . Know that $E > (\log^2 N)$ by Karmarkar-Karp
$D$	Algorithm degree (in either Efron-Stein sense or usual polynomial sense.)	Large	Higher degree means more complexity. Want to show even complex algorithms fail.
$\varepsilon$	Complement of correlation/resample probability; $(g, g')$ are $(1 - \varepsilon)$ -correlated.	Small	$\varepsilon$ is “distance” between $g, g'$ . Want to show that small changes in disorder lead to “breaking” of landscape.
$\eta$	Algorithm instability; $\mathcal{A}$ is stable if $\ \mathcal{A}(g) - \mathcal{A}(g')\  \leq \eta\sqrt{N}$ , for $(g, g')$ $(1 - \varepsilon)$ -correlated.	Large	Large $\eta$ indicates a more unstable algorithm; want to show that even weakly stable algorithms fail.

Table 1: Explanation of Parameters

## 5. Conditional Landscape Obstruction

We start with a bound on the geometry of the binary hypercube.

Lemma 5.1: Fix  $x \in \Sigma_N$ . Then, the number of  $x'$  within distance  $\eta\sqrt{N}$  of  $x$  is

$$\left| \left\{ x' \in \Sigma_N \mid \|x - x'\| \leq \eta\sqrt{N} \right\} \right| = \exp_2 \left( Nh \left( \frac{\eta^2}{4} \right) + O(\log_2 N) \right),$$

where  $h(x) = x \log_2 \left( \frac{1}{x} \right) + (1 - x) \log_2 \left( \frac{1}{1-x} \right)$  is the binary entropy function.

*Proof:* Let  $k$  be the number of coordinates which differ between  $x$  and  $x'$  (i.e. the Hamming distance). Then we have  $\|x - x'\|^2 = 4k$ , so for  $\|x - x'\| \leq \eta\sqrt{N}$  requires that  $k \leq N \frac{\eta^2}{4}$ . Given  $k$ , there are  $\binom{N}{k}$  choices for the coordinates of  $x$  to flip, giving a count of

$$\sum_{k \leq \frac{N\eta^2}{4}} \binom{N}{k} = \sum_{k \leq \frac{N\eta^2}{4}} \exp_2 \left( Nh \left( \frac{k}{N} \right) + O(\log_2 N) \right) \leq \exp_2 \left( Nh \left( \frac{\eta^2}{4} \right) + O(\log_2 N) \right).$$

Here, the first equality follows from the Stirling approximation for  $\binom{N}{k}$ , and the second follows because the sum is over  $O(N)$  terms. Moreover, for  $k \leq \frac{N}{2}$ ,  $\frac{k}{N} \leq \frac{\eta^2}{4}$  over the range of this sum.  $\square$

**Lemma 5.2:** Suppose that  $K \leq N/2$ , and let  $h(x) = -x \log(x) - (1-x) \log(x)$  be the binary entropy function. Then, for  $p := K/N$ ,

$$\sum_{k \leq K} \binom{N}{k} \leq \exp(Nh(p)) \leq \exp \left( 2Np \log \left( \frac{1}{p} \right) \right).$$

*Proof:* Consider a  $\text{Bin}(N, p)$  random variable  $S$ . Summing its PMF from 0 to  $K$ , we have

$$1 \geq \mathbb{P}(S \leq K) = \sum_{k \leq K} \binom{N}{k} p^k (1-p)^{N-k} \geq \sum_{k \leq K} \binom{N}{k} p^K (1-p)^{N-K}.$$

Here, the last inequality follows from the fact that  $p \leq (1-p)$ , and we multiply each term by  $\left(\frac{p}{1-p}\right)^{K-k} < 1$ . Now rearrange to get

$$\begin{aligned} \sum_{k \leq K} \binom{N}{k} &\leq p^{-K} (1-p)^{-(N-K)} \\ &= \exp(-K \log(p) - (N-K) \log(1-p)) \\ &= \exp \left( N \cdot \left( -\frac{K}{N} \log(p) - \left( \frac{N-K}{N} \right) \log(1-p) \right) \right) \\ &= \exp(N(-p \log(p) - (1-p) \log(p))) = \exp(Nh(p)). \end{aligned}$$

The final equality then follows from the bound  $h(p) \leq 2p \log(1/p)$  for  $p \leq \frac{1}{2}$ .  $\square$

**Lemma 5.3:** Fix  $x \in \Sigma_N$ , and let  $\eta \leq \frac{1}{\sqrt{2}}$ . Then the number of  $x'$  within distance  $2\eta\sqrt{N}$  of  $x$  is

$$\left| \{x' \in \Sigma_N \mid \|x - x'\| \leq 2\eta\sqrt{N}\} \right| \leq \exp_2(2\eta^2 \log_2(1/\eta^2)N)$$

*Proof:* Let  $k$  be the number of coordinates which differ between  $x$  and  $x'$  (i.e. the Hamming distance). We have  $\|x - x'\|^2 = 4k$ , so  $\|x - x'\| \leq 2\eta\sqrt{N}$  iff  $k \leq N\eta^2$ . Moreover, for  $\eta \leq \frac{1}{\sqrt{2}}$ ,  $k \leq \frac{N}{2}$ . Thus, by Lemma 5.2, we get

$$\sum_{k \leq N\eta^2} \binom{N}{k} \leq \exp_2(Nh(\eta^2)) \leq \exp_2 \left( 2\eta^2 \log_2 \left( \frac{1}{\eta^2} \right) N \right)$$

$\square$

Next, we can consider what this probability is in the case of correlated Normals.

**Lemma 5.4:** Suppose  $(g, g')$  are  $(1 - \varepsilon)$ -correlated Normal vectors, and let  $x \in \Sigma_N$ . Then

$$\mathbb{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g) \leq 2^{-E+O(\log_2 \varepsilon N)}.$$

*Proof:* Let  $\tilde{g}$  be an independent Normal vector to  $g$ , and observe that  $g'$  can be represented as  $g' = pg + \sqrt{1 - p^2}\tilde{g}$ , for  $p = 1 - \varepsilon$ . Thus,  $\langle g', x \rangle = p\langle g, x \rangle + \sqrt{1 - p^2}\langle \tilde{g}, x \rangle$ . Observe  $\langle g, x \rangle$  is constant given  $g$ , and  $\langle \tilde{g}, x \rangle$  is a Normal r.v. with mean 0 and variance  $N$ , so  $\langle g', x \rangle \sim \mathcal{N}(p\langle g, x \rangle, (1 - p^2)N)$ . This random variable is nondegenerate for  $(1 - p^2)N > 0$ , with density bounded above as

$$\varphi_g(z) = \frac{1}{\sqrt{2\pi(1 - p^2)N}} e^{-\frac{(z - p\langle g, x \rangle)^2}{2(1 - p^2)N}} \leq \frac{1}{\sqrt{2\pi(1 - p^2)N}}.$$

Following the remainder of the proof of Lemma 5.5, we conclude that

$$\mathbb{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g) \leq \sqrt{\frac{2}{\pi(1 - p^2)N}} 2^{-E} = 2^{-E+O(\log_2(1 - p^2)N)} = 2^{-E+O(\log_2 \varepsilon N)}.$$

The last line follows as  $(1 - p^2)N \leq 2(1 - p)N = 2\varepsilon N$ .  $\square$

Note for instance that here  $\varepsilon$  can be exponentially small in  $E$  (i.e.  $\varepsilon = \exp_2(-E/10)$ ), which for the case  $E = \Theta(N)$  implies  $\varepsilon$  can be exponentially small in  $N$ .

First, we consider the probability of a solution being optimal for a resampled instance.

**Lemma 5.5:** Suppose  $(g, g')$  are  $(1 - \varepsilon)$ -resampled Normal vectors, and let  $x \in \Sigma_N$ . Then,

$$\mathbb{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g, g \neq g') \leq 2^{-M+O(1)}.$$

*Proof:* Let  $S = \{i \in [N] : g_i \neq g'_i\}$  be the set of indices where  $g$  and  $g'$  differ. We can express

$$\langle g', x' \rangle = \sum_{i \in [N]} g'_i x_i = \sum_{i \notin S} g_i x_i + \sum_{i \in S} g'_i x_i \sim \mathcal{N}\left(\sum_{i \notin S} g_i x_i, |S|\right).$$

Let  $\mu := \sum_{i \notin S} g_i x_i$ . The conditional distribution of interest can now be expressed as  $\mathbb{P}(|\langle g', x' \rangle| \leq 2^{-E} \mid g, |S| \geq 1)$ . Given  $|S| \geq 1$ , the quantity  $\langle g', x' \rangle$  is a nondegenerate random variable, with density bounded above as

$$\varphi_{g, |S|}(z) = \frac{1}{\sqrt{2\pi|S|}} e^{-\frac{(z - \mu)^2}{2|S|}} \leq \frac{1}{\sqrt{2\pi|S|}} \leq \frac{1}{\sqrt{2\pi}}.$$

Hence, the quantity of interest can be bounded as

$$\mathbb{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g, g \neq g') \leq \int_{|z| \leq 2^{-E}} \varphi_{g, |S|}(z) dz \leq \sqrt{\frac{2}{\pi}} 2^{-E} = 2^{-E+O(1)}.$$

□

In this case, we can compute the probability that  $g = g'$  as

$$\mathbb{P}(g = g') = \prod_{i=1}^N \mathbb{P}(g_i = g_{i'}) = (1 - \varepsilon)^N,$$

which for  $\varepsilon \ll 1$  is approximately  $1 - N\varepsilon$ . Thus, for  $\varepsilon \gg \omega(\frac{1}{N})$ , we have

$$\mathbb{P}(|\langle g', x \rangle| \leq 2^{-E} \mid g) \leq 2^{-E+O(1)}$$

### 5.1. Proof of Low-Degree Hardness.

Let  $\delta > 0$ . Let  $E$  be a sequence of energy levels. Assume for sake of contradiction that  $p_{\text{solve}} \geq \Omega(1)$ . Let  $g, g'$  be  $(1 - \varepsilon)$ -resampled/ $(1 - \varepsilon)$ -correlated problem instances. We define the following events:

$$S_{\text{solve}} = \{\mathcal{A}(g) \in S(\delta; g), \mathcal{A}(g') \in S(\delta; g')\}$$

$$S_{\text{stable}} = \{\|\mathcal{A}(g) - \mathcal{A}(g')\| \leq 2\eta\sqrt{N}\}$$

$$S_{\text{ogp}} = \{\text{for } x \text{ depending only on } g, \exists x' \in S(\delta; g') \text{ such that } \|x - x'\| \leq \eta\sqrt{N}\}$$

To set the remaining parameters, choose  $\varepsilon = \omega(\frac{1}{N})$  such that  $\varepsilon D = o(1)$ . Then, choose  $\eta$  such that  $(h^{-1}(\delta))^2 \gg \eta^4 \gg \varepsilon D$ . With this, the previous landscape obstructions give the following.

**Lemma 5.1.1:** For any  $\omega(\log^2 N) \leq E \leq \Theta(N)$ , there exist choices of  $\varepsilon, \eta$  (depending on  $N, E$ ) such that  $\mathbb{P}(S_{\text{ogp}}) = o(1)$ .

*Proof:* Observe that

$$\mathbb{P}(S_{\text{ogp}}) = \mathbb{E}[\mathbb{P}(S_{\text{ogp}} \mid g)]. \quad (5.1)$$

Conditional on  $g$ , we can compute  $\mathbb{P}(S_{\text{ogp}} \mid g) = \mathbb{P}(\exists x' \in S(E; g'), \|x - x'\| \leq 2\eta\sqrt{N})$  by setting  $x = \mathcal{A}(g)$  (so  $x$  only depends on  $g$ ), and union bounding Lemma 5.5 over the  $x'$  within  $2\eta\sqrt{N}$  of  $x$ , as per Lemma 5.1:

$$\mathbb{P}(S_{\text{ogp}} \mid g) \leq \exp_2(-E + N\eta^2 \log_2(1/\eta^2) + O(1)).$$

We want to choose  $\eta$  such that

$$-E + N\eta^2 \log_2(1/\eta^2) = -\Omega(N)$$

$$\frac{E}{N} > \eta^2 \log(1/\eta^2)$$

Using the fact that  $\sqrt{2x} \geq -x \log_2 x$ , it suffices to pick  $\eta^2$  with

$$\frac{E}{N} > 2\eta,$$

so  $\eta^2 := \frac{E^2}{2N^2}$  is a valid choice.

By the choice of  $\eta^2 \ll (h^{-1})(\delta) \asymp 1$ , this bound gives  $\mathbb{P}(S_{\text{ogp}}|g) \leq \exp_2(-O(N)) = o(1)$ .

Integrating over  $g$  gives the overall bound.  $\square$

When  $CD\varepsilon N^2 = \omega_{N(1)}$  (i.e. goes to infinity),

$$\begin{aligned} \mathbb{P}(S_{\text{stable}}) &\leq \frac{16CD\varepsilon N^2}{16\eta^4 N^2} \\ &= \frac{CD\varepsilon}{\eta^4} = \frac{4CD\varepsilon N^4}{E^4} \end{aligned}$$

$D\varepsilon \rightarrow 0$  same as  $D = o(\frac{1}{\varepsilon}) = o(N)$ .

Lemma 5.1.2:  $\mathbb{P}(S_{\text{solve}}, S_{\text{stable}}) \leq \mathbb{P}(S_{\text{ogp}}) = o(1)$ .

*Proof:* The first inequality follows from definition, with  $x = \mathcal{A}(g)$  and  $x' = \mathcal{A}(g')$ . For the second, observe that

$$\mathbb{P}(S_{\text{ogp}}) = \mathbb{E}[\mathbb{P}(S_{\text{ogp}} \mid g)].$$

Now, let  $M = \delta N$ , we can compute  $\mathbb{P}(S_{\text{ogp}} \mid g) = \mathbb{P}(\exists x' \in S(\delta; g'), \|x - x'\| \leq \eta\sqrt{N})$  by setting  $x = \mathcal{A}(g)$  (so  $x$  only depends on  $g$ ), and union bounding Lemma 5.5 over the  $x'$  within  $\eta\sqrt{N}$  of  $x$ , as per Lemma 5.1:

$$\mathbb{P}(S_{\text{ogp}} \mid g) \leq \exp_2\left(-\delta N + Nh\left(\frac{\eta^2}{4}\right) + O(\log_2 N)\right).$$

By the choice of  $\eta^2 \ll (h^{-1})(\delta) \asymp 1$ , this bound gives  $\mathbb{P}(S_{\text{ogp}}|g) \leq \exp_2(-O(N)) = o(1)$ .

Integrating over  $g$  gives the overall bound.  $\square$

However, by the choice of parameters above, we also have

$$\begin{aligned} \mathbb{P}(S_{\text{solve}}, S_{\text{stable}}) &\geq \mathbb{P}(S_{\text{solve}}) + \mathbb{P}(S_{\text{stable}}) - 1 \\ &\geq p_{\text{solve}}^4 + p_{\text{unstable}} \geq \Omega(1) - o(1) = \Omega(1), \end{aligned} \tag{5.2}$$

which is a contradiction.

## 6. Randomized Rounding Things

Claim: no two adjacent points on  $\Sigma_N$  (or pairs within  $k = O(1)$  distance) which are both good solutions to the same problem. The reason is that this would require a subset of  $k$  signed coordinates  $\pm g_{\{i_1\}}, \dots, \pm g_{\{i_k\}}$  to have small sum, and there are only  $2^k \text{ binom}\{N\}\{k\}l = O(N^k)$

possibilities, each of which is centered Gaussian with variance at least 1, so the smallest is typically of order  $\Omega(N^{\{-k\}})$ .

Thus, rounding would destroy the solution.

- Say we're in the case where rounding changes the solution. (i.e. rounding moves  $x$  to point that is not the closest point  $x_*$ , whp.)
- Let  $p_1, \dots, p_N$  be the probabilities of disagreeing with  $x_*$  on each coordinate.
  - We know that  $\sum p_i$  diverges (this is equivalent to the statement that rounding will change the solution whp).
  - Reason: for each coord, chance of staying at that coordinate is  $e^{-\Theta(p_i)}$ .
- For each  $i$ , flip coin with heads prob  $2p_i$ , and keep all the heads.
  - By Borel-Cantelli type argument, typical number of heads will be  $\omega(1)$ .
- For every coin with a head, change coord with prob 50%, if tails, keep coord.
  - Randomized rounding in artificially difficult way. (I.e. this multistage procedure accomplishes the same thing as randomized rounding.)
- Now, randomized rounding is done by choosing a random set of  $\omega(1)$  coordinates, and making those iid Uniform in  $\{-1, 1\}$ .
- Pick a large constant (e.g. 100), and only randomize the first 100 heads, and condition on the others (i.e. choose the others arbitrarily). Note that since  $100l = \omega(1)$ , there are at least 100 heads whp.
- Now rounded point is random point in 100 dimensional subcube, but at most one of them is a good solution by the claim at the top of the page.
- Combining, the probability for rounding to give a good solution is at most  $o(1) + 2^{\{-100\}}$ . Since 100 is arbitrary, this is  $o(1)$  by sending parameters to 0 and/or infinity in the right order.

## Bibliography

- [1] D. Achlioptas and A. Coja-Oghlan, "Algorithmic Barriers from Phase Transitions," in *2008 49th Annual IEEE Symposium on Foundations of Computer Science*, Oct. 2008, pp. 793–802. doi: 10.1109/FOCS.2008.11.
- [2] D. Achlioptas and F. Ricci-Tersenghi, "On the Solution-Space Geometry of Random Constraint Satisfaction Problems." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/cs/0611052>
- [3] L. Addario-Berry, L. Devroye, G. Lugosi, and R. I. Oliveira, "Local Optima of the Sherrington-Kirkpatrick Hamiltonian." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1712.07775>
- [4] B. Alidaee, F. Glover, G. A. Kochenberger, and C. Rego, "A New Modeling and Solution Approach for the Number Partitioning Problem," *Journal of Applied Mathematics and Decision Sciences*, vol. 2005, no. 2, pp. 113–121, Jan. 2005, doi: 10.1155/JAMDS.2005.113.
- [5] M. F. Argüello, T. A. Feo, and O. Goldschmidt, "Randomized Methods for the Number Partitioning Problem," *Computers & Operations Research*, vol. 23, no. 2, pp. 103–111, Feb. 1996, doi: 10.1016/0305-0548(95)E0020-L.

- [6] L. Asproni, D. Caputo, B. Silva, G. Fazzi, and M. Magagnini, “Accuracy and Minor Embedding in Subqubo Decomposition with Fully Connected Large Problems: A Case Study about the Number Partitioning Problem,” *Quantum Machine Intelligence*, vol. 2, no. 1, p. 4, Jun. 2020, doi: 10.1007/s42484-020-00014-w.
- [7] B. Aubin, W. Perkins, and L. Zdeborová, “Storage Capacity in Symmetric Binary Perceptrons,” *Journal of Physics A: Mathematical and Theoretical*, vol. 52, no. 29, p. 294003, Jul. 2019, doi: 10.1088/1751-8121/ab227a.
- [8] A. S. Bandeira, A. Perry, and A. S. Wein, “Notes on Computational-to-Statistical Gaps: Predictions Using Statistical Physics.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1803.11132>
- [9] N. Bansal, “Constructive Algorithms for Discrepancy Minimization.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1002.2259>
- [10] B. Barak, S. B. Hopkins, J. Kelner, P. K. Kothari, A. Moitra, and A. Potechin, “A Nearly Tight Sum-of-Squares Lower Bound for the Planted Clique Problem.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1604.03084>
- [11] H. Bauke, S. Franz, and S. Mertens, “Number Partitioning as a Random Energy Model,” *Journal of Statistical Mechanics: Theory and Experiment*, vol. 2004, no. 4, p. P4003, Apr. 2004, doi: 10.1088/1742-5468/2004/04/P04003.
- [12] M. Bayati, D. Gamarnik, and P. Tetali, “Combinatorial Approach to the Interpolation Method and Scaling Limits in Sparse Random Graphs,” *The Annals of Probability*, vol. 41, no. 6, Nov. 2013, doi: 10.1214/12-AOP816.
- [13] Q. Berthet and P. Rigollet, “Computational Lower Bounds for Sparse PCA.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1304.0828>
- [14] S. Boettcher and S. Mertens, “Analysis of the Karmarkar-Karp Differencing Algorithm,” *The European Physical Journal B*, vol. 65, no. 1, pp. 131–140, Sep. 2008, doi: 10.1140/epjb/e2008-00320-9.
- [15] C. Borgs, J. Chayes, and B. Pittel, “Phase Transition and Finite-size Scaling for the Integer Partitioning Problem,” *Random Structures & Algorithms*, vol. 19, no. 3–4, pp. 247–288, Oct. 2001, doi: 10.1002/rsa.10004.
- [16] M. Brennan and G. Bresler, “Optimal Average-Case Reductions to Sparse PCA: From Weak Assumptions to Strong Hardness.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1902.07380>
- [17] M. Brennan, G. Bresler, and W. Huleihel, “Reducibility and Computational Lower Bounds for Problems with Planted Sparse Structure.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1806.07508>
- [18] K. Chandrasekaran and S. Vempala, “Integer Feasibility of Random Polytopes.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1111.4649>



- [19] W.-K. Chen, D. Gamarnik, D. Panchenko, and M. Rahman, “Suboptimality of Local Algorithms for a Class of Max-Cut Problems,” *The Annals of Probability*, vol. 47, no. 3, May 2019, doi: 10.1214/18-AOP1291.
- [20] E. G. Coffman Jr., M. R. Garey, and D. S. Johnson, “An Application of Bin-Packing to Multiprocessor Scheduling,” *SIAM Journal on Computing*, vol. 7, no. 1, pp. 1–17, Feb. 1978, doi: 10.1137/0207001.
- [21] E. G. Coffman and G. S. Lueker, *Probabilistic Analysis of Packing and Partitioning Algorithms*. in Wiley-Interscience Series in Discrete Mathematics and Optimization. New York: J. Wiley & sons, 1991.
- [22] A. Coja-Oghlan and C. Efthymiou, “On Independent Sets in Random Graphs,” *Random Structures & Algorithms*, vol. 47, no. 3, pp. 436–486, Oct. 2015, doi: 10.1002/rsa.20550.
- [23] D. Corus, P. S. Oliveto, and D. Yazdani, “Artificial Immune Systems Can Find Arbitrarily Good Approximations for the NP-hard Number Partitioning Problem,” *Artificial Intelligence*, vol. 274, pp. 180–196, Sep. 2019, doi: 10.1016/j.artint.2019.03.001.
- [24] I. Cultura, I. Gent, and T. Walsh, “Phase Transitions and Annealed Theories: Number Partitioning as a Case Study,” Jun. 2000.
- [25] Y. Deshpande and A. Montanari, “Improved Sum-of-Squares Lower Bounds for Hidden Clique and Hidden Submatrix Problems.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1502.06590>
- [26] I. Diakonikolas, D. M. Kane, and A. Stewart, “Statistical Query Lower Bounds for Robust Estimation of High-dimensional Gaussians and Gaussian Mixtures.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1611.03473>
- [27] V. Feldman, E. Grigorescu, L. Reyzin, S. Vempala, and Y. Xiao, “Statistical Algorithms and a Lower Bound for Detecting Planted Clique.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1201.1214>
- [28] F. F. Ferreira and J. F. Fontanari, “Probabilistic Analysis of the Number Partitioning Problem,” *Journal of Physics A: Mathematical and General*, vol. 31, no. 15, p. 3417, Apr. 1998, doi: 10.1088/0305-4470/31/15/007.
- [29] D. Gamarnik and E. C. Kızıldağ, “Algorithmic Obstructions in the Random Number Partitioning Problem.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2103.01369>
- [30] D. Gamarnik, E. C. Kızıldağ, W. Perkins, and C. Xu, “Algorithms and Barriers in the Symmetric Binary Perceptron Model.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2203.15667>
- [31] D. Gamarnik and E. Kizildag, “Computing the Partition Function of the Sherrington-Kirkpatrick Model Is Hard on Average,” *The Annals of Applied Probability*, vol. 31, no. 3, Jun. 2021, doi: 10.1214/20-AAP1625.

- [32] D. Gamarnik, A. Jagannath, and A. S. Wein, “Hardness of Random Optimization Problems for Boolean Circuits, Low-Degree Polynomials, and Langevin Dynamics.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2004.12063>
- [33] D. Gamarnik and I. Zadik, “High-Dimensional Regression with Binary Coefficients. Estimating Squared Error and a Phase Transition.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1701.04455>
- [34] D. Gamarnik and I. Zadik, “The Landscape of the Planted Clique Problem: Dense Subgraphs and the Overlap Gap Property.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1904.07174>
- [35] D. Gamarnik and M. Sudan, “Limits of Local Algorithms over Sparse Random Graphs.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/1304.1831>
- [36] D. Gamarnik and A. Jagannath, “The Overlap Gap Property and Approximate Message Passing Algorithms for  $\mathbb{Z}_p$ -Spin Models.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1911.06943>
- [37] D. Gamarnik, “The Overlap Gap Property: A Geometric Barrier to Optimizing over Random Structures,” *Proceedings of the National Academy of Sciences*, vol. 118, no. 41, p. e2108492118, Oct. 2021, doi: 10.1073/pnas.2108492118.
- [38] D. Gamarnik, A. Jagannath, and S. Sen, “The Overlap Gap Property in Principal Submatrix Recovery,” *Probability Theory and Related Fields*, vol. 181, no. 4, pp. 757–814, Dec. 2021, doi: 10.1007/s00440-021-01089-7.
- [39] D. Gamarnik and M. Sudan, “Performance of Sequential Local Algorithms for the Random NAE- $\mathbb{Z}_K$ -SAT Problem,” *SIAM Journal on Computing*, vol. 46, no. 2, pp. 590–619, Jan. 2017, doi: 10.1137/140989728.
- [40] D. Gamarnik and I. Zadik, “Sparse High-Dimensional Linear Regression. Algorithmic Barriers and a Local Search Algorithm.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1711.04952>
- [41] M. R. Garey and D. S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*. in A Series of Books in the Mathematical Sciences. New York: W. H. Freeman, 1979.
- [42] I. P. Gent and T. Walsh, “Analysis of Heuristics for Number Partitioning,” *Computational Intelligence*, vol. 14, no. 3, pp. 430–451, 1998, doi: 10.1111/0824-7935.00069.
- [43] C. Harshaw, F. Sävje, D. Spielman, and P. Zhang, “Balancing Covariates in Randomized Experiments with the Gram-Schmidt Walk Design.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/1911.03071>
- [44] T. Hastie, R. Tibshirani, and J. Friedman, *The Elements of Statistical Learning*. in Springer Series in Statistics. New York, NY: Springer New York, 2009. doi: 10.1007/978-0-387-84858-7.

- [45] H. Hatami, L. Lovász, and B. Szegedy, “Limits of Locally–Globally Convergent Graph Sequences,” *Geometric and Functional Analysis*, vol. 24, no. 1, pp. 269–296, Feb. 2014, doi: 10.1007/s00039-014-0258-7.
- [46] R. Hoberg, H. Ramadas, T. Rothvoss, and X. Yang, “Number Balancing Is as Hard as Minkowski’s Theorem and Shortest Vector.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/1611.08757>
- [47] S. B. Hopkins, P. K. Kothari, A. Potechin, P. Raghavendra, T. Schramm, and D. Steurer, “The Power of Sum-of-Squares for Detecting Hidden Structures.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1710.05017>
- [48] S. Hopkins, “Statistical Inference and the Sum of Squares Method,” 2018.
- [49] S. B. Hopkins, J. Shi, and D. Steurer, “Tensor Principal Component Analysis via Sum-of-Squares Proofs.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1507.03269>
- [50] B. Huang and M. Sellke, “Strong Low Degree Hardness for Stable Local Optima in Spin Glasses.” Accessed: Jan. 30, 2025. [Online]. Available: <http://arxiv.org/abs/2501.06427>
- [51] M. Jerrum, “Large Cliques Elude the Metropolis Process,” *Random Structures & Algorithms*, vol. 3, no. 4, pp. 347–359, Jan. 1992, doi: 10.1002/rsa.3240030402.
- [52] D. S. Johnson, C. R. Aragon, L. A. McGeoch, and C. Schevon, “Optimization by Simulated Annealing: An Experimental Evaluation; Part I, Graph Partitioning,” *Operations Research*, vol. 37, no. 6, pp. 865–892, 1989, Accessed: Mar. 15, 2025. [Online]. Available: <http://www.jstor.org/stable/171470>
- [53] D. S. Johnson, C. R. Aragon, L. A. McGeoch, and C. Schevon, “Optimization by Simulated Annealing: An Experimental Evaluation; Part II, Graph Coloring and Number Partitioning,” *Operations Research*, vol. 39, no. 3, pp. 378–406, 1991, Accessed: Mar. 15, 2025. [Online]. Available: <http://www.jstor.org/stable/171393>
- [54] N. Karmarkar, R. M. Karp, G. S. Lueker, and A. M. Odlyzko, “Probabilistic Analysis of Optimum Partitioning,” *Journal of Applied Probability*, vol. 23, no. 3, pp. 626–645, 1986, doi: 10.2307/3214002.
- [55] N. Karmarkar and R. M. Karp, “The Differencing Method of Set Partitioning,” 1983. Accessed: Mar. 15, 2025. [Online]. Available: <https://www2.eecs.berkeley.edu/Pubs/TechRpts/1983/6353.html>
- [56] M. Kearns, “Efficient Noise-Tolerant Learning from Statistical Queries,” *Journal of the ACM*, vol. 45, no. 6, pp. 983–1006, Nov. 1998, doi: 10.1145/293347.293351.
- [57] E. C. Kızıldağ, “Planted Random Number Partitioning Problem.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2309.15115>
- [58] J. Kojić, “Integer Linear Programming Model for Multidimensional Two-Way Number Partitioning Problem,” *Computers & Mathematics with Applications*, vol. 60, no. 8, pp. 2302–2308, Oct. 2010, doi: 10.1016/j.camwa.2010.08.024.

- [59] R. E. Korf, "From Approximate to Optimal Solutions: A Case Study of Number Partitioning," in *Proceedings of the 14th International Joint Conference on Artificial Intelligence - Volume 1*, in IJCAI'95. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc., Aug. 1995, pp. 266–272.
- [60] R. E. Korf, "A Complete Anytime Algorithm for Number Partitioning," *Artificial Intelligence*, vol. 106, no. 2, pp. 181–203, Dec. 1998, doi: 10.1016/S0004-3702(98)00086-1.
- [61] R. E. Korf, "Multi-Way Number Partitioning," in *Proceedings of the 21st International Joint Conference on Artificial Intelligence*, in IJCAI'09. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc., Jul. 2009, pp. 538–543.
- [62] P. K. Kothari, R. Mori, R. O'Donnell, and D. Witmer, "Sum of Squares Lower Bounds for Refuting Any CSP." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1701.04521>
- [63] J. Kratica, J. Kojić, and A. Savić, "Two Metaheuristic Approaches for Solving Multidimensional Two-Way Number Partitioning Problem," *Computers & Operations Research*, vol. 46, pp. 59–68, Jun. 2014, doi: 10.1016/j.cor.2014.01.003.
- [64] A. M. Krieger, D. Azriel, and A. Kapelner, "Nearly Random Designs with Greatly Improved Balance," *Biometrika*, vol. 106, no. 3, pp. 695–701, Sep. 2019, doi: 10.1093/biomet/asz026.
- [65] D. Kunisky, A. S. Wein, and A. S. Bandeira, "Notes on Computational Hardness of Hypothesis Testing: Predictions Using the Low-Degree Likelihood Ratio." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1907.11636>
- [66] J. Lauer and N. Wormald, "Large Independent Sets in Regular Graphs of Large Girth," *Journal of Combinatorial Theory, Series B*, vol. 97, no. 6, pp. 999–1009, Nov. 2007, doi: 10.1016/j.jctb.2007.02.006.
- [67] A. Levy, H. Ramadas, and T. Rothvoss, "Deterministic Discrepancy Minimization via the Multiplicative Weight Update Method." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1611.08752>
- [68] S. Lovett and R. Meka, "Constructive Discrepancy Minimization by Walking on The Edges." Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1203.5747>
- [69] G. S. Lueker, "A Note on the Average-Case Behavior of a Simple Differencing Method for Partitioning," *Operations Research Letters*, vol. 6, no. 6, pp. 285–287, Dec. 1987, doi: 10.1016/0167-6377(87)90044-7.
- [70] R. Meka, A. Potechin, and A. Wigderson, "Sum-of-Squares Lower Bounds for Planted Clique," in *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing*, Portland Oregon USA: ACM, Jun. 2015, pp. 87–96. doi: 10.1145/2746539.2746600.
- [71] R. Merkle and M. Hellman, "Hiding Information and Signatures in Trapdoor Knapsacks," *IEEE Transactions on Information Theory*, vol. 24, no. 5, pp. 525–530, Sep. 1978, doi: 10.1109/TIT.1978.1055927.
- [72] S. Mertens, "The Easiest Hard Problem: Number Partitioning." Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/cond-mat/0310317>

- [73] S. Mertens, “A Physicist's Approach to Number Partitioning,” *Theoretical Computer Science*, vol. 265, no. 1, pp. 79–108, Aug. 2001, doi: 10.1016/S0304-3975(01)00153-0.
- [74] M. Mézard, T. Mora, and R. Zecchina, “Clustering of Solutions in the Random Satisfiability Problem,” *Physical Review Letters*, vol. 94, no. 19, p. 197205, May 2005, doi: 10.1103/PhysRevLett.94.197205.
- [75] W. Michiels, J. Korst, E. Aarts, and J. Van Leeuwen, “Performance Ratios for the Differencing Method Applied to the Balanced Number Partitioning Problem,” *STACS 2003*, vol. 2607. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 583–595, 2003. doi: 10.1007/3-540-36494-3\_51.
- [76] R. O'Donnell, “Analysis of Boolean Functions.” Accessed: Mar. 15, 2025. [Online]. Available: <http://arxiv.org/abs/2105.10386>
- [77] P. Raghavendra, T. Schramm, and D. Steurer, “High-Dimensional Estimation via Sum-of-Squares Proofs.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1807.11419>
- [78] M. Rahman and B. Virag, “Local Algorithms for Independent Sets Are Half-Optimal,” *The Annals of Probability*, vol. 45, no. 3, May 2017, doi: 10.1214/16-AOP1094.
- [79] T. Rothvoss, “Constructive Discrepancy Minimization for Convex Sets.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1404.0339>
- [80] V. Santucci, M. Baioletti, and G. Di Bari, “An Improved Memetic Algebraic Differential Evolution for Solving the Multidimensional Two-Way Number Partitioning Problem,” *Expert Systems with Applications*, vol. 178, p. 114938, Sep. 2021, doi: 10.1016/j.eswa.2021.114938.
- [81] R. H. Storer, S. W. Flanders, and S. David Wu, “Problem Space Local Search for Number Partitioning,” *Annals of Operations Research*, vol. 63, no. 4, pp. 463–487, Aug. 1996, doi: 10.1007/BF02156630.
- [82] L.-H. Tsai, “Asymptotic Analysis of an Algorithm for Balanced Parallel Processor Scheduling,” *SIAM Journal on Computing*, vol. 21, no. 1, pp. 59–64, Feb. 1992, doi: 10.1137/0221007.
- [83] P. Turner, R. Meka, and P. Rigollet, “Balancing Gaussian Vectors in High Dimension.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/1910.13972>
- [84] M. J. Wainwright, *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. in Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press, 2019. doi: 10.1017/9781108627771.
- [85] A. S. Wein, “Optimal Low-Degree Hardness of Maximum Independent Set.” Accessed: Mar. 16, 2025. [Online]. Available: <http://arxiv.org/abs/2010.06563>
- [86] J. Wen *et al.*, “Optical Experimental Solution for the Multiway Number Partitioning Problem and Its Application to Computing Power Scheduling,” *Science China Physics, Mechanics & Astronomy*, vol. 66, no. 9, p. 290313, Sep. 2023, doi: 10.1007/s11433-023-2147-3.

- [87] B. Yakir, “The Differencing Algorithm LDM for Partitioning: A Proof of a Conjecture of Karmarkar and Karp,” *Mathematics of Operations Research*, vol. 21, no. 1, pp. 85–99, Feb. 1996, doi: 10.1287/moor.21.1.85.
- [88] L. Zdeborová and F. Krzakala, “Statistical Physics of Inference: Thresholds and Algorithms,” *Advances in Physics*, vol. 65, no. 5, pp. 453–552, Sep. 2016, doi: 10.1080/00018732.2016.1211393.