

SLDH For NPP

Strong Low Degree Hardness for the Number Partitioning Problem

Rushil Mallarapu¹

Harvard University, Cambridge, MA 02138

Abstract. Meow Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat voluptatem. Ut enim aequale doleamus animo, cum corpore dolemus, fieri tamen permagna accessio potest, si aliquod aeternum et infinitum impendere malum nobis opinemur. Quod idem licet transferre in voluptatem, ut postea variari voluptas distinguere possit, augeri amplificarique non possit. At etiam Athenis, ut e patre audiebam facete et urbane Stoicos irridente, statua est in quo a nobis philosophia defensa et collaudata est, cum id, quod maxime placeat, facere possimus, omnis voluptas assumenda est, omnis dolor repellendus. Temporibus autem quibusdam et aut officiis debitis aut rerum necessitatibus saepe eveniet, ut et voluptates repudiandae sint et molestiae non recusandae. Itaque earum rerum defuturum, quas natura non depravata desiderat. Et quem ad me accedis, saluto: 'chaere,' inquam, 'Titel' lictores, turma omnis chorusque: 'chaere, Titel' hinc hostis mi Albucius, hinc inimicus. Sed iure Mucius. Ego autem mirari satis non queo unde hoc sit tam insolens domesticarum rerum fastidium. Non est omnino hic docendi locus; sed ita prorsus existimo, neque eum Torquatum, qui hoc primus cognomen invenerit, aut torquem illum hosti detraxisse, ut aliquam ex eo est consecutus? – Laudem et caritatem, quae sunt vitae.

¹Written under the joint supervision of Professor Mark Sellke and Professor Subhabrata Sen.

Acknowledgments

Meow Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat voluptatem. Ut enim aequale doleamus animo, cum corpore dolemus, fieri tamen permagna accessio potest, si aliquod aeternum et infinitum impendere malum nobis opinemur. Quod idem licet transferre in voluptatem, ut postea variari voluptas distinguique possit, augeri amplificarique non possit. At etiam Athenis, ut e patre audiebam facete et urbane Stoicos irridente, statua est in quo a nobis philosophia defensa et collaudata est, cum id, quod maxime placeat, facere possimus, omnis voluptas assumenda est, omnis dolor repellendus. Temporibus autem quibusdam et aut officiis debitis aut rerum necessitatibus saepe eveniet, ut et voluptates repudiandae sint et molestiae non recusandae. Itaque earum rerum defuturum, quas natura non depravata desiderat. Et quem ad me accedis, saluto: 'chaere,' inquam, 'Tite!' lictores, turma omnis chorusque: 'chaere, Tite!' hinc hostis mi Albucius, hinc inimicus. Sed iure Mucius. Ego autem mirari satis non queo unde hoc sit tam insolens domesticarum rerum fastidium. Non est omnino hic docendi locus; sed ita prorsus existimo, neque eum Torquatum, qui hoc primus cognomen invenerit, aut torquem illum hosti detraxisse, ut aliquam ex eo est consecutus? – Laudem et caritatem, quae sunt vitae sine metu degendae praesidia firmissima. – Filium morte multavit. – Si sine causa, nollem me ab eo delectari, quod ista Platonis, Aristoteli, Theophrasti orationis ornamenta neglexerit. Nam illud quidem physici, credere aliquid esse minimum, quod profecto numquam putavisset, si a Polyaeno, familiari suo, geometrica discere maluisset quam illum etiam ipsum dedocere. Sol Democrito magnus videtur, quippe homini erudito in geometriaque perfecto, huic pedalis fortasse; tantum enim esse omnino in nostris poetis aut inertissimae segnitiae est aut fastidii delicatissimi. Mihi quidem videtur, inermis ac nudus est. Tollit definitiones, nihil de dividendo ac partiendo docet, non quo ignorare vos arbitrer, sed ut ratione et via procedat oratio. Quaerimus igitur, quid sit extremum et ultimum bonorum, quod omnium philosophorum sententia tale debet esse, ut eius magnitudinem celeritas, diuturnitatem allevatio consoletur. Ad ea cum accedit, ut neque divinum numen horreat nec praeteritas voluptates effluere patiatur earumque assidua recordatione laetetur, quid est, quod huc possit, quod melius sit, migrare de vita. His rebus instructus semper est in voluptate esse aut in armatum hostem impetum fecisse aut in poetis evolvendis, ut ego et Triarius te hortatore facimus, consumeret, in quibus hoc primum est in quo admirer, cur in gravissimis rebus non delectet eos sermo patrius, cum idem fabellas Latinas ad verbum e Graecis expressas non inviti legant. Quis enim tam inimicus paene nomini Romano est, qui Ennii Medeam aut Antiopam Pacuvii spernat aut reiciat, quod se isdem Euripidis fabulis delectari dicat, Latinas litteras oderit? Synephebos ego, inquit, potius Caecillii aut Andriam Terentii quam utramque Menandri legam? A quibus tantum dissentio, ut, cum Sophocles vel optime scripserit Electram, tamen male conversam Atilii mihi legendam putem, de quo Lucilius: 'ferreum scriptorem', verum, opinor, scriptorem tamen, ut legendus sit. Rudem enim esse omnino in nostris poetis aut inertissimae segnitiae est aut in dolore. Omnis autem privatione doloris putat Epicurus.

Contents

Acknowledgments	2
1 Introduction	4
1.1 History	5
1.2 Statistical-to-Computational Gap	5
1.3 Overlap Gap Property	6
1.4 Our Results	6
1.5 Existing Results	6
1.6 Our Results	7
1.7 Notation and Conventions	7
1.7.1 Glossary:	8
2 Low-Degree Algorithms	9
2.1 Coordinate Degree and L^2 Stability	9
2.2 Hermite Polynomials	13
2.3 Stability of Low-Degree Algorithms	15
3 Proof of Strong Low-Degree Hardness	17
3.1 Hardness for Low Degree Polynomial Algorithms	17
3.2 Proof for Low Coordinate-Degree Algorithms	21
3.3 Summary of Parameters	25
4 Extensions to Real-Valued Algorithms	27
4.1 Solutions repel meow	27
4.2 Proof of Hardness for Close Algorithms	28
4.3 Truly Random Rounding	31
5 Conclusion	34
5.1 Future Work	34
Bibliography	35

1 Introduction

Suppose we have N items, each with associated weights. How should we divide these items into two groups such that the sum of their weights is as close as possible? Alternatively, is it possible to divide these items into two groups such that the absolute difference of the sum of their weights is below a certain threshold? This question is known in statistics, physics, and computer science as the *number partitioning problem (NPP)*, and has been the subject of intense study from the 1970s to the present day.

Let g_1, \dots, g_N be N real numbers. The *number partitioning problem (NPP)* asks: what is the subset A of $[N] := \{1, 2, \dots, N\}$ such that the sum of the g_i for $i \in A$ and the sum of the remaining g_i are as close as possible? More formally, the A we want to find is the one minimizing the discrepancy

$$\left| \sum_{i \in A} g_i - \sum_{i \notin A} g_i \right|.$$

When rephrased as a decision problem (i.e., whether there exists an A such that the discrepancy is zero, or sufficiently small), the NPP is NP-complete; this can be shown by reduction from the subset sum problem. The NPP is also one of the six basic NP-complete problems of Garey and Johnson, and of those, the only one to deal with numbers (Garey and Johnson 1979, § 3.1).

(talk about modifications and variants?)

The number partitioning problem can be rephrased in the following way. Let our instance g_1, \dots, g_N be identified with a point $g \in \mathbf{R}^N$. Then, a choice of $A \subseteq [N]$ is equivalent to choosing a point x in the N -dimensional binary hypercube $\Sigma_N := \{\pm 1\}^N$, and the discrepancy of x is now $|\langle g, x \rangle|$. The goal is now to find the x minimizing this discrepancy:

$$\min_{x \in \Sigma_N} |\langle g, x \rangle|.$$

The number partitioning problem and algorithms designed to solve it have myriad practical applications.

Early work by Coffman, Garey, and Johnson, as well as by Tsai, looked at utilizing such algorithms for multiprocessor scheduling: dividing a group of tasks of approximately known runtimes across a pool of processors (Coffman, Garey, and Johnson 1978). Coffman and Lueker also describe how the NPP can be applied as a framework for allocating material stocks, such as steel coils in factories, paintings in museums, or advertisements in newspapers (Coffman and Lueker 1991).

On the other hand, in 1976, Merkle and Hellman devised one of the earliest public key cryptography schemes, deriving its hardness from their belief that a variant of the NPP was computationally difficult to solve – at the time, it was not yet known whether the NPP was NP-complete or not (Merkle and Hellman 1978). Their proposal was for the receiver, say Alice, to generate as a public key

N natural numbers (a_1, \dots, a_N) , with N typically around 100 and each a_i around 200 bits long. Then, to encrypt a N -bit message, $x = (x_1, \dots, x_N)$, with $x_i \in \{0, 1\}$, the sender, say Bob, could compute

$$b := \sum_{i \in N} a_i x_i,$$

and send the ciphertext b to Alice. Any eavesdropper would know a_1, \dots, a_N , as well as b , and decrypting the message involved finding a subset of the a_i adding up to b : this is the *knapsack problem*, which is NP-complete. However, such NP-completeness is only a worst-case hardness guarantee; Merkle and Hellman’s scheme involved Alice choosing a_1, \dots, a_N by cryptographically scrambling a sequence (a'_1, \dots, a'_N) for which solving the NPP was easy, enabling the receiver to practically decrypt the message x from the ciphertext b . In 1984, Shamir – one of the developers of the RSA cryptosystem still in use today – showed that one could exploit this public key generation process to reduce the “hard” knapsack problem to one which was solvable in polynomial time, rendering the Merkle-Hellman scheme insecure (Shamir 1982). While today, Merkle-Hellman is but a footnote in the history of cryptography, it demonstrates the importance of looking beyond worst-case hardness and expanding complexity theory to describe the difficulty of the average problem instance.

One particularly important application of the NPP in statistics comes from the design of *randomized controlled trials*. Consider N individuals, each with a set of covariate information $g_i \in \mathbf{R}^d$. Then the problem is to divide them into a treatment group (denoted A_+) and a control group (denoted A_-), subject each to different conditions, and evaluate the responses. In order for such a trial to be accurate, it is necessary to ensure that the covariates across both groups are roughly the same; in our notation, this equates to finding an A_+ (with $A_- := [N] \setminus A_+$) to minimize

$$\min_{A_+ \subseteq [N]} \left\| \sum_{i \in A_+} g_i - \sum_{i \in A_-} g_i \right\|_{\infty}.$$

This multidimensional extension of the NPP is often termed the *vector balancing problem (VBP)*, and many algorithms for solving the NPP/VBP come from designing such randomized controlled trials (Krieger, Azriel, and Kapelner 2019; Harshaw et al. 2023).

An orthogonal extension to the NPP is the *multiway number partitioning problem (MWNPP)*: here we want to partition g_1, \dots, g_N into M subsets such that the within-subset sums are mutually close. While what “mutually close” might mean is ()

Other applications.

- Circuit design, etc.

Two questions of interest:

1. What is optimal solution.
2. How to find optimal solution.

1.1 History

1.2 Statistical-to-Computational Gap

Non-planted models:

- Random constraint satisfaction: (Mézard, Mora, and Zecchina 2005; Achlioptas and Coja-Oghlan 2008; Kothari et al. 2017).
- Maximum independent sets in sparse random graphs (Gamarnik and Sudan 2013; Coja-Oghlan and Efthymiou 2015).
- Largest submatrix (Gamarnik and Li 2016)
- p -spin model: (Gamarnik and Jagannath 2019; Montanari 2019)
- diluted p -spin model: (Chen et al. 2019)

Planted models:

- matrix principal component analysis (Berthet and Rigollet 2013; Lesieur, Krzakala, and Zdeborová 2015; Lesieur, Krzakala, and Zdeborová 2015)
- tensor PCA (Hopkins, Shi, and Steurer 2015; Hopkins et al. 2017; Arous, Gheissari, and Jagannath 2020)
- high dimensional linear regression (Gamarnik and Zadik 2019a; 2019b)
- planted clique problem (Jerrum 1992; Deshpande and Montanari 2015; Meka, Potechin, and Wigderson 2015; Barak et al. 2016; Gamarnik and Zadik 2019c)

Evidence of hardness:

Failure of MCMC: (Huang and Sellke 2025; Jerrum 1992) Failure of AMP: (Zdeborová and Krzakala 2016; Bandeira, Perry, and Wein 2018) Reductions from planted clique - (Berthet and Rigollet 2013; Brennan and Bresler 2019; Brennan, Bresler, and Huleihel 2019) Lower bounds against Sum of Squares hierarchy: (Hopkins, Shi, and Steurer 2015; Hopkins et al. 2017; Raghavendra, Schramm, and Steurer 2019; Barak et al. 2016) Lower bounds in statistical query model: (Kearns 1998; Diakonikolas, Kane, and Stewart 2017; Feldman et al. 2016) Low degree methods, and low degree likelihood ratio: (Hopkins 2018; Kunisky, Wein, and Bandeira 2019)

1.3 Overlap Gap Property

1.4 Our Results

Low degree heuristic: degree D algorithms are a proxy for the class of $e^{\tilde{O}(D)}$ -time algorithms.

1.5 Existing Results

1. $X_i, 1 \leq i \leq n$ i.i.d. uniform from $\{1, 2, \dots, M := 2^m\}$, with $\kappa := \frac{m}{n}$, then phase transition going from $\kappa < 1$ to $\kappa > 1$.
2. Average case, X_i i.i.d. standard Normal.
3. Karmarkar [KKLO86] - NPP value is $\Theta(\sqrt{N}2^{-N})$ whp as $N \rightarrow \infty$ (doesn't need Normality).
4. Best polynomial-time algorithm: Karmarkar-Karp [KK82] - Discrepancy $O(N^{-\alpha \log N}) = 2^{-\Theta(\log^2 N)}$ whp as $N \rightarrow \infty$
5. PDM (paired differencing) heuristic - fails for i.i.d. uniform inputs with objective $\Theta(n^{-1})$ (Lueker).
6. LDM (largest differencing) heuristic - works for i.i.d. Uniforms, with $n^{-\Theta(\log n)}$ (Yakir, with constant $\alpha = \frac{1}{2 \ln 2}$ calculated non-rigorously by Boettcher and Mertens).
7. Krieger - $O(n^{-2})$ for balanced partition.
8. Hoberg [HHRY17] - computational hardness for worst-case discrepancy, as poly-time oracle that can get discrepancy to within $O(2^{\sqrt{n}})$ would be oracle for Minkowski problem.

9. Gamarnik-Kizildag: Information-theoretic guarantee $E_n = n$, best computational guarantee $E_n = \Theta(\log^2 n)$.
10. Existence of m -OGP for $m = O(1)$ and $E_n = \Theta(n)$.
11. Absence for $\omega(1) \leq E_n = o(n)$
12. Existence for $\omega(\sqrt{n \log_2 n}) \leq E_n \leq o(n)$ for $m = \omega_{n(1)}$ (with changing η, β)
 1. While OGP not ruled out for $E_n \leq \omega(\sqrt{n \log_2 n})$, argued that it is tight.
13. For $\varepsilon \in (0, \frac{1}{5})$, no stable algorithm can solve $\omega(n \log^{-\frac{1}{5}+\varepsilon} n) \leq E_n \leq o(n)$
14. Possible to strengthen to $E_n = \Theta(n)$ (as $2^{-\Theta(n)} \leq 2^{-o(n)}$)

1.6 Our Results

1.7 Notation and Conventions

Definition 1.1. Let $x \in \Sigma_N$. The *energy* of x (with respect to the instance g) is

$$E(x; g) := -\log_2 |\langle g, x \rangle|.$$

The *solution set* $S(E; g)$ is the set of all $x \in \Sigma_N$ that have energy at least E , i.e. that satisfy

$$|\langle g, x \rangle| \leq 2^{-E}. \quad (1.1)$$

- This terminology is motivated by the statistical physics literature, wherein random optimization problems are often reframed as energy maximization over a random landscape (Mertens 2001).
- Observe that minimizing the discrepancy $|\langle g, x \rangle|$ corresponds to maximizing the energy E .

Conventions:

1. On \mathbf{R}^N we write $\|\cdot\|_2 = \|\cdot\|$ for the Euclidean norm, and $\|\cdot\|_1$ for the ℓ^1 norm.
2. If $x \in \mathbf{R}^N$ and $S \subseteq [N]$, then x_S is vector with

$$(x_S)_i = \begin{cases} x_i & i \in S, \\ 0 & \text{else.} \end{cases}$$

In particular, for $x, y \in \mathbf{R}^N$,

$$\langle x_S, y \rangle = \langle x, y_S \rangle = \langle x_S, y_S \rangle.$$

3. meow
4. $B(x, r) = \{y \in \mathbf{R}^N : \|y - x\| < r\}$ is ℓ^2 unit ball.
5. Recall by Jensen's inequality that for any real numbers d_1, \dots, d_n , we have

$$\left(\sum_{i=1}^n d_i \right)^2 \leq n \sum_{i=1}^n d_i^2.$$

We will use this in the following way: suppose $x^{(1)}, \dots, x^{(n)}, x^{(n+1)}$ are n vectors in \mathbf{R}^N . Then

$$\|x^{(1)} - x^{(n+1)}\|^2 \leq \left(\sum_{i=1}^n \|x^{(i)} - x^{(i+1)}\| \right)^2 \leq n \sum_{i=1}^n \|x^{(i)} - x^{(i+1)}\|^2 \quad (1.2)$$

Throughout we will make key use of the following lemma:

Lemma 1.2 (Normal Small-Probability Estimate). *Let $E, \sigma^2 > 0$, and μ, Z be random variables with $Z \mid \mu \sim \mathcal{N}(\mu, \sigma^2)$. for σ^2 a constant. Then*

$$\mathbf{P}(|Z| \leq 2^{-E} \mid \mu) \leq \exp_2\left(-E - \frac{1}{2} \log_2(\sigma^2) + O(1)\right). \quad (1.3)$$

Proof: Observe that conditional on μ , the distribution of Z is bounded as

$$\varphi_{Z|\mu}(z) \leq \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(z-\mu)^2}{2\sigma^2}} \leq (2\pi\sigma^2)^{-1/2}.$$

Integrating over $|z| \leq 2^{-E}$ then gives (1.3), via

$$\mathbf{P}(|Z| \leq 2^{-E}) = \int_{|z| \leq 2^{-E}} (2\pi\sigma^2)^{-1/2} dz \leq 2^{-E - \frac{1}{2} \log_2(2\pi\sigma^2) + 1}. \quad \square$$

Note that this is decreasing function of σ^2 , e.g. it's bounded by $\exp_2(-E - \frac{1}{2} \log_2(\min \sigma^2))$ (this bound is trivial unless $\sigma^2 \Rightarrow \gamma > 0$).

Lemma 1.3. *Suppose that $K \leq N/2$, and let $h(x) = -x \log_2(x) - (1-x) \log_2(x)$ be the binary entropy function. Then, for $p := K/N$,*

$$\sum_{k \leq K} \binom{N}{k} \leq \exp_2(Nh(p)) \leq \exp_2\left(2Np \log_2\left(\frac{1}{p}\right)\right).$$

Proof: Consider a $\text{Bin}(N, p)$ random variable S . Summing its PMF from 0 to K , we have

$$1 \geq \mathbf{P}(S \leq K) = \sum_{k \leq K} \binom{N}{k} p^k (1-p)^{N-k} \geq \sum_{k \leq K} \binom{N}{k} p^K (1-p)^{N-K}.$$

Here, the last inequality follows from the fact that $p \leq (1-p)$, and we multiply each term by $\left(\frac{p}{1-p}\right)^{K-k} \leq 1$. Now rearrange to get

$$\begin{aligned} \sum_{k \leq K} \binom{N}{k} &\leq p^{-K} (1-p)^{-(N-K)} \\ &= \exp_2(-K \log_2(p) - (N-K) \log_2(1-p)) \\ &= \exp_2\left(N \cdot \left(-\frac{K}{N} \log_2(p) - \left(\frac{N-K}{N}\right) \log_2(1-p)\right)\right) \\ &= \exp_2(N \cdot (-p \log_2(p) - (1-p) \log_2(1-p))) = \exp_2(Nh(p)). \end{aligned}$$

The final equality then follows from the bound $h(p) \leq 2p \log_2(1/p)$ for $p \leq 1/2$. \square

1.7.1 Glossary:

1. “instance”/“disorder” - g , instance of the NPP problem
2. “discrepancy” - for a given g , value of $\min_{x \in \Sigma_N} |\langle g, x \rangle|$
3. “energy” - negative exponent of discrepancy, i.e. if discrepancy is 2^{-E} , then energy is E . Lower energy indicates “worse” discrepancy.
4. “near-ground state”/“approximate solution”

2 Low-Degree Algorithms

For our purposes, an *algorithm* is a function which takes as input a problem instance $g \in \mathbf{R}^N$ and outputs some $x \in \Sigma_N$. This definition can be extended to functions giving outputs on \mathbf{R}^N , and rounding to a vertex on the hypercube Σ_N . Alternatively, we could consider *randomized algorithms* via taking as additional input some randomness ω independent of the problem instance. However, most of our analysis will focus on the deterministic case.

To further restrict the category of algorithms considered, we specifically restrict to *low degree algorithms*. Compared to analytically-defined classes of algorithms (e.g. Lipschitz), these algorithms have a regular algebraic structure that we can exploit to precisely control their stability properties. In particular, our goal is to show *strong low degree hardness*, in the sense of (Huang and Sellke 2025, Def. 3).

Definition 2.1 (Strong Low-Degree Hardness). A random search problem, namely a N -indexed sequence of input vectors $y_N \in \mathbf{R}^{d_N}$ and random subsets $S_N = S_{N(y_N)} \subseteq \Sigma_N$, exhibits *strong low degree hardness up to degree $D \leq o(D_N)$* if, for all sequences of degree $o(D_N)$ algorithms (\mathcal{A}_N) with $\mathbf{E}\|\mathcal{A}(y_N)\|^2 \leq O(N)$, we have

$$\mathbf{P}(\mathcal{A}(y_N) \in S_N) \leq o(1).$$

In addition, degree D polynomials are a heuristic proxy for the class of $e^{\widetilde{O}(D)}$ -time algorithms (Hopkins 2018; Kothari et al. 2017). Thus, strong low degree hardness up to $o(N)$ can be thought of as evidence of requiring exponential (i.e. $e^{\Omega(N)}$) time to find globally optimal solutions.

For the case of NPP, we consider two distinct notions of degree. One is traditional polynomial degree, which has an intuitive interpretation, but the other, known in the literature as “coordinate degree,” is a more flexible notion which can be applied to a much broader class of algorithms. As we will see in Section 3, these classes of algorithms exhibit quantitatively different behavior, in line with existing heuristics for the “brittleness” of NPP.

2.1 Coordinate Degree and L^2 Stability

First, we consider a general class of putative algorithms, where the notion of “degree” corresponds to how many variables can interact nonlinearly with each other. Given this notion, deriving stability bounds becomes a straightforward piece of functional analysis. To start, recall the notion of L^2 functions:

Definition 2.2. Let π be a probability distribution on \mathbf{R} . The L^2 space $L^2(\mathbf{R}^N, \pi^{\otimes N})$ is the space of functions $f : \mathbf{R}^N \rightarrow \mathbf{R}$ with finite L^2 norm.

$$\mathbf{E}[f^2] := \int_{x=(x_1, \dots, x_n) \in \mathbf{R}^N} f(x)^2 d\pi^{\otimes N}(x) < \infty.$$

Alternatively, this is the space of L^2 functions of N i.i.d. random variables x_i , distributed as π .

Note that this is an extremely broad class of functions; for instance, all bounded functions are L^2 .

Given any function $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$, we can consider how it depends on various subsets of the N input coordinates. In principle, everything about f should be reflected in how it acts on all possible such subsets. To formalize this intuition, define the following coordinate projection:

Definition 2.3. Let $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ and $J \subseteq [N]$, with $\bar{J} = [N] \setminus J$. The *projection of f onto J* is the function $f^{\subseteq J} : \mathbf{R}^N \rightarrow \mathbf{R}$ given by

$$f^{\subseteq J}(x) = \mathbf{E}[f(x_1, \dots, x_n) \mid x_i, i \in J] = \mathbf{E}[f(x) \mid x_J]$$

Intuitively $f^{\subseteq J}$ is f with the \bar{J} coordinates re-randomized, so $f^{\subseteq J}$ only depends on the coordinates in J . However, depending on how f accounts for higher-order interactions, it might be the case that $f^{\subseteq J}$ is fully described by some $f^{\subseteq J'}$, for $J' \subsetneq J$. What we really want is to decompose f as

$$f = \sum_{S \subseteq [N]} f^{\subseteq S} \quad (2.1)$$

where each $f^{\subseteq S}$ only depends on the coordinates in S , but not any smaller subset. That is, if $T \not\subseteq S$ and g depends only on the coordinates in T , then $\langle f^{\subseteq S}, g \rangle = 0$.

This decomposition, often called the *Efron-Stein, orthogonal, or Hoeffding* decomposition, does indeed exist, and exhibits the following combinatorial construction. Our presentation largely follows (O'Donnell 2021, § 8.3), as well as the paper (Kunisky 2024a).

The motivating fact is that for any $J \subseteq [N]$, we should have

$$f^{\subseteq J} = \sum_{S \subseteq J} f^{\subseteq S}. \quad (2.2)$$

Intuitively, $f^{\subseteq J}$ captures everything about f depending on the coordinates in J , and each $f^{\subseteq S}$ captures precisely the interactions within each subset S of J . The construction of $f^{\subseteq S}$ proceeds by inverting this formula.

First, we consider the case $J = \emptyset$. It is clear that $f^{\subseteq \emptyset} = f^{\subseteq \emptyset}$, which, by Definition 2.3 is the constant function $\mathbf{E}[f]$. Next, if $J = \{j\}$ is a singleton, (2.2) gives

$$f^{\subseteq \{j\}} = f^{\subseteq \emptyset} + f^{\subseteq \{j\}},$$

and as $f^{\subseteq \{j\}}(x) = \mathbf{E}[f \mid x_j]$, we get

$$f^{\subseteq \{j\}} = \mathbf{E}[f \mid x_j] - \mathbf{E}[f].$$

This function only depends on x_j ; all other coordinates are averaged over, thus measuring how the expectation of f changes given x_j .

Continuing on to sets of two coordinates, some brief manipulation gives, for $J = \{i, j\}$,

$$\begin{aligned} f^{\subseteq \{i, j\}} &= f^{\subseteq \emptyset} + f^{\subseteq \{i\}} + f^{\subseteq \{j\}} + f^{\subseteq \{i, j\}} \\ &= f^{\subseteq \emptyset} + (f^{\subseteq \{i\}} - f^{\subseteq \emptyset}) + (f^{\subseteq \{j\}} - f^{\subseteq \emptyset}) + f^{\subseteq \{i, j\}} \\ \therefore f^{\subseteq \{i, j\}} &= f^{\subseteq \{i, j\}} - f^{\subseteq \{i\}} - f^{\subseteq \{j\}} + f^{\subseteq \emptyset}. \end{aligned}$$

We can imagine that this accounts for the two-way interaction of i and j , namely $f^{\subseteq\{i,j\}} = \mathbf{E}[f \mid x_i, x_j]$, while “correcting” for the one-way effects of x_i and x_j individually. Inductively, we can continue on and define all the $f^{=J}$ via inclusion-exclusion, as

$$f^{=J} := \sum_{S \subseteq J} (-1)^{|J|-|S|} f^{\subseteq S} = \sum_{S \subseteq J} (-1)^{|J|-|S|} \mathbf{E}[f \mid x_S].$$

This construction, along with some direct calculations, leads to the following theorem on Efron-Stein decompositions:

Theorem 2.4 ((O'Donnell 2021, Thm 8.35)). *Let $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$. Then f has a unique Efron-Stein decomposition as*

$$f = \sum_{S \subseteq [N]} f^{=S}$$

where the functions $f^{=S} \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ satisfy

1. $f^{=S}$ depends only on the coordinates in S ;
2. if $T \subsetneq S$ and $g \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ only depends on coordinates in T , then $\langle f^{=S}, g \rangle = 0$.

In addition, this decomposition has the following properties:

3. Condition 2. holds whenever $S \not\subseteq T$.
4. The decomposition is orthogonal: $\langle f^{=S}, f^{=T} \rangle = 0$ for $S \neq T$.
5. $\sum_{S \subseteq T} f^{=S} = f^{=T}$.
6. For each $S \subseteq [N]$, $f \mapsto f^{=S}$ is a linear operator.

In summary, this decomposition of any $L^2(\mathbf{R}^N, \pi^{\otimes N})$ function into its different interaction levels not only uniquely exists, but is an orthogonal decomposition, enabling us to apply tools from elementary Fourier analysis.

Theorem 2.4 further implies that we can define subspaces of $L^2(\mathbf{R}^N, \pi^{\otimes N})$ (see also (Kunisky 2024a, § 1.3))

$$\begin{aligned} V_J &:= \{f \in L^2(\mathbf{R}^N, \pi^{\otimes N}) : f = f^{\subseteq J}\}, \\ V_{\leq D} &:= \sum_{\substack{J \subseteq [N] \\ |J| \leq D}} V_J. \end{aligned} \tag{2.3}$$

These capture functions which only depend on some subset of coordinates, or some bounded number of coordinates. Note that $V_{[N]} = V_{\leq N} = L^2(\mathbf{R}^N, \pi^{\otimes N})$.

With this, we can define the notion of “coordinate degree”:

Definition 2.5. The *coordinate degree* of a function $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ is

$$\text{cdeg}(f) := \max\{|S| : S \subseteq [N], f^{=S} \neq 0\} = \min\{D : f \in V_{\leq D}\}$$

If $f = (f_1, \dots, f_M) : \mathbf{R}^N \rightarrow \mathbf{R}^M$ is a multivariate function, then

$$\text{cdeg}(f) := \max_{i \in [M]} \text{cdeg}(f_i).$$

Intuitively, the coordinate degree is the maximum size of (nonlinear) multivariate interaction that f accounts for. Of course, this degree is also bounded by N , very much unlike polynomial degree. Note as a special case that any multivariate polynomial of degree D has coordinate degree at most D . As an example, the function $x_1 + x_2$ has both polynomial degree and coordinate degree 1, while $x_1 + x_2^2$ has polynomial degree 2 and coordinate degree 1. We are especially interested in algorithms coming from functions in $V_{\leq D}$, which we term *low coordinate degree algorithms*.

As we are interested in how these function behaves under small changes in its input, we are led to consider the following “noise operator,” which lets us measures the effect of small changes in the input on the coordinate decomposition. First, we need the following notion of distance between problem instances:

Definition 2.6. For $p \in [0, 1]$, and $x \in \mathbf{R}^N$, we say $y \in \mathbf{R}^N$ is *p-resampled from* x , denoted $y \sim \pi_p^{\otimes N}(x)$, if y is chosen as follows: for each $i \in [N]$, independently,

$$y_i = \begin{cases} x_i & \text{with probability } p \\ \text{drawn from } \pi & \text{with probability } 1 - p \end{cases}$$

We say (x, y) are a *p-resampled pair*.

Note that being *p-resampled* and being *p-correlated* are rather different - for one, there is a nonzero probability that, for π a continuous probability distribution, $x = y$ when they are *p-resampled*, even though this a.s. never occurs if they were *p-correlated*.

Definition 2.7. For $p \in [0, 1]$, the *noise operator* T_p is the linear operator on $L^2(\mathbf{R}^N, \pi^{\otimes N})$ defined by

$$T_p f(x) = \mathbf{E}_{y \sim \pi_p^{\otimes N}(x)}[f(y)]$$

In particular, $\langle f, T_p f \rangle = \mathbf{E}_{(x, y) \text{ p-resampled}}[f(x) \cdot f(y)]$.

This noise operator changes the Efron-Stein decomposition, and hence the behavior of low coordinate degree functions, in a controlled way:

Lemma 2.8. Let $p \in [0, 1]$ and $f \in L^2(\mathbf{R}^N, \pi^{\otimes N})$ have Efron-Stein decomposition $f = \sum_{S \subseteq [N]} f^{=S}$. Then

$$T_p f(x) = \sum_{S \subseteq [N]} p^{|S|} f^{=S}.$$

Proof: Let J denote a *p-random subset* of $[N]$, i.e. with J formed by including each $i \in [N]$ independently with probability p . By definition, $T_p f(x) = \mathbf{E}_J[f^{\subseteq J}(x)]$ (i.e. pick a random subset of coordinates to fix, and re-randomize the rest). We know by [Theorem 2.4](#) that $f^{\subseteq J} = \sum_{S \subseteq J} f^{=S}$, so

$$T_p f(x) = \mathbf{E}_J \left[\sum_{S \subseteq J} f^{=S} \right] = \sum_{S \subseteq [N]} \mathbf{E}_J[I(S \subseteq J)] \cdot f^{=S} = \sum_{S \subseteq [N]} p^{|S|} f^{=S},$$

since for a fixed $S \subseteq [N]$, the probability that $S \subseteq J$ is $p^{|S|}$. □

Thus, we can derive the following stability bound on low coordinate degree functions.

Theorem 2.9. Let $p \in [0, 1]$ and let $f = (f_1, \dots, f_M) : \mathbf{R}^N \rightarrow \mathbf{R}^M$ be a multivariate function with coordinate degree D and each $f_i \in L^2(\mathbf{R}^N, \pi^{\otimes N})$. Suppose that (x, y) are a p -resampled pair under $\pi^{\otimes N}$, and $\mathbf{E}\|f(x)\|^2 = 1$. Then

$$\mathbf{E}\|f(x) - f(y)\|^2 \leq 2(1 - p^D) \leq 2(1 - p)D. \quad (2.4)$$

Proof: Observe that

$$\begin{aligned} \mathbf{E}\|f(x) - f(y)\|^2 &= \mathbf{E}\|f(x)\|^2 + \mathbf{E}\|f(y)\|^2 - 2\mathbf{E}\langle f(x), f(y) \rangle \\ &= 2 - 2\left(\sum_i \mathbf{E}[f_i(x)f_i(y)]\right) \\ &= 2 - 2\left(\sum_i \langle f_i, T_p f_i \rangle\right). \end{aligned} \quad (2.5)$$

Here, we have for each $i \in [M]$ that

$$\langle f_i, T_p f_i \rangle = \left\langle \sum_{S \subseteq [N]} f_i^{\neg S}, \sum_{S \subseteq [N]} p^{|S|} f_i^{\neg S} \right\rangle = \sum_{S \subseteq [N]} p^{|S|} \|f_i^{\neg S}\|^2,$$

by Lemma 2.8 and orthogonality. Now, as each f_i has coordinate degree at most D , the sum above can be taken only over $S \subseteq [N]$ with $0 \leq |S| \leq D$, giving the bound

$$p^D \mathbf{E}[f_i(x)^2] \leq \langle f_i, T_p f_i \rangle = \mathbf{E}[f_i(x) \cdot T_p f_i(x)] \leq \mathbf{E}[f_i(x)^2].$$

Summing up over i , and using that $\mathbf{E}\|f(x)\|^2 = 1$, gives

$$p^D \leq \sum_i \langle f_i, T_p f_i \rangle = \mathbf{E}\langle f(x), f(y) \rangle \leq 1.$$

Finally, we can substitute into (2.5) to get²

$$\mathbf{E}\|f(x) - f(y)\|^2 \leq 2 - 2p^D = 2(1 - p^D) \leq 2(1 - p)D. \quad \square$$

2.2 Hermite Polynomials

Alternatively, we can consider the much more restrictive (but more concrete) class of honest polynomials. When considered as functions of independent Normal variables, such functions admit a simple description in terms of *Hermite polynomials*, which enables us to prove similar bounds as Theorem 2.9. This theory is much more classical, so we encourage the interested reader to see (O'Donnell 2021, § 11) for details.

Definition 2.10. Let γ_N be the N -dimensional standard Normal measure on \mathbf{R}^N . Then the N -dimensional Gaussian space is the space $L^2(\mathbf{R}^N, \gamma^N)$ of L^2 functions of N i.i.d. standard Normal r.v.s.

Note that under the usual L^2 inner product, $\langle f, g \rangle = \mathbf{E}[f \cdot g]$, this is a separable Hilbert space.

It is a well-known fact that the monomials $1, z, z^2, \dots$ form a complete basis for $L^2(\mathbf{R}, \gamma)$ (O'Donnell 2021, Thm 11.22). However, these are far from an orthonormal “Fourier” basis; for

²The last inequality follows from $(1 - p^D) = (1 - p)(1 + p + p^2 + \dots + p^{D-1})$; the bound is tight for $p \approx 1$.

instance, we know $\mathbf{E}[z^2] = 1$ for $z \sim \mathcal{N}(0, 1)$. By the Gram-Schmidt process, these monomials can be converted into the (*normalized*) *Hermite polynomials* h_j for $j \geq 0$, given as

$$h_0(z) = 1, \quad h_1(z) = z, \quad h_2(z) = \frac{z^2 - 1}{\sqrt{2}}, \quad h_3(z) = \frac{z^3 - 3z}{\sqrt{6}}, \quad \dots \quad (2.6)$$

Note here that each h_j is a degree j polynomial. With these, we have:

Theorem 2.11 ((O'Donnell 2021, Prop 11.30)). *The Hermite polynomials $(h_j)_{j \geq 0}$ form a complete orthonormal basis for $L^2(\mathbf{R}, \gamma)$.*

To extend this to $L^2(\mathbf{R}^N, \gamma^N)$, we can take products. For a multi-index $\alpha \in \mathbb{N}^N$, we define the multivariate Hermite polynomial $h_\alpha : \mathbf{R}^N \rightarrow \mathbf{R}$ as

$$h_\alpha(z) := \prod_{j=1}^N h_{\alpha_j}(z_j).$$

The degree of h_α is clearly $|\alpha| = \sum_j \alpha_j$.

Theorem 2.12. *The Hermite polynomials $(h_\alpha)_{\alpha \in \mathbb{N}^N}$ form a complete orthonormal basis for $L^2(\mathbf{R}^N, \gamma^N)$. In particular, every $f \in L^2(\mathbf{R}^N, \gamma^N)$ has a unique expansion in L^2 norm as*

$$f(z) = \sum_{\alpha \in \mathbb{N}^N} \hat{f}(\alpha) h_\alpha(z).$$

As a consequence of the uniqueness of the expansion in , we see that polynomials are their own Hermite expansion. Namely, let $H^{\leq k} \subseteq L^2(\mathbf{R}^N, \gamma^N)$ be the subset of multivariate polynomials of degree at most k . Then, any $f \in H^{\leq k}$ can be Hermite expanded as

$$f(z) = \sum_{\alpha \in \mathbb{N}^N} \hat{f}(\alpha) h_\alpha(z) = \sum_{|\alpha| \leq k} \hat{f}(\alpha) h_\alpha(z).$$

Thus, $H^{\leq k}$ is the closed linear span of the set $\{h_\alpha : |\alpha| \leq k\}$.

When working with honest polynomials, the traditional notion of correlation is a much more natural measure of “distance” between inputs:

Definition 2.13. Let (x, y) be N -dimensional standard Normal vectors. We say (x, y) are *p-correlated* if (x_i, y_i) are *p-correlated* for each $i \in [N]$, and these pairs are mutually independent.

In a similar way to the Efron-Stein case, we can consider the resulting “noise operator,” as a way of measuring the effect on a function of a small change in the input.

Definition 2.14. For $p \in [0, 1]$, the *Gaussian noise operator* T_p is the linear operator on $L^2(\mathbf{R}^N, \gamma^N)$, given by

$$T_p f(x) = \mathbf{E}_{y \text{ p-correlated to } x} [f(y)] = \mathbf{E}_{y \sim \mathcal{N}(0, I_N)} \left[f\left(px + \sqrt{1-p^2}y\right) \right]$$

This operator admits a more classical description in terms of the Ornstein-Uhlenbeck semigroup, but we will not need that connection here. As it happens, a straightforward computation with the Normal moment generating function gives the following:

Lemma 2.15 ((O'Donnell 2021, Prop 11.37)). Let $p \in [0, 1]$ and $f \in L^2(\mathbf{R}^N, \gamma^N)$. Then $T_p f$ has Hermite expansion

$$T_p f = \sum_{\alpha \in \mathbb{N}^N} p^{|\alpha|} \hat{f}(\alpha) h_\alpha$$

and in particular,

$$\langle f, T_p f \rangle = \sum_{\alpha \in \mathbb{N}^N} p^{|\alpha|} \hat{f}(\alpha)^2.$$

With this in hand, we can prove a similar stability bound to [Theorem 2.9](#).

Theorem 2.16. Let $p \in [0, 1]$ and let $f = (f_1, \dots, f_M) : \mathbf{R}^N \rightarrow \mathbf{R}^M$ be a multivariate polynomial with degree D . Suppose that (x, y) are a p -correlated pair of standard Normal vectors, and $\mathbf{E}\|f(x)\|^2 = 1$. Then

$$\mathbf{E}\|f(x) - f(y)\|^2 \leq 2(1 - p^D) \leq 2(1 - p)D. \quad (2.7)$$

Proof: The proof is almost identical to that of [Theorem 2.9](#) (see also [\(Gamarnik, Jagannath, and Wein 2022, Lem. 3.4\)](#)). The main modification is to realize that for each f_i , having degree at most D implies that $\hat{f}_i(\alpha) = 0$ for $|\alpha| > D$. Thus, as $p^D \leq p^s \leq 1$ for all $s \leq D$, we can apply [Lemma 2.15](#) to get

$$p^D \mathbf{E}[f_i(x)^2] \leq \langle f_i, T_p f_i \rangle = \sum_{\alpha \in \mathbb{N}^N : |\alpha| \leq D} p^{|\alpha|} \hat{f}_i(\alpha)^2 \leq \mathbf{E}[f_i(x)^2].$$

From there, the proof proceeds as before. □

As a comparison to the case for functions with coordinate degree D , notice that [Theorem 2.16](#) gives, generically, a much looser bound. In exchange, being able to use p -correlation as a “metric” on the input domain will turn out to offer significant strengthenings in the arguments which follow, justifying equal consideration of both classes of functions.

2.3 Stability of Low-Degree Algorithms

With these notions of low degree functions/polynomials in hand, we can consider algorithms based on such functions.

Definition 2.17. A (randomized) algorithm is a measurable function $\mathcal{A} : (g, \omega) \mapsto x^* \in \Sigma^N$, where $\omega \in \Omega_N$ is an independent random variable. Such an \mathcal{A} is *deterministic* if it does not depend on ω .

In practice, we want to consider \mathbf{R}^N -valued algorithms as opposed to Σ_N -valued ones to avoid the resulting restrictions on the component functions. These can then be converted to Σ_N -valued algorithms by some rounding procedure. We discuss the necessary extensions to handling this rounding in [Section 4](#).

Definition 2.18. A polynomial algorithm is an algorithm $\mathcal{A}(g, \omega)$ where each coordinate of $\mathcal{A}(g, \omega)$ is given by a polynomial in the N entries of g . If \mathcal{A} is a polynomial algorithm, we say it has degree D if each coordinate has degree at most D (with at least one equality).

We can broaden the notion of polynomial algorithms (with their obvious notion of degree) to algorithms with a well-defined notion of coordinate degree:

Definition 2.19. Suppose an algorithm $\mathcal{A}(g, \omega)$ is such that each coordinate of $\mathcal{A}(-, \omega)$ is in $L^2(\mathbf{R}^N, \pi^{\otimes N})$. Then, the *coordinate degree* of \mathcal{A} is the maximum coordinate degree of each of its coordinate functions.

By the low degree heuristic, these algorithms can be interpreted as a proxy for time N^D -algorithms, unlike classes based off of their stability properties, such as Lipschitz/Hölder continuous algorithms. Yet in addition to this interpretability, these algorithms also have accessible stability bounds:

Proposition 2.20 (Low-Degree Stability – (Huang and Sellke 2025, Prop. 1.9)). *Suppose we have a deterministic algorithm \mathcal{A} with degree (resp. coordinate degree) $\leq D$ and norm $\mathbf{E}\|\mathcal{A}(g)\|^2 \leq CN$. Then, for inputs g, g' which are $(1 - \varepsilon)$ -correlated (resp. $(1 - \varepsilon)$ -resampled),*

$$\mathbf{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^2 \leq 2CD\varepsilon N, \quad (2.8)$$

and thus

$$\mathbf{P}\left(\|\mathcal{A}(g) - \mathcal{A}(g')\| \geq 2\sqrt{\eta N}\right) \leq \frac{CD\varepsilon}{2\eta} \asymp \frac{D\varepsilon}{\eta} \quad (2.9)$$

Proof: Let $C' := \mathbf{E}\|\mathcal{A}(g)\|^2$, and define the rescaling $\mathcal{A}' := \mathcal{A}/\sqrt{C'}$. Then, by Theorem 2.16 (or Theorem 2.9, in the low coordinate degree case), we have

$$\mathbf{E}\|\mathcal{A}'(g) - \mathcal{A}'(g')\|^2 = \frac{1}{C'} \mathbf{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^2 \leq 2D\varepsilon.$$

Multiplying by C' gives (2.8) (as $C' \leq CN$). Finally, (2.9) follows from Markov's inequality. \square

3 Proof of Strong Low-Degree Hardness

In this section, we prove strong low degree hardness for both low degree polynomial algorithms and algorithms with low Efron-Stein degree.

For now, we consider Σ_N -valued deterministic algorithms. We discuss the extension to \mathbf{R}^N -valued algorithms in [Section 4](#). As outlined in [Section 1.6](#), we show that TODO.

The key argument is as follows. Fix some energy levels E , depending on N . Suppose we have a Σ_N -valued, deterministic algorithm \mathcal{A} given by a degree D polynomial (resp. an Efron-Stein degree D function), and we have two instances $g, g' \sim \mathcal{N}(0, I_N)$ which are $(1 - \varepsilon)$ -correlated (resp. $(1 - \varepsilon)$ -resampled), for $\varepsilon > 0$. Say $\mathcal{A}(g) = x \in \Sigma_N$ is a solution with energy at least E , i.e. it “solves” this NPP instance. For ε close to 0, $\mathcal{A}(g') = x'$ will be close to x , by low degree stability. However, by adjusting parameters carefully, we can make it so that with high probability (exponential in E), there are no solutions to g' close to x . By application of a correlation bound on the probability of solving any fixed instance, we can conclude that with high probability, \mathcal{A} can’t find solutions to NPP with energy E .

Our argument utilizes what can be thought of as a “conditional” version of the overlap gap property. Traditionally, the overlap gap property is a global obstruction: one shows that with high probability, one cannot find a tuple of good solutions to a family of correlated instances which are all roughly the same distance apart. Here, however, we show a local obstruction - we condition on being able to solve a single instance, and show that after a small change to the instance, we cannot guarantee any solutions will exist close to the first one. This is an instance of the “brittleness,” so to speak, that makes NPP so frustrating to solve; even small changes in the instance break the landscape geometry, so that even if solutions exist, there’s no way to know where they’ll end up.

First moment details meow.

We start with some setup which will apply, with minor modifications depending on the nature of the algorithm in consideration, to all of the energy regimes in discussion. After proving some preliminary estimates, we establish the existence of our conditional landscape obstruction, which is of independent interest. Finally, we conclude by establishing low degree hardness in both the linear and sublinear energy regimes.

Explain more meow.

3.1 Hardness for Low Degree Polynomial Algorithms

First, consider the case of \mathcal{A} being a polynomial algorithm with degree D .

Let g, g' be $(1 - \varepsilon)$ -correlated standard Normal r.v.s, and let $x \in \Sigma_N$ depend only on g . Furthermore, let $\eta > 0$ be a parameter which will be chosen in a manner specified later. We define the following events:

$$S_{\text{solve}} = \{\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g')\} \quad (3.1)$$

$$\begin{aligned}
S_{\text{stable}} &= \{\|\mathcal{A}(g) - \mathcal{A}(g')\| \leq 2\sqrt{\eta N}\} \\
S_{\text{cond}}(x) &= \left\{ \nexists x' \in S(E; g') \text{ such that } \|x - x'\| \leq 2\sqrt{\eta N} \right\}
\end{aligned} \tag{3.1}$$

Intuitively, the first two events ask that the algorithm solves both instances and is stable, respectively. The last event, which depends on x , corresponds to the conditional landscape obstruction: for an x depending only on g , there is no solution to g' which is close to x .

Lemma 3.1. *We have, for $x := \mathcal{A}(g)$, $S_{\text{solve}} \cap S_{\text{stable}} \cap S_{\text{cond}}(x) = \emptyset$.*

Proof: Suppose that S_{solve} and S_{stable} both occur. Letting $x := \mathcal{A}(g)$ (which only depends on g) and $x' := \mathcal{A}(g')$, we have that $x' \in S(E; g')$ while also being within distance $2\sqrt{\eta N}$ of x . This contradicts $S_{\text{cond}}(x)$, thus completing the proof. \square

First, define $p_{\text{solve}}^{\text{cor}}$ as the probability that the algorithm solves a single random instance:

$$p_{\text{solve}}^{\text{cor}} = \mathbf{P}(\mathcal{A}(g) \in S(E; g)). \tag{3.2}$$

Then, we have the following correlation bound, which allows us to avoid union bounding over instances:

Lemma 3.2. *For g, g' being $(1 - \varepsilon)$ -correlated, we have*

$$\mathbf{P}(S_{\text{solve}}) = \mathbf{P}(\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g')) \geq (p_{\text{solve}}^{\text{cor}})^2$$

Proof: Let $\tilde{g}, g^{(0)}, g^{(1)}$ be three i.i.d. copies of g , and observe that g, g' are jointly representable as

$$g = \sqrt{1 - \varepsilon} \tilde{g} + \sqrt{\varepsilon} g^{(0)}, \quad g' = \sqrt{1 - \varepsilon} \tilde{g} + \sqrt{\varepsilon} g^{(1)}.$$

Thus, since g, g' are conditionally independent given \tilde{g} , we have

$$\begin{aligned}
\mathbf{P}(\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g')) &= \mathbf{E}[\mathbf{P}(\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g') \mid \tilde{g})] \\
&= \mathbf{E}[\mathbf{P}(\mathcal{A}(g) \in S(E; g) \mid \tilde{g})^2] \\
&\geq \mathbf{E}[\mathbf{P}(\mathcal{A}(g) \in S(E; g) \mid \tilde{g})]^2 = (p_{\text{solve}}^{\text{cor}})^2,
\end{aligned}$$

where the last line follows by Jensen's inequality. \square

Moreover, let us define $p_{\text{unstable}}^{\text{cor}}$ and $p_{\text{cond}}^{\text{cor}}(x)$ by

$$p_{\text{unstable}}^{\text{cor}} = 1 - \mathbf{P}(S_{\text{stable}}), \quad p_{\text{cond}}^{\text{cor}}(x) = 1 - \mathbf{P}(S_{\text{cond}}(x)).$$

In addition, define

$$p_{\text{cond}}^{\text{cor}} := \max_{x \in \Sigma_N} p_{\text{cond}}^{\text{cor}}(x). \tag{3.3}$$

By **Lemma 3.1**, we know that for $x := \mathcal{A}(g)$

$$\mathbf{P}(S_{\text{solve}}) + \mathbf{P}(S_{\text{stable}}) + \mathbf{P}(S_{\text{cond}}(x)) \leq 2,$$

and rearranging, we get that

$$(p_{\text{solve}}^{\text{cor}})^2 \leq p_{\text{unstable}}^{\text{cor}} + p_{\text{cond}}^{\text{cor}} \tag{3.4}$$

Our proof follows by showing that, for appropriate choices of ε and η , depending on D , E , and N , we have $p_{\text{unstable}}^{\text{cor}}, p_{\text{cond}}^{\text{cor}} = o(1)$.

To this end, we start by bounding the size of neighborhoods on Σ_N .

Proposition 3.3 (Hypercube Neighborhood Size). *Fix $x \in \Sigma_N$, and let $\eta \leq 1/2$. Then the number of x' within distance $2\sqrt{\eta N}$ of x is*

$$\left| \left\{ x' \in \Sigma_N : \|x - x'\| \leq 2\eta\sqrt{N} \right\} \right| \leq \exp_2(2\eta \log_2(1/\eta)N)$$

Proof: Let k be the number of coordinates which differ between x and x' (i.e. the Hamming distance). We have $\|x - x'\|^2 = 4k$, so $\|x - x'\| \leq 2\sqrt{\eta N}$ iff $k \leq N\eta$. Moreover, for $\eta \leq \frac{1}{2}$, $k \leq \frac{N}{2}$. Thus, by [Lemma 1.3](#), we get

$$\sum_{k \leq N\eta} \binom{N}{k} \leq \exp_2(Nh(\eta)) \leq \exp_2(2\eta \log_2(1/\eta)N). \quad \square$$

This shows that within a small neighborhood of any $x \in \Sigma_N$, the number of nearby points is exponential in N , with a more nontrivial dependence on η . The question is how many of these are solutions to a correlated/resampled instance.

First, we consider the conditional probability of any fixed $x \in \Sigma_N$ solving a $(1 - \varepsilon)$ -correlated problem instance g' , given g :

Putting together these bounds, we conclude the following fundamental estimates of $p_{\text{cond}}^{\text{cor}}$, i.e. of the failure of our conditional landscape obstruction.

Proposition 3.4 (Fundamental Estimate – Correlated Case). *Assume that (g, g') are $(1 - \varepsilon)$ -correlated standard Normal vectors. Then, for any x only depending on g ,*

$$p_{\text{cond}}^{\text{cor}}(x) := \mathbf{P} \left(\exists x' \in S(E; g') \text{ such that } \|x - x'\| \leq 2\sqrt{\eta N} \right) \leq \exp_2 \left(-E - \frac{1}{2} \log_2(\varepsilon) + 2\eta \log_2 \left(\frac{1}{\eta} \right) N + O(\log_2 N) \right).$$

Proof: For each x' within distance $2\sqrt{\eta N}$ of x , let

$$I_{x'} := I(x \in S(E; g')) = I(|\langle g', x' \rangle| \leq 2^{-E}),$$

so that

$$p_{\text{cond}}^{\text{cor}}(x) = \mathbf{E} \left[\sum_{\|x - x'\| \leq 2\sqrt{\eta N}} \mathbf{E}[I_{x'} \mid g] \right] = \mathbf{E} \left[\sum_{\|x - x'\| \leq 2\sqrt{\eta N}} \mathbf{P}(|\langle g', x' \rangle| \leq 2^{-E} \mid g) \right] \quad (3.5)$$

To bound the inner probability, let \tilde{g} be a Normal vector independent to g and set $p = 1 - \varepsilon$. Observe that g' can be represented as $g' = pg + \sqrt{1 - p^2}\tilde{g}$, so, $\langle g', x' \rangle = p\langle g, x' \rangle + \sqrt{1 - p^2}\langle \tilde{g}, x' \rangle$. We know $\langle \tilde{g}, x' \rangle \sim \mathcal{N}(0, N)$, so conditional on g , we have $\langle g', x' \rangle \mid g \sim \mathcal{N}(p\langle g, x' \rangle, (1 - p^2)N)$. Note that $\langle g', x' \rangle$ is nondegenerate for $(1 - p^2)N \geq \varepsilon N > 0$; thus by [Lemma 1.2](#), we get

$$\mathbf{P}(|\langle g', x' \rangle| \leq 2^{-E} \mid g) \leq \exp_2 \left(-E - \frac{1}{2} \log_2(\varepsilon) + O(\log_2 N) \right). \quad (3.6)$$

Finally, by [Proposition 3.3](#), the number of terms in the sum [\(3.5\)](#) is bounded by $\exp_2(2\eta \log_2(1/\eta)N)$, so given that [\(3.6\)](#) is independent of g , we conclude that

$$p_{\text{cond}}^{\text{cor}}(x) \leq \exp_2\left(-E + -\frac{1}{2}\log_2(\varepsilon) + 2\eta \log_2\left(\frac{1}{\eta}\right)N + O(\log_2 N)\right). \quad \square$$

Note for instance that ε can be exponentially small in E (e.g. $\varepsilon = \exp_2(-E/10)$), which for the case $E = \Theta(N)$ implies ε can be exponentially small in N .

Transition para meow.

Throughout this section, we let $E = \delta N$ for some $\delta > 0$, and aim to rule out the existence of low degree algorithms achieving these energy levels. This corresponds to the statistically optimal regime, as per [\(Karmarkar et al. 1986\)](#). These results roughly correspond to those in [\(Gamarnik and Kızıldağ 2021, Thm. 3.2\)](#), although their result applies to stable algorithms more generally, and does not show a low degree hardness-type result.

Theorem 3.5. *Let $\delta > 0$ and $E = \delta N$, and let g, g' be $(1 - \varepsilon)$ -correlated standard Normal r.v.s. Then, for any degree $D \leq o(\exp_2(\delta N/2))$ polynomial algorithm \mathcal{A} (with $\mathbb{E}\|\mathcal{A}(g)\|^2 \leq CN$), there exist $\varepsilon, \eta > 0$ such that $p_{\text{solve}}^{\text{cor}} = o(1)$.*

Proof: Recall from [\(3.4\)](#) that it suffices to show that both $p_{\text{cond}}^{\text{cor}}$ and $p_{\text{unstable}}^{\text{cor}}$ go to zero. Further, by [\(3.3\)](#) and [Proposition 3.4](#), we have

$$p_{\text{cond}}^{\text{cor}} \leq \exp_2\left(-E - \frac{1}{2}\log_2(\varepsilon) + 2\eta \log_2\left(\frac{1}{\eta}\right)N + O(\log_2 N)\right)$$

Thus, first choose η sufficiently small, such that $2\eta \log_2(1/\eta) < \delta/4$ – this results in η being independent of N . Next, choose $\varepsilon = \exp_2(-\delta N/2)$. This gives

$$p_{\text{cond}}^{\text{cor}} \leq \exp_2\left(-\delta N - \frac{1}{2}\left(-\frac{\delta N}{2}\right) + \frac{\delta N}{4} + O(\log_2 N)\right) = \exp_2\left(-\frac{\delta N}{2} + O(\log_2 N)\right) = o(1).$$

Moreover, for $D \leq o(\exp_2(\delta N/2))$, we get by [Proposition 2.20](#) that

$$p_{\text{unstable}}^{\text{cor}} \leq \frac{CD\varepsilon}{2\eta} \asymp \frac{D\varepsilon}{\eta} \asymp D \cdot \exp_2\left(-\frac{\delta N}{2}\right) \rightarrow 0.$$

By [\(3.4\)](#), we conclude that $(p_{\text{solve}}^{\text{cor}})^2 \leq p_{\text{unstable}}^{\text{cor}} + p_{\text{cond}}^{\text{cor}} = o(1)$, thus completing the proof. \square

Remark that this implies poly algs are really bad, requiring double exponential time. meow.

Next, we let $\omega(\log_2 N) \leq E \leq o(N)$.

Theorem 3.6. *Let $\omega(\log_2^2 N) \leq E \leq o(N)$, and let g, g' be $(1 - \varepsilon)$ -correlated standard Normal r.v.s. Then, for any polynomial algorithm \mathcal{A} with degree $D \leq o(\exp_2(E/4))$ (and with $\mathbb{E}\|\mathcal{A}(g)\|^2 \leq CN$), there exist $\varepsilon, \eta > 0$ such that $p_{\text{solve}}^{\text{cor}} = o(1)$.*

Proof: As in [Theorem 3.5](#), it suffices to show that both $p_{\text{cond}}^{\text{cor}}$ and $p_{\text{unstable}}^{\text{cor}}$ go to zero. To do this, we choose

$$\varepsilon = \exp_2\left(-\frac{E}{2}\right), \quad \eta = \frac{E}{16N \log_2(N/E)}. \quad (3.7)$$

With this choice of η , some simple analysis shows that for $\frac{E}{N} \ll 1$, we have that

$$\frac{E}{4N} > 2\eta \log_2(1/\eta).$$

Thus, by [Proposition 3.4](#), we get

$$\begin{aligned} p_{\text{cond}}^{\text{cor}} &\leq \exp_2\left(-E - \frac{1}{2} \log_2(\varepsilon) + 2\eta \log_2\left(\frac{1}{\eta}\right)N + O(\log_2 N)\right) \\ &\leq \exp_2\left(-E + \frac{E}{4} + \frac{E}{4} + O(\log_2 N)\right) = \exp_2\left(-\frac{E}{2} + O(\log_2 N)\right) = o(1). \end{aligned}$$

where the last equality follows as $E \gg \log_2 N$. Then, by [Proposition 2.20](#), the choice of $D = o(\exp_2(E/4))$ gives

$$\begin{aligned} p_{\text{unstable}}^{\text{cor}} &\leq \frac{CD\varepsilon}{2\eta} \asymp \frac{D\varepsilon N \log_2(N/E)}{E} \\ &= \frac{D \exp_2(-E/2) N \log_2(N/E)}{E} \leq \frac{D \exp_2(-E/2) N \log_2(N)}{E} \\ &\leq D \exp_2\left(-\frac{E}{2} + \log_2(N) + \log_2 \log_2(N) - \log_2(E)\right) \\ &\leq \exp_2\left(-\frac{E}{4} + \log_2(N) + \log_2 \log_2(N) - \log_2(E)\right) = o(1), \end{aligned}$$

again, as $E \gg \log_2 N$. Ergo, by [\(3.4\)](#), $(p_{\text{solve}}^{\text{cor}})^2 \leq p_{\text{unstable}}^{\text{cor}} + p_{\text{cond}}^{\text{cor}} = o(1)$, as desired. \square

3.2 Proof for Low Coordinate-Degree Algorithms

Next, let \mathcal{A} have coordinate degree D . We now want g, g' to be $(1 - \varepsilon)$ -resampled standard Normals. We define the following events.

$$\begin{aligned} S_{\text{diff}} &= \{g \neq g'\} \\ S_{\text{solve}} &= \{\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g')\} \\ S_{\text{stable}} &= \{\|\mathcal{A}(g) - \mathcal{A}(g')\| \leq 2\sqrt{\eta N}\} \\ S_{\text{cond}}(x) &= \left\{ \nexists x' \in S(E; g') \text{ such that } \|x - x'\| \leq 2\sqrt{\eta N} \right\} \end{aligned} \quad (3.8)$$

Note that these are the same events as [\(3.1\)](#), along with an event to ensure that g' is nontrivially resampled from g .

Lemma 3.7. *For g, g' being $(1 - \varepsilon)$ -resampled, $\mathbf{P}(S_{\text{diff}}) = 1 - (1 - \varepsilon)^N \leq \varepsilon N$.*

Proof: Follows from calculation:

$$\mathbf{P}(g = g') = \prod_{i=1}^N \mathbf{P}(g_i = g_{i'}) = (1 - \varepsilon)^N \quad \square$$

Lemma 3.8. *We have, for $x = \mathcal{A}(g)$, $S_{\text{diff}} \cap S_{\text{solve}} \cap S_{\text{stable}} \cap S_{\text{cond}}(x) = \emptyset$.*

Proof: This follows from [Lemma 3.1](#), noting that the proof did not use that $g \neq g'$ almost surely. \square

We should interpret this as saying $S_{\text{solve}}, S_{\text{stable}}, S_{\text{cond}}$ are all mutually exclusive, conditional on $g \neq g'$.

The previous definition of $p_{\text{solve}}^{\text{cor}}$ in [\(3.2\)](#), which we now term $p_{\text{solve}}^{\text{res}}$, remains valid. In particular, we have

Lemma 3.9. *For g, g' being $(1 - \varepsilon)$ -resampled, we have*

$$\mathbf{P}(S_{\text{solve}}) = \mathbf{P}(\mathcal{A}(g) \in S(E; g), \mathcal{A}(g') \in S(E; g')) \geq (p_{\text{solve}}^{\text{res}})^2$$

Proof: Let $\tilde{g}, g^{(0)}, g^{(1)}$ be three i.i.d. copies of g , and let J be a random subset of $[N]$ where each coordinate is included with probability $1 - \varepsilon$. Then, g, g' are jointly representable as

$$g = \tilde{g}_J + g_{[N] \setminus J}^{(0)}, \quad g' = \tilde{g}_J + g_{[N] \setminus J}^{(1)}$$

where \tilde{g}_J denotes the vector with coordinates \tilde{g}_i if $i \in J$ and 0 else. Thus g and g' are conditionally independent, given (\tilde{g}, J) , and the proof concludes as in [Lemma 3.2](#). \square

Let us slightly redefine $p_{\text{unstable}}^{\text{res}}$ and $p_{\text{cond}}^{\text{res}}(x)$ by

$$p_{\text{unstable}}^{\text{res}} = 1 - \mathbf{P}(S_{\text{stable}} \mid S_{\text{diff}}), \quad p_{\text{cond}}^{\text{res}}(x) = 1 - \mathbf{P}(S_{\text{cond}}(x) \mid S_{\text{diff}}). \quad (3.9)$$

This is necessary as when $g = g'$, S_{stable} always holds and $S_{\text{cond}}(x)$ always fails. Note however that if we knew that $\mathbf{P}(S_{\text{diff}}) = 1$, which is always the case for g, g' being $(1 - \varepsilon)$ -correlated, these definitions agree with what we had in [\(3.4\)](#). Again, we can define $p_{\text{cond}}^{\text{res}}$ via [\(3.3\)](#), i.e. as the maximum of $p_{\text{cond}}^{\text{res}}(x)$ over Σ_N .

Now, by [Lemma 3.8](#), we know that for $x = \mathcal{A}(g)$, $\mathbf{P}(S_{\text{solve}}, S_{\text{stable}}, S_{\text{cond}}(x) \mid S_{\text{diff}}) = 0$, so

$$\mathbf{P}(S_{\text{solve}} \mid S_{\text{diff}}) + \mathbf{P}(S_{\text{stable}} \mid S_{\text{diff}}) + \mathbf{P}(S_{\text{cond}}(x) \mid S_{\text{diff}}) \leq 2.$$

Thus, rearranging and multiplying by $\mathbf{P}(S_{\text{diff}})$ (so as to apply [Lemma 3.9](#)) gives

$$(p_{\text{solve}}^{\text{res}})^2 \leq \mathbf{P}(S_{\text{diff}}) \cdot (p_{\text{unstable}}^{\text{res}} + p_{\text{cond}}^{\text{res}}) \quad (3.10)$$

As before, our proof follows by showing that, for appropriate choices of ε and η , depending on D, E , and N , that $p_{\text{unstable}}^{\text{res}}, p_{\text{cond}}^{\text{res}} = o(1)$. However, this also requires us to choose $\varepsilon \gg \frac{1}{N}$, so as to ensure that $g \neq g'$, as otherwise $p_{\text{unstable}}^{\text{res}}, p_{\text{cond}}^{\text{res}}$ would be too large. This restriction on ε effectively limits us from showing hardness for algorithms with degree larger than $o(N)$, as we will see shortly.

First, we bound the same probability of a fixed x solving a resampled instance. Here, we need to condition on the resampled instance being different, as otherwise the probability in question can be made to be 1 if x was chosen to solve g .

Proposition 3.10 (Fundamental Estimate – Resampled Case). *Assume that (g, g') are $(1 - \varepsilon)$ -resampled standard Normal vectors. Then, for any x only depending on g ,*

$$p_{\text{cond}}^{\text{res}}(x) = \mathbf{P} \left(\exists x' \in S(E; g') \text{ such that } \left| \begin{array}{l} \|x - x'\| \leq 2\sqrt{\eta N} \\ g \neq g' \end{array} \right. \right) \leq \exp_2 \left(-E + 2\eta \log_2 \left(\frac{1}{\eta} \right) N + O(1) \right).$$

Proof: We follow the setup of proof of [Proposition 3.4](#). For each x' within distance $2\sqrt{\eta N}$ of x , let

$$I_{x'} := I(x \in S(E; g')) = I(|\langle g', x' \rangle| \leq 2^{-E}),$$

so that

$$\begin{aligned} p_{\text{cond}}^{\text{res}}(x) &= \mathbf{E} \left[\sum_{\|x - x'\| \leq 2\sqrt{\eta N}} \mathbf{E}[I_{x'} \mid g, g \neq g'] \right] \\ &= \mathbf{E} \left[\sum_{\|x - x'\| \leq 2\sqrt{\eta N}} \mathbf{P}(|\langle g', x' \rangle| \leq 2^{-E} \mid g, g \neq g') \mid g \neq g' \right] \end{aligned} \quad (3.11)$$

Again, to bound the inner probability, let \tilde{g} be a Normal vector independent to g . Let $J \subseteq [N]$ be a random subset where each $i \in J$ with probability $1 - \varepsilon$, independently, so g' can be represented as $g' = g_J + \tilde{g}_{[N] \setminus J}$. For a fixed x' and conditional on (g, J) , we know that $\langle \tilde{g}_{[N] \setminus J}, x' \rangle$ is $\mathcal{N}(0, N - |J|)$ and $\langle g_J, x' \rangle$ is deterministic. That is,

$$\langle g', x' \rangle \mid (g, J) \sim \mathcal{N}(\langle g_J, x' \rangle, N - |J|).$$

Conditioning on $g \neq g'$ is equivalent to conditioning on $|J| < N$, so $N - |J| \geq 1$. Thus, applying [Lemma 1.2](#) and integrating over all valid choices of J gives

$$\mathbf{P}(|\langle g', x' \rangle| \leq 2^{-E} \mid g, g \neq g') \leq \exp_2(-E + O(1)). \quad (3.12)$$

By [Proposition 3.3](#), the number of terms in the sum (3.11) is bounded by $\exp_2(2\eta \log_2(1/\eta)N)$, so summing (3.12) allows us to conclude that

$$p_{\text{cond}}^{\text{res}}(x) \leq \exp_2 \left(-E + 2\eta \log_2 \left(\frac{1}{\eta} \right) N + O(1) \right). \quad \square$$

Note that in contrast to [Proposition 3.4](#), this bound doesn't involve ε at all, but the condition $g \neq g'$ requires $\varepsilon = \omega(1/N)$ to hold almost surely, by [Lemma 3.7](#).

With this, we can show strong low degree hardness for low coordinate degree algorithms at energy levels $E = \Theta(N)$.

Theorem 3.11. *Let $\delta > 0$ and $E = \delta N$, and let g, g' be $(1 - \varepsilon)$ -resampled standard Normal r.v.s. Then, for any algorithm \mathcal{A} with coordinate degree $D \leq o(N)$ and $\mathbf{E}\|\mathcal{A}(g)\|^2 \leq CN$, there exist $\varepsilon, \eta > 0$ such that $p_{\text{solve}}^{\text{res}} = o(1)$.*

Proof: Recall from (3.10) that it suffices to show that both $p_{\text{cond}}^{\text{res}}$ and $p_{\text{unstable}}^{\text{res}}$ go to zero, while $\mathbf{P}(S_{\text{diff}}) \approx 1$. By [Lemma 3.7](#), the latter condition is satisfied for $\varepsilon = \omega(1/N)$. Thus, pick

$$\varepsilon = \frac{\log_2(N/D)}{N}. \quad (3.13)$$

Note that this satisfies $N\varepsilon = \log_2(N/D) \gg 1$, for $D = o(N)$. Next, choose η such that $2\eta \log_2(1/\eta) < \delta/4$ —again, this results in η being independent of N . As the bound in [Proposition 3.10](#) is independent of x , we get

$$p_{\text{cond}}^{\text{res}} \leq \exp_2\left(-\delta N + \frac{\delta N}{4} + O(1)\right) = o(1).$$

Moreover, for $D \leq o(N)$, [Proposition 2.20](#) now gives

$$p_{\text{unstable}}^{\text{res}} \leq \frac{CD\varepsilon}{2\eta} \asymp D \cdot \frac{\log_2(N/D)}{N} \rightarrow 0,$$

as $x \log_2(1/x) \rightarrow 0$ for $x \ll 1$. By [\(3.10\)](#), we conclude that $(p_{\text{solve}}^{\text{res}})^2 \leq \mathbf{P}(S_{\text{diff}}) \cdot (p_{\text{unstable}}^{\text{res}} + p_{\text{cond}}^{\text{res}}) = o(1)$, thus completing the proof. \square

Sublinear case. We now consider sublinear energy levels, ranging from $(\log_2 N)^2 \ll E \ll N$. Note here that we have to increase our lower bound to $(\log_2 N)^2$ as opposed to $\log_2 N$ from [Theorem 3.6](#), to address the requirement that $\varepsilon = \omega(1/N)$.

Theorem 3.12. *Let $\omega((\log_2 N)^2) \leq E \leq o(N)$, and let g, g' be $(1 - \varepsilon)$ -resampled standard Normal r.v.s. Then, for any algorithm \mathcal{A} with coordinate degree $D \leq o(E/(\log_2 N)^2)$ and $\mathbf{E}\|\mathcal{A}(g)\|^2 \leq CN$, there exist $\varepsilon, \eta > 0$ such that $p_{\text{solve}}^{\text{res}} = o(1)$.*

Proof: As in [Theorem 3.11](#), choose ε as in [\(3.13\)](#), so that $\varepsilon = \omega(1/N)$ and $\mathbf{P}(S_{\text{diff}}) \approx 1$. However, to account for $E \leq o(N)$, we need to adjust η as $N \rightarrow \infty$. Thus, choose η as in [\(3.7\)](#): this ensures that $\varepsilon = \omega(1/N)$ and that $2\eta \log_2(1/\eta) < E/4N$ for $E \ll N$. By [Proposition 3.10](#), this guarantees that

$$p_{\text{cond}}^{\text{res}} \leq \exp_2\left(-E + 2\eta \log_2\left(\frac{1}{\eta}\right)N + O(1)\right) \leq \exp_2\left(-\frac{3E}{4} + O(1)\right) = o(1).$$

The low coordinate degree requirement $D \leq o(E/(\log_2 N)^2)$ plus [Proposition 2.20](#) now gives

$$\begin{aligned} p_{\text{unstable}}^{\text{res}} &\leq \frac{CD\varepsilon}{2\eta} \asymp \frac{D\varepsilon N \log_2(N/E)}{E} \\ &= \frac{D \log_2(N/D) \log_2(N/E)}{E} \leq \frac{D(\log_2 N)^2}{E} = o(1). \end{aligned}$$

By [\(3.10\)](#), $(p_{\text{solve}}^{\text{res}})^2 \leq \mathbf{P}(S_{\text{diff}}) \cdot (p_{\text{unstable}}^{\text{res}} + p_{\text{cond}}^{\text{res}}) = o(1)$, thus completing the proof. \square

3.3 Summary of Parameters

Parameter	Meaning	Desired Direction	Intuition
N	Dimension	Large	Showing hardness <i>asymptotically</i> , want “bad behavior” to pop up in low dimensions.
E	Solution energy; want to find x such that $ \langle g, x \rangle \leq 2^{-E}$	Small	Smaller E implies weaker solutions, and can consider full range of $1 \ll E \ll N$. Know that $E > (\log^2 N)$ by (Karmarkar and Karp 1983)
D	Algorithm degree (in either Efron-Stein sense or usual polynomial sense.)	Large	Higher degree means more complexity. Want to show even complex algorithms fail.
ε	Complement of correlation/resample probability; (g, g') are $(1 - \varepsilon)$ -correlated.	Small	ε is “distance” between g, g' . Want to show that small changes in disorder lead to “breaking” of landscape.
η	Algorithm instability; \mathcal{A} is stable if $\ \mathcal{A}(g) - \mathcal{A}(g')\ \leq 2\sqrt{\eta N}$, for (g, g') close.	Large	Large η indicates a more unstable algorithm; want to show that even weakly stable algorithms fail.

Table 1: Explanation of Parameters

4 Extensions to Real-Valued Algorithms

With [Section 3](#), we have established strong low degree hardness for both low degree polynomial algorithms and low coordinate degree algorithms. However, our stability analysis assumed that the algorithms in question were Σ_N -valued. In this section, we show that this assumption is not in fact as restrictive as it might appear.

Throughout, let \mathcal{A} denote an \mathbf{R}^N -valued algorithm. We want to show that

- I. No low degree \mathcal{A} can reliably output points *close* – within constant distance – to a solution,
- II. No Σ_N -valued algorithm $\tilde{\mathcal{A}}$ coming from randomly rounding the output of \mathcal{A} , which changes an $\omega(1)$ number of coordinates, can find a solution with nonvanishing probability.

In principle, the first possibility fails via the same analysis as in [Section 3](#), while the second fails because the landscape of solutions to any given NPP instance is sparse.

Why are these the only two possibilities? For \mathcal{A} to provide a way to actually solve the NPP, we must be able to turn its outputs on \mathbf{R}^N into points on Σ_N . If \mathcal{A} could output points within an constant distance (independent of the instance) of a solution, then we could convert \mathcal{A} into a Σ_N -valued algorithm by manually computing the energy of all points close to its output and returning the energy-maximizing point.

However, the more common way to convert a \mathbf{R}^N -valued algorithm into a Σ_N -valued one is by rounding the outputs, as in [\(Huang and Sellke 2025\)](#). Doing this directly can lead to difficulties in performing the stability analysis. In our case, though, if we know no \mathcal{A} can reliably output points within constant distance of a solution, then any rounding scheme which only flips $O(1)$ many coordinates will assuredly fail. Thus, the only rounding schemes worth considering are those which flip $\omega(1)$ many coordinates.

We first describe a landscape obstruction to finding multiple solutions at the same energy level for a random NPP instance. Then, we show hardness in both of the aforementioned cases. meow.

4.1 Solutions repel meow

Introduce section meow.

No two adjacent points on Σ_N (or pairs within $k = O(1)$ distance) which are both good solutions to the same problem.

Proposition 4.1. *Fix distinct points $x, x' \in \Sigma_N$ and let $g \sim \mathcal{N}(0, I_N)$ be a random instance. Then,*

$$\mathbf{P}(x, x' \in S(E; g)) \leq \exp_2(-E + O(1)) = \exp_2(-E + O(1)).$$

Proof: For $x \neq x'$, let $J \subseteq [N]$ denote the subset of coordinates in which x, x' differ, i.e. $x_J \neq x'_J$. In particular, we can write

$$x = x_{[N] \setminus J} + x_J, \quad x' = x_{[N] \setminus J} - x_J.$$

Thus, for a fixed pair (x, x') , if $-2^{-E} \leq \langle g, x \rangle, \langle g, x' \rangle \leq 2^{-E}$, we can expand this into

$$\begin{aligned} -2^{-E} &\leq \langle g, x_{[N] \setminus J} \rangle + \langle g, x_J \rangle \leq 2^{-E}, \\ -2^{-E} &\leq \langle g, x_{[N] \setminus J} \rangle - \langle g, x_J \rangle \leq 2^{-E}. \end{aligned}$$

Multiplying the lower equation by -1 and adding the resulting inequalities gives $|\langle g, x_J \rangle| \leq 2^{-E}$. Note that $\langle g, x_J \rangle \sim \mathcal{N}(0, |J|)$ (and is nondegenerate, as $|J| > 0$). By [Lemma 1.2](#) and the following remark, it follows that

$$\mathbf{P}(x, x' \in S(E; g)) \leq \mathbf{P}(|\langle g, x_J \rangle| \leq 2^{-E}) \leq \exp_2(-E + O(1)). \quad \square$$

Remarks on theorem below meow.

Theorem 4.2 (Solutions Can't Be Close). *Consider any distances $k = \Omega(1)$ and energy levels $E \gg k \log_2 N$. Then for any instance g , there are no pairs of distinct solutions $x, x' \in S(E; g)$ with $\|x - x'\| \leq 2\sqrt{k}$ (i.e. within k coordinate flips of each other) with high probability.*

Proof: Observe that by [Proposition 4.1](#), finding a pair of distinct solutions within distance $2\sqrt{k}$ implies finding some subset of at most k coordinates $J \subset [N]$ of g and $|J|$ signs x_J such that $|\langle g_J, x_J \rangle|$ is small. For any g , there are at most 2^k choices of signs and, by ([Vershynin 2018, Exer. 0.0.5](#)), there are

$$\sum_{1 \leq k' \leq k} \binom{N}{k'} \leq \left(\frac{eN}{k} \right)^k \leq (eN)^k = 2^{O(k \log_2 N)}$$

choices of such subsets. Union bounding [Proposition 4.1](#) over these $\exp_2 O(k \log_2 N)$ choices, we get

$$\mathbf{P} \left(\begin{array}{l} \exists x, x' \text{ s.t.} \\ \text{(a) } \|x - x'\| \leq 2\sqrt{k}, \\ \text{(b) } x, x' \in S(E; g) \end{array} \right) \leq \mathbf{P} \left(\begin{array}{l} \exists J \subset [N], x_J \in \{\pm 1\}^{|J|} \text{ s.t.} \\ \text{(a) } |J| \leq k, \\ \text{(b) } |\langle g_J, x_J \rangle| \leq \exp_2(-E) \end{array} \right) \leq \exp_2(-E + O(k \log_2 N)) = o(1).$$

Note that the last equality holds as $E \gg k \log_2 N$. \square

4.2 Proof of Hardness for Close Algorithms

Throughout this section, fix some distance $r = O(1)$. Consider the event that the \mathbf{R}^N -valued \mathcal{A} outputs a point close to a solution for an instance g :

$$S_{\text{close}}(r) = \left\{ \begin{array}{l} \exists \hat{x} \in S(E; g) \text{ s.t.} \\ \mathcal{A}(g) \in B(\hat{x}, r) \end{array} \right\} = \{B(\mathcal{A}(g), r) \cap S(E; g) \neq \emptyset\}$$

Note that as r is fixed (potentially depending on \mathcal{A} , but independent of N or g), we can convert \mathcal{A} into a Σ_N -valued algorithm by considering the corners of Σ_N within constant distance of $\mathcal{A}(g)$.

Definition 4.3. Let $r > 0$ and \mathcal{A} be an \mathbf{R}^N -valued algorithm. Define $\widehat{\mathcal{A}}_r$ to be the Σ_N -valued algorithm defined by

$$\widehat{\mathcal{A}}_r(g) := \operatorname{argmin}_{x' \in B(\mathcal{A}(g), r) \cap \Sigma_N} |\langle g, x' \rangle|. \quad (4.2)$$

If $B(\mathcal{A}(g), r) \cap \Sigma_N = \emptyset$, then set $\widehat{\mathcal{A}}_r(g) := (1/g_1, 0, \dots)$, which always has energy 0.

Observe that $S_{\text{close}(r)}$ occurring is the same as $\widehat{\mathcal{A}}_r$ finding a solution for g . In addition, note that practically speaking, computing $\widehat{\mathcal{A}}_r$ requires additionally computing the energy of $O(1)$ -many points on Σ_N . This requires only an additional $O(N)$ operations.

Recall from [Section 2.3](#) that if \mathcal{A} is low degree (or low coordinate degree) then we can derive useful stability bounds for its outputs. Luckily, this modification $\widehat{\mathcal{A}}_r$ of \mathcal{A} also are stable, with slightly modified bounds.

Lemma 4.4. *Suppose that $\mathbf{E}\|\mathcal{A}(g)\|^2 \leq CN$ and that \mathcal{A} has degree $\leq D$ (resp. coordinate degree $\leq D$), and let (g, g') be $(1 - \varepsilon)$ -correlated (resp. $(1 - \varepsilon)$ -resampled). Then $\widehat{\mathcal{A}}_r$ as defined above has*

$$\mathbf{E}\|\widehat{\mathcal{A}}_r(g) - \widehat{\mathcal{A}}_r(g')\|^2 \leq 6CD\varepsilon N + 6r^2.$$

In particular, we have

$$\mathbf{P}\left(\|\widehat{\mathcal{A}}_r(g) - \widehat{\mathcal{A}}_r(g')\| \geq 2\sqrt{\eta N}\right) \leq \frac{3CD\varepsilon}{2\eta} + \frac{3r^2}{2\eta N}. \quad (4.3)$$

Proof: Observe by the triangle inequality, as per [\(1.2\)](#), that

$$\|\widehat{\mathcal{A}}_r(g) - \widehat{\mathcal{A}}_r(g')\|^2 \leq 3\left(\|\widehat{\mathcal{A}}_r(g) - \mathcal{A}(g)\|^2 + \|\mathcal{A}(g) - \mathcal{A}(g')\|^2 + \|\mathcal{A}(g') - \widehat{\mathcal{A}}_r(g')\|^2\right).$$

By [Proposition 2.20](#), we know $\mathbf{E}\|\mathcal{A}(g) - \mathcal{A}(g')\|^2 \leq 6CD\varepsilon N$. Moreover, we know that $\|\widehat{\mathcal{A}}_r(g) - \mathcal{A}(g)\| \leq r$ by definition, so the remaining terms can be bounded by $3r^2$ each deterministically. Finally, [\(4.2\)](#) follows from Markov's inequality. \square

Of course, computing $\widehat{\mathcal{A}}_r$ is certainly never polynomial, and does not preserve any low coordinate degree assumptions in a controllable way. Thus, we cannot directly hope for [Theorem 3.5](#), [Theorem 3.6](#), [Theorem 3.11](#), or [Theorem 3.12](#) to hold meow

We show for \mathcal{A} being a \mathbf{R}^N -valued, low coordinate degree algorithm and any $r = O(1)$, low degree hardness still holds for $\widehat{\mathcal{A}}_r$. Note that by a similar argument, we can show hardness in the case that \mathcal{A} is a low degree polynomial algorithm, but we omit the proof meow.

We recall the setup from [Section 3.2](#). Let g, g' be $(1 - \varepsilon)$ -resampled standard Normal vectors. Define the following events:

$$\begin{aligned} S_{\text{diff}} &= \{g \neq g'\} \\ S_{\text{solve}} &= \{\widehat{\mathcal{A}}_r(g) \in S(E; g), \widehat{\mathcal{A}}_r(g') \in S(E; g')\} \\ S_{\text{stable}} &= \{\|\widehat{\mathcal{A}}_r(g) - \widehat{\mathcal{A}}_r(g')\| \leq 2\sqrt{\eta N}\} \\ S_{\text{cond}}(x) &= \left\{ \nexists x' \in S(E; g') \text{ such that } \|x - x'\| \leq 2\sqrt{\eta N} \right\} \end{aligned} \quad (4.4)$$

These are the same events as in [\(3.8\)](#), just adapted to $\widehat{\mathcal{A}}_r$. In particular, [Lemma 3.8](#) holds unchanged.

Moreover, we can define

$$\hat{p}_{\text{solve}}^{\text{cor}} = \mathbf{P}\left(\widehat{\mathcal{A}}_r(g) \in S(E; g)\right) = \mathbf{P}(S_{\text{close}}(r)), \quad (4.5)$$

as well as

$$\hat{p}_{\text{unstable}}^{\text{cor}} = 1 - \mathbf{P}(S_{\text{stable}} \mid S_{\text{diff}}), \quad \hat{p}_{\text{cond}}^{\text{cor}}(x) = 1 - \mathbf{P}(S_{\text{cond}}(x) \mid S_{\text{diff}}),$$

along with $\hat{p}_{\text{cond}}^{\text{cor}} := \max_{x \in \Sigma_N} \hat{p}_{\text{cond}}^{\text{cor}}(x)$, echoing (3.9).

Observe that as $\hat{p}_{\text{cond}}^{\text{cor}}$ makes no reference to any algorithm, the bound in [Proposition 3.10](#) holds without change. Moreover, [Lemma 4.4](#) lets us control $\hat{p}_{\text{unstable}}^{\text{cor}}$. The final piece needed is an appropriate analog of [Lemma 3.9](#).

Lemma 4.5. *For g, g' being $(1 - \varepsilon)$ -resampled, we have*

$$\mathbf{P}(S_{\text{solve}}) = \mathbf{P}(\widehat{\mathcal{A}}_r(g) \in S(E; g), \widehat{\mathcal{A}}_r(g') \in S(E; g')) \geq (\hat{p}_{\text{solve}}^{\text{cor}})^2$$

Proof: Observe that, letting $+$ denote Minkowski sum, we have that

$$\{\widehat{\mathcal{A}}_r(g) \in S(E; g)\} = \{\mathcal{A}(g) \in S(E; g) + B(0, r)\}.$$

Expanding $S(E; g)$, the proof concludes as in [Lemma 3.9](#). \square

Theorem 4.6. *Let $\omega((\log_2 N)^2) \leq E \leq \Theta(N)$, and let g, g' be $(1 - \varepsilon)$ -resampled standard Normal r.v.s. Consider any $r = O(1)$ and \mathbf{R}^N -valued \mathcal{A} with $\mathbf{E}\|\mathcal{A}(g)\|^2 \leq CN$, and assume in addition that*

- (a) *if $E = \delta N = \Theta(N)$ for $\delta > 0$, then \mathcal{A} has coordinate degree $D \leq o(N)$;*
- (b) *if $(\log_2 N)^2 \ll E \ll N$, then \mathcal{A} has coordinate degree $D \leq o(E/(\log_2 N)^2)$.*

Let $\widehat{\mathcal{A}}_r$ be defined as in [Definition 4.3](#). Then there exist $\varepsilon, \eta > 0$ such that

$$\hat{p}_{\text{solve}}^{\text{cor}} = \mathbf{P}(\widehat{\mathcal{A}}_r(g) \in S(E; g)) = o(1).$$

Proof: First, by [Lemma 3.8](#), the appropriate adjustment of (3.10) holds, namely that

$$(\hat{p}_{\text{solve}}^{\text{cor}})^2 \leq \mathbf{P}(S_{\text{diff}}) \cdot (\hat{p}_{\text{unstable}}^{\text{cor}} + \hat{p}_{\text{cond}}^{\text{cor}}). \quad (4.6)$$

To ensure $\mathbf{P}(S_{\text{diff}}) \approx 1$, we begin by following (3.13) and choosing $\varepsilon = \log_2(N/D)/N$. Moreover, following the proof of [Theorem 3.11](#) and [Theorem 3.12](#), we know that choosing

$$\eta = \begin{cases} O(1) \text{ s.t. } 2\eta \log_2(1/\eta) < \delta/4 & E = \delta N, \\ \frac{E}{16N \log_2(N/E)} & E = o(N) \end{cases}$$

in conjunction with [Proposition 3.10](#), guarantees that

$$\hat{p}_{\text{cond}}^{\text{cor}} \leq \exp_2\left(-\frac{3E}{4} + O(1)\right) = o(1).$$

Finally, note that in the linear case, when $\eta = O(1)$, $\frac{r^2}{\eta N} = o(1)$ trivially. In the sublinear case, for $\eta = E/(16N \log_2(N/E))$, we instead get

$$\eta N = \frac{E}{16 \log_2(N/E)} \geq \frac{E}{16 \log_2 N} = \omega(1),$$

as $E \gg (\log_2 N)^2$. Thus, applying the properly modified [Lemma 4.4](#) with these choices of ε, η , we see that $\hat{p}_{\text{unstable}}^{\text{cor}} = o(1)$. By (4.6), we conclude that $\hat{p}_{\text{solve}}^{\text{cor}} = o(1)$, as desired. \square

Talk about implications meow.

4.3 Truly Random Rounding

At this point, one might wonder whether, while deterministic algorithms fail, perhaps a randomized rounding scheme could save us, maybe by assigning small values to coordinates which it was less confident in. However, this approach is blunted by the same brittleness of the NPP landscape that established the conditional obstruction of [Proposition 3.4](#) and [Proposition 3.10](#). In particular, by [Theorem 4.2](#), if you have a subcube of Σ_N nonconstant but bounded dimension, then with high probability at most one of those points will be a solution.

For this section, again let \mathcal{A} be a deterministic \mathbf{R}^N -valued algorithm. Moreover, assume we are searching for solutions with energy between $(\log_2 N)^2 \ll E \leq N$; note that for lower values of E , algorithms like [\(Karmarkar and Karp 1983\)](#) already achieve discrepancies of $N^{O(\log_2 N)}$ energy in polynomial time.

To start, for any $x \in \mathbf{R}^N$, we write x^* for the coordinate-wise signs of x , i.e.

$$x_i^* := \begin{cases} +1 & x_i > 0, \\ -1 & x_i \leq 0. \end{cases}$$

We can define the modified alg $\mathcal{A}^*(g) := \mathcal{A}(g)^*$.

Remark 4.7. meow \mathcal{A}^* fails and is still degree D lcdf, even if it stops being a polynomial. Bounds on D worsen, but only to what you'd expect. Note that if \mathcal{A} has coordinate degree D , then \mathcal{A}^* also has coordinate degree D . As a deterministic Σ_N -valued algorithm, strong low degree hardness as proved in the previous section applies.

In contrast to deterministically taking signs of the outputs of \mathcal{A} (which corresponds to deterministically rounding the outputs of \mathcal{A} to Σ_N), we can consider passing the output of \mathcal{A} through a randomized rounding scheme. Let $\text{round}(x, \omega) : \mathbf{R}^N \times \Omega \rightarrow \Sigma_N$ denote any randomized rounding function, with randomness ω independent of the input. We will often suppress the ω in the notation, and treat $\text{round}(x)$ as a Σ_N -valued random variable. Given such a randomized rounding function, we can describe it in the following way. Let $p_1(x), \dots, p_N(x)$ be defined by

$$p_i(x) := \max\left(\mathbf{P}(\text{round}(x)_i \neq x_i^*), \frac{1}{2}\right). \quad (4.7)$$

We need to guarantee that each $p_i(x) \leq 1/2$ for the following alternative description of $\text{round}(x)$:

Lemma 4.8. Fix $x \in \mathbf{R}^N$, and draw N coin flips $I_{x,i} \sim \text{Bern}(2p_i(x))$ and N signs $S_i \sim \text{Unif}\{\pm 1\}$, all mutually independent, and define the random variable $\tilde{x} \in \Sigma_N$ by

$$\tilde{x}_i := S_i I_{x,i} + (1 - I_{x,i}) x_i^*.$$

Then $\tilde{x} \sim \text{round}(x)$.

Proof: Conditioning on $I_{x,i}$, we can check that

$$\mathbf{P}(\tilde{x}_i \neq x_i) = 2p_i(x) \cdot \mathbf{P}(\tilde{x}_i = x_i \mid I_{x,i} = 1) + (1 - 2p_i(x)) \cdot \mathbf{P}(\tilde{x}_i \neq x_i \mid I_{x,i} = 0) = p_i(x).$$

Thus, $\mathbf{P}(\tilde{x}_i = x_i^*) = \mathbf{P}(\text{round}(x)_i = x_i^*)$, as claimed. \square

By [Lemma 4.8](#), we can redefine $\text{round}(x)$ to be \tilde{x} as constructed without loss of generality.

It thus makes sense to define $\tilde{\mathcal{A}}(g) := \text{round}(\mathcal{A}(g))$, which is now (a) Σ_N -valued and (b) randomized only in the transition from \mathbf{R}^N to Σ_N (i.e., the rounding doesn't depend directly on g , only the output $x = \mathcal{A}(g)$). We should expect that if $\tilde{\mathcal{A}} = \mathcal{A}^*$ (e.g., if \mathcal{A} outputs values far outside the cube $[-1, 1]^N$) with high probability, then low degree hardness will still apply, as \mathcal{A}^* is deterministic. However, in general, any $\tilde{\mathcal{A}}$ which differs from \mathcal{A}^* will fail to solve g with high probability. This is independent of any assumptions on \mathcal{A} : any rounding scheme will introduce so much randomness that \tilde{x} will be effectively a random point, which has a vanishing probability of being a solution because of how sparse and disconnected the NPP landscape is.

To see this, we first show that any randomized rounding scheme as in [Lemma 4.8](#) which a.s. differs from picking the closest point deterministically will resample a diverging number of coordinates.

Lemma 4.9. *Fix $x \in \mathbf{R}^N$, and let $p_1(x), \dots, p_N(x)$ be defined as in [\(4.7\)](#). Then $\tilde{x} \neq x^*$ with high probability iff $\sum_i p_i(x) = \omega(1)$. Moreover, assuming that $\sum_i p_i(x) = \omega(1)$, the number of coordinates in which \tilde{x} is resampled diverges almost surely.*

Proof: Recall that for $x \in [0, 1/2]$, $\log_2(1 - x) = \Theta(x)$. Thus, as each coordinate of x is rounded independently, we can compute

$$\mathbf{P}(\tilde{x} = x^*) = \prod_i (1 - p_i(x)) = \exp_2 \left(\sum_i \log_2(1 - p_i(x)) \right) \leq \exp_2 \left(-\Theta \left(\sum_i p_i(x) \right) \right).$$

Thus, $\mathbf{P}(\tilde{x} = x^*) = o(1)$ iff $\sum_i p_i(x) = \omega(1)$, as claimed.

Next, following the construction of \tilde{x} in [Lemma 4.8](#), let $E_i = \{I_{x,i} = 1\}$ be the event that \tilde{x}_i is resampled from $\text{Unif}\{\pm 1\}$, independently of x_i^* . The E_i are independent, so by Borel-Cantelli, $\sum_i \mathbf{P}(E_i) = 2 \sum_i p_i(x) = \omega(1)$ implies \tilde{x}_i is resampled infinitely often with probability 1, thus completing the proof. \square

With this, we can apply [Theorem 4.2](#), which shows that solutions resist clustering at a rate related to their energy level (i.e. higher energy solutions are pushed further apart), to conclude that any $\tilde{\mathcal{A}}$ which is not equal to \mathcal{A}^* with high probability fails to find solutions.

Theorem 4.10. *Let $x = \mathcal{A}(g)$, and define x^*, \tilde{x} , etc., as previously. Moreover, assume that for any x in the possible outputs of \mathcal{A} , we have $\sum_i p_i(x) = \omega(1)$. Then, for any $E \geq \omega((\log_2 N)^2)$, we have*

$$\mathbf{P}(\tilde{\mathcal{A}}(g) \in S(E; g)) = \mathbf{P}(\tilde{x} \in S(E; g)) \leq o(1).$$

Proof: Following the characterization of \tilde{x} in [Lemma 4.8](#), let $K := \max(\log_2 N, \sum_i I_{x,i})$. By assumption on $\sum_i p_i(x)$ and [Lemma 4.9](#), we know K , which is at least the number of coordinates which are resampled, is bounded as $1 \ll K \leq \log_2 N$, for any possible $x = \mathcal{A}(g)$. Now, let $S \subseteq [N]$ denote the set of the first K coordinates to be resampled, so that $K = |S|$. Consider now

$$\mathbf{P}(\tilde{x} \in S(E; g) \mid \tilde{x}_{[N] \setminus S}),$$

where we fix the coordinates outside of S and let \tilde{x} be uniformly sampled from a K -dimensional subcube of Σ_N . All such \tilde{x} are within distance $2\sqrt{K}$ of each other, so by **Theorem 4.2**, the probability there is more than one such $\tilde{x} \in S(E; g)$ is bounded by

$$\exp_2(-E + O(K \log_2 N)) \leq \exp_2(-E + O((\log_2 N)^2)) = o(1),$$

by assumption on E . Thus, the probability that any of the \tilde{x} is in $S(E; g)$ is bounded by 2^{-K} , whence

$$\mathbf{P}(\tilde{x} \in S(E; g)) = \mathbf{E}[\mathbf{P}(\tilde{x} \in S(E; g) \mid \tilde{x}_{[N] \setminus S})] \leq 2^{-K} \leq o(1). \quad \square$$

meow discuss possible extensions of randomization schemes and whether you expect those to work instead.

5 Conclusion

Meow

5.1 Future Work

Bibliography

- Achlioptas, Dimitris, and Amin Coja-Oghlan. 2008. “Algorithmic Barriers from Phase Transitions”. In *2008 49th Annual IEEE Symposium on Foundations of Computer Science*, 793–802. <https://doi.org/10.1109/FOCS.2008.11>.
- Achlioptas, Dimitris, and Federico Ricci-Tersenghi. 2006. “On the Solution-Space Geometry of Random Constraint Satisfaction Problems”. December 15, 2006. <https://doi.org/10.48550/arXiv.cs/0611052>.
- Addario-Berry, Louigi, Luc Devroye, Gabor Lugosi, and Roberto Imbuzeiro Oliveira. 2017. “Local Optima of the Sherrington-Kirkpatrick Hamiltonian”. December 21, 2017. <https://doi.org/10.48550/arXiv.1712.07775>.
- Alidaee, Bahram, Fred Glover, Gary A. Kochenberger, and Cesar Rego. 2005. “A New Modeling and Solution Approach for the Number Partitioning Problem”. *Journal of Applied Mathematics and Decision Sciences* 2005 (2): 113–21. <https://doi.org/10.1155/JAMDS.2005.113>.
- Argüello, Michael F., Thomas A. Feo, and Olivier Goldschmidt. 1996. “Randomized Methods for the Number Partitioning Problem”. *Computers & Operations Research* 23 (2): 103–11. [https://doi.org/10.1016/0305-0548\(95\)E0020-L](https://doi.org/10.1016/0305-0548(95)E0020-L).
- Arous, Gérard Ben, Reza Gheissari, and Aukosh Jagannath. 2020. “Algorithmic Thresholds for Tensor PCA”. *The Annals of Probability* 48 (4): 2052–87. <https://doi.org/10.1214/19-AOP1415>.
- Asproni, Luca, Davide Caputo, Blanca Silva, Giovanni Fazzi, and Marco Magagnini. 2020. “Accuracy and Minor Embedding in Subqubo Decomposition with Fully Connected Large Problems: A Case Study About the Number Partitioning Problem”. *Quantum Machine Intelligence* 2 (1): 4. <https://doi.org/10.1007/s42484-020-00014-w>.
- Aubin, Benjamin, Will Perkins, and Lenka Zdeborová. 2019. “Storage Capacity in Symmetric Binary Perceptrons”. *Journal of Physics A: Mathematical and Theoretical* 52 (29): 294003. <https://doi.org/10.1088/1751-8121/ab227a>.
- Bandeira, Afonso S., Amelia Perry, and Alexander S. Wein. 2018. “Notes on Computational-to-Statistical Gaps: Predictions Using Statistical Physics”. April 20, 2018. <https://doi.org/10.48550/arXiv.1803.11132>.
- Bansal, Nikhil. 2010. “Constructive Algorithms for Discrepancy Minimization”. August 9, 2010. <https://doi.org/10.48550/arXiv.1002.2259>.
- Barak, Boaz, Samuel B. Hopkins, Jonathan Kelner, Pravesh K. Kothari, Ankur Moitra, and Aaron Potechin. 2016. “A Nearly Tight Sum-of-Squares Lower Bound for the Planted Clique Problem”. April 12, 2016. <https://doi.org/10.48550/arXiv.1604.03084>.

- Bauke, Heiko, Silvio Franz, and Stephan Mertens. 2004. “Number Partitioning as a Random Energy Model”. *Journal of Statistical Mechanics: Theory and Experiment* 2004 (4): P4003. <https://doi.org/10.1088/1742-5468/2004/04/P04003>.
- Bayati, Mohsen, David Gamarnik, and Prasad Tetali. 2013. “Combinatorial Approach to the Interpolation Method and Scaling Limits in Sparse Random Graphs”. *The Annals of Probability* 41 (6). <https://doi.org/10.1214/12-AOP816>.
- Berthet, Quentin, and Philippe Rigollet. 2013. “Computational Lower Bounds for Sparse PCA”. April 26, 2013. <https://doi.org/10.48550/arXiv.1304.0828>.
- Bismuth, Samuel, Vladislav Makarov, Erel Segal-Halevi, and Dana Shapira. 2024. “Partitioning Problems with Splittings and Interval Targets”. September 18, 2024. <https://doi.org/10.48550/arXiv.2204.11753>.
- Boettcher, S., and S. Mertens. 2008. “Analysis of the Karmarkar-Karp Differencing Algorithm”. *The European Physical Journal B* 65 (1): 131–40. <https://doi.org/10.1140/epjb/e2008-00320-9>.
- Borgs, Christian, Jennifer Chayes, and Boris Pittel. 2001. “Phase Transition and Finite-size Scaling for the Integer Partitioning Problem”. *Random Structures & Algorithms* 19 (3–4): 247–88. <https://doi.org/10.1002/rsa.10004>.
- Brennan, Matthew, and Guy Bresler. 2019. “Optimal Average-Case Reductions to Sparse PCA: From Weak Assumptions to Strong Hardness”. February 20, 2019. <https://doi.org/10.48550/arXiv.1902.07380>.
- Brennan, Matthew, Guy Bresler, and Wasim Huleihel. 2019. “Reducibility and Computational Lower Bounds for Problems with Planted Sparse Structure”. November 18, 2019. <https://doi.org/10.48550/arXiv.1806.07508>.
- Chandrasekaran, Karthekeyan, and Santosh Vempala. 2013. “Integer Feasibility of Random Polytopes”. August 23, 2013. <https://doi.org/10.48550/arXiv.1111.4649>.
- Chen, Wei-Kuo, David Gamarnik, Dmitry Panchenko, and Mustazee Rahman. 2019. “Suboptimality of Local Algorithms for a Class of Max-Cut Problems”. *The Annals of Probability* 47 (3). <https://doi.org/10.1214/18-AOP1291>.
- Coffman, E. G., Jr., M. R. Garey, and D. S. Johnson. 1978. “An Application of Bin-Packing to Multiprocessor Scheduling”. *SIAM Journal on Computing* 7 (1): 1–17. <https://doi.org/10.1137/0207001>.
- Coffman, Edward Grady, and George S. Lueker. 1991. *Probabilistic Analysis of Packing and Partitioning Algorithms*. Wiley-Interscience Series in Discrete Mathematics and Optimization. New York: J. Wiley & sons.
- Coja-Oghlan, Amin, and Charilaos Efthymiou. 2015. “On Independent Sets in Random Graphs”. *Random Structures & Algorithms* 47 (3): 436–86. <https://doi.org/10.1002/rsa.20550>.
- Corus, Dogan, Pietro S. Oliveto, and Donya Yazdani. 2019. “Artificial Immune Systems Can Find Arbitrarily Good Approximations for the NP-hard Number Partitioning Problem”. *Artificial Intelligence* 274 (September):180–96. <https://doi.org/10.1016/j.artint.2019.03.001>.
- Deshpande, Yash, and Andrea Montanari. 2015. “Improved Sum-of-Squares Lower Bounds for Hidden Clique and Hidden Submatrix Problems”. February 23, 2015. <https://doi.org/10.48550/arXiv.1502.06590>.

- Diakonikolas, Ilias, Daniel M. Kane, and Alistair Stewart. 2017. “Statistical Query Lower Bounds for Robust Estimation of High-dimensional Gaussians and Gaussian Mixtures”. May 17, 2017. <https://doi.org/10.48550/arXiv.1611.03473>.
- Feldman, Vitaly, Elena Grigorescu, Lev Reyzin, Santosh Vempala, and Ying Xiao. 2016. “Statistical Algorithms and a Lower Bound for Detecting Planted Clique”. August 15, 2016. <https://doi.org/10.48550/arXiv.1201.1214>.
- Ferreira, F. F., and J. F. Fontanari. 1998. “Probabilistic Analysis of the Number Partitioning Problem”. *Journal of Physics A: Mathematical and General* 31 (15): 3417. <https://doi.org/10.1088/0305-4470/31/15/007>.
- Gamarnik, David. 2021. “The Overlap Gap Property: A Geometric Barrier to Optimizing over Random Structures”. *Proceedings of the National Academy of Sciences* 118 (41): e2108492118. <https://doi.org/10.1073/pnas.2108492118>.
- Gamarnik, David, and Aukosh Jagannath. 2019. “The Overlap Gap Property and Approximate Message Passing Algorithms for \mathbb{Z}_2 -Spin Models”. November 26, 2019. <https://doi.org/10.48550/arXiv.1911.06943>.
- Gamarnik, David, and Eren Kizildag. 2021. “Computing the Partition Function of the Sherrington-Kirkpatrick Model Is Hard on Average”. *The Annals of Applied Probability* 31 (3). <https://doi.org/10.1214/20-AAP1625>.
- Gamarnik, David, and Eren C. Kızıldağ. 2021. “Algorithmic Obstructions in the Random Number Partitioning Problem”. March 2, 2021. <https://doi.org/10.48550/arXiv.2103.01369>.
- Gamarnik, David, and Quan Li. 2016. “Finding a Large Submatrix of a Gaussian Random Matrix”. February 26, 2016. <https://doi.org/10.48550/arXiv.1602.08529>.
- Gamarnik, David, and Madhu Sudan. 2013. “Limits of Local Algorithms over Sparse Random Graphs”. April 5, 2013. <https://doi.org/10.48550/arXiv.1304.1831>.
- Gamarnik, David, and Madhu Sudan. 2017. “Performance of Sequential Local Algorithms for the Random NAE- \mathbb{Z}_2 -SAT Problem”. *SIAM Journal on Computing* 46 (2): 590–619. <https://doi.org/10.1137/140989728>.
- Gamarnik, David, and Ilias Zadik. 2019b. “High-Dimensional Regression with Binary Coefficients. Estimating Squared Error and a Phase Transition”. September 25, 2019. <https://doi.org/10.48550/arXiv.1701.04455>.
- Gamarnik, David, and Ilias Zadik. 2019a. “Sparse High-Dimensional Linear Regression. Algorithmic Barriers and a Local Search Algorithm”. September 22, 2019. <https://doi.org/10.48550/arXiv.1711.04952>.
- Gamarnik, David, and Ilias Zadik. 2019c. “The Landscape of the Planted Clique Problem: Dense Subgraphs and the Overlap Gap Property”. December 30, 2019. <https://doi.org/10.48550/arXiv.1904.07174>.
- Gamarnik, David, Aukosh Jagannath, and Subhabrata Sen. 2021. “The Overlap Gap Property in Principal Submatrix Recovery”. *Probability Theory and Related Fields* 181 (4): 757–814. <https://doi.org/10.1007/s00440-021-01089-7>.

- Gamarnik, David, Aukosh Jagannath, and Alexander S. Wein. 2022. “Hardness of Random Optimization Problems for Boolean Circuits, Low-Degree Polynomials, And Langevin Dynamics”. January 26, 2022. <https://doi.org/10.48550/arXiv.2004.12063>.
- Gamarnik, David, Eren C. Kızıldağ, Will Perkins, and Changji Xu. 2022. “Algorithms and Barriers in the Symmetric Binary Perceptron Model”. March 29, 2022. <https://doi.org/10.48550/arXiv.2203.15667>.
- Garey, Michael R., and David S. Johnson. 1979. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. A Series of Books in the Mathematical Sciences. New York: W. H. Freeman.
- Gent, Ian P., and Toby Walsh. 1998. “Analysis of Heuristics for Number Partitioning”. *Computational Intelligence* 14 (3): 430–51. <https://doi.org/10.1111/0824-7935.00069>.
- Gent, Ian, and Toby Walsh. 2000. “Phase Transitions and Annealed Theories: Number Partitioning as a Case Study”. *Instituto Cultura*, June.
- Harshaw, Christopher, Fredrik Sävje, Daniel Spielman, and Peng Zhang. 2023. “Balancing Covariates in Randomized Experiments with the Gram-Schmidt Walk Design”. November 14, 2023. <https://doi.org/10.48550/arXiv.1911.03071>.
- Hastie, Trevor, Robert Tibshirani, and Jerome Friedman. 2009. *The Elements of Statistical Learning*. Springer Series in Statistics. New York, NY: Springer New York. <https://doi.org/10.1007/978-0-387-84858-7>.
- Hatami, Hamed, László Lovász, and Balázs Szegedy. 2014. “Limits of Locally–Globally Convergent Graph Sequences”. *Geometric and Functional Analysis* 24 (1): 269–96. <https://doi.org/10.1007/s00039-014-0258-7>.
- Hoberg, Rebecca, Harishchandra Ramadas, Thomas Rothvoss, and Xin Yang. 2016. “Number Balancing Is as Hard as Minkowski’s Theorem and Shortest Vector”. November 29, 2016. <https://doi.org/10.48550/arXiv.1611.08757>.
- Hopkins, Samuel. 2018. “Statistical Inference and the Sum of Squares Method.”
- Hopkins, Samuel B., Pravesh K. Kothari, Aaron Potechin, Prasad Raghavendra, Tselil Schramm, and David Steurer. 2017. “The Power of Sum-of-Squares for Detecting Hidden Structures”. October 13, 2017. <https://doi.org/10.48550/arXiv.1710.05017>.
- Hopkins, Samuel B., Jonathan Shi, and David Steurer. 2015. “Tensor Principal Component Analysis via Sum-of-Squares Proofs”. July 12, 2015. <https://doi.org/10.48550/arXiv.1507.03269>.
- Huang, Brice, and Mark Sellke. 2025. “Strong Low Degree Hardness for Stable Local Optima in Spin Glasses”. January 11, 2025. <https://doi.org/10.48550/arXiv.2501.06427>.
- Huang, Han, and Elchanan Mossel. 2025. “Optimal Low Degree Hardness for Broadcasting on Trees”. February 7, 2025. <https://doi.org/10.48550/arXiv.2502.04861>.
- Jerrum, Mark. 1992. “Large Cliques Elude the Metropolis Process”. *Random Structures & Algorithms* 3 (4): 347–59. <https://doi.org/10.1002/rsa.3240030402>.
- Johnson, David S., Cecilia R. Aragon, Lyle A. McGeoch, and Catherine Schevon. 1989. “Optimization by Simulated Annealing: An Experimental Evaluation; Part I, Graph Partitioning”. *Operations Research* 37 (6): 865–92. <http://www.jstor.org/stable/171470>.

- Johnson, David S., Cecilia R. Aragon, Lyle A. McGeoch, and Catherine Schevon. 1991. "Optimization by Simulated Annealing: An Experimental Evaluation; Part II, Graph Coloring and Number Partitioning". *Operations Research* 39 (3): 378–406. <http://www.jstor.org/stable/171393>.
- Karmarkar, Narendra, and Richard M. Karp. 1983. "The Differencing Method of Set Partitioning". <https://www2.eecs.berkeley.edu/Pubs/TechRpts/1983/6353.html>.
- Karmarkar, Narendra, Richard M. Karp, George S. Lueker, and Andrew M. Odlyzko. 1986. "Probabilistic Analysis of Optimum Partitioning". *Journal of Applied Probability* 23 (3): 626–45. <https://doi.org/10.2307/3214002>.
- Kearns, Michael. 1998. "Efficient Noise-Tolerant Learning from Statistical Queries". *Journal of the ACM* 45 (6): 983–1006. <https://doi.org/10.1145/293347.293351>.
- Kojić, Jelena. 2010. "Integer Linear Programming Model for Multidimensional Two-Way Number Partitioning Problem". *Computers & Mathematics with Applications* 60 (8): 2302–8. <https://doi.org/10.1016/j.camwa.2010.08.024>.
- Korf, Richard E. 1995. "From Approximate to Optimal Solutions: A Case Study of Number Partitioning". In *Proceedings of the 14th International Joint Conference on Artificial Intelligence - Volume 1*, 266–72. IJCAI'95. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc.
- Korf, Richard E. 1998. "A Complete Anytime Algorithm for Number Partitioning". *Artificial Intelligence* 106 (2): 181–203. [https://doi.org/10.1016/S0004-3702\(98\)00086-1](https://doi.org/10.1016/S0004-3702(98)00086-1).
- Korf, Richard E. 2009. "Multi-Way Number Partitioning". In *Proceedings of the 21st International Joint Conference on Artificial Intelligence*, 538–43. IJCAI'09. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc.
- Kothari, Pravesh K., Ryuhei Mori, Ryan O'Donnell, and David Witmer. 2017. "Sum of Squares Lower Bounds for Refuting Any CSP". January 17, 2017. <https://doi.org/10.48550/arXiv.1701.04521>.
- Kratika, Jozef, Jelena Kojić, and Aleksandar Savić. 2014. "Two Metaheuristic Approaches for Solving Multidimensional Two-Way Number Partitioning Problem". *Computers & Operations Research* 46 (June):59–68. <https://doi.org/10.1016/j.cor.2014.01.003>.
- Krieger, A M, D Azriel, and A Kapelner. 2019. "Nearly Random Designs with Greatly Improved Balance". *Biometrika* 106 (3): 695–701. <https://doi.org/10.1093/biomet/asz026>.
- Kunisky, Dmitriy. 2024a. "Low Coordinate Degree Algorithms I: Universality of Computational Thresholds for Hypothesis Testing". March 12, 2024. <https://doi.org/10.48550/arXiv.2403.07862>.
- Kunisky, Dmitriy. 2024b. "Low Coordinate Degree Algorithms II: Categorical Signals and Generalized Stochastic Block Models". December 30, 2024. <https://doi.org/10.48550/arXiv.2412.21155>.
- Kunisky, Dmitriy, Alexander S. Wein, and Afonso S. Bandeira. 2019. "Notes on Computational Hardness of Hypothesis Testing: Predictions Using the Low-Degree Likelihood Ratio". July 26, 2019. <https://doi.org/10.48550/arXiv.1907.11636>.
- Kızıldağ, Eren C. 2023. "Planted Random Number Partitioning Problem". September 26, 2023. <https://doi.org/10.48550/arXiv.2309.15115>.

- Lauer, Joseph, and Nicholas Wormald. 2007. “Large Independent Sets in Regular Graphs of Large Girth”. *Journal of Combinatorial Theory, Series B* 97 (6): 999–1009. <https://doi.org/10.1016/j.jctb.2007.02.006>.
- Lesieur, Thibault, Florent Krzakala, and Lenka Zdeborova. 2015. “Phase Transitions in Sparse PCA”. In *2015 IEEE International Symposium on Information Theory (ISIT)*, 1635–39. <https://doi.org/10.1109/ISIT.2015.7282733>.
- Lesieur, Thibault, Florent Krzakala, and Lenka Zdeborová. 2015. “MMSE of Probabilistic Low-Rank Matrix Estimation: Universality with Respect to the Output Channel”. In *2015 53rd Annual Allerton Conference on Communication, Control, And Computing (Allerton)*, 680–87. <https://doi.org/10.1109/ALLERTON.2015.7447070>.
- Levy, Avi, Harishchandra Ramadas, and Thomas Rothvoss. 2017. “Deterministic Discrepancy Minimization via the Multiplicative Weight Update Method”. March 11, 2017. <https://doi.org/10.48550/arXiv.1611.08752>.
- Lovett, Shachar, and Raghu Meka. 2012. “Constructive Discrepancy Minimization by Walking on The Edges”. October 11, 2012. <https://doi.org/10.48550/arXiv.1203.5747>.
- Lueker, George S. 1987. “A Note on the Average-Case Behavior of a Simple Differencing Method for Partitioning”. *Operations Research Letters* 6 (6): 285–87. [https://doi.org/10.1016/0167-6377\(87\)90044-7](https://doi.org/10.1016/0167-6377(87)90044-7).
- Meka, Raghu, Aaron Potechin, and Avi Wigderson. 2015. “Sum-of-Squares Lower Bounds for Planted Clique”. In *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing*, 87–96. Portland Oregon USA: ACM. <https://doi.org/10.1145/2746539.2746600>.
- Merkle, R., and M. Hellman. 1978. “Hiding Information and Signatures in Trapdoor Knapsacks”. *IEEE Transactions on Information Theory* 24 (5): 525–30. <https://doi.org/10.1109/TIT.1978.1055927>.
- Mertens, Stephan. 2001. “A Physicist's Approach to Number Partitioning”. *Theoretical Computer Science, Phase Transitions in Combinatorial Problems*, 265 (1): 79–108. [https://doi.org/10.1016/S0304-3975\(01\)00153-0](https://doi.org/10.1016/S0304-3975(01)00153-0).
- Mertens, Stephan. 2003. “The Easiest Hard Problem: Number Partitioning”. October 18, 2003. <https://doi.org/10.48550/arXiv.cond-mat/0310317>.
- Michiels, Wil, Jan Korst, Emile Aarts, and Jan Van Leeuwen. 2003. “Performance Ratios for the Differencing Method Applied to the Balanced Number Partitioning Problem”. Edited by Helmut Alt, Michel Habib, Gerhard Goos, Juris Hartmanis, and Jan Van Leeuwen. *STACS 2003*. Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/3-540-36494-3_51.
- Montanari, Andrea. 2019. “Optimization of the Sherrington-Kirkpatrick Hamiltonian”. April 5, 2019. <https://doi.org/10.48550/arXiv.1812.10897>.
- Montanari, Andrea, and Alexander S. Wein. 2024. “Equivalence of Approximate Message Passing and Low-Degree Polynomials in Rank-One Matrix Estimation”. October 10, 2024. <https://doi.org/10.48550/arXiv.2212.06996>.
- Mézard, M., T. Mora, and R. Zecchina. 2005. “Clustering of Solutions in the Random Satisfiability Problem”. *Physical Review Letters* 94 (19): 197205. <https://doi.org/10.1103/PhysRevLett.94.197205>.

- O'Donnell, Ryan. 2021. "Analysis of Boolean Functions". May 21, 2021. <https://doi.org/10.48550/arXiv.2105.10386>.
- Raghavendra, Prasad, Tselil Schramm, and David Steurer. 2019. "High-Dimensional Estimation via Sum-of-Squares Proofs". August 6, 2019. <https://doi.org/10.48550/arXiv.1807.11419>.
- Rahman, Mustazee, and Balint Virag. 2017. "Local Algorithms for Independent Sets Are Half-Optimal". *The Annals of Probability* 45 (3). <https://doi.org/10.1214/16-AOP1094>.
- Rothvoss, Thomas. 2016. "Constructive Discrepancy Minimization for Convex Sets". April 12, 2016. <https://doi.org/10.48550/arXiv.1404.0339>.
- Santucci, Valentino, Marco Baioletti, and Gabriele Di Bari. 2021. "An Improved Memetic Algebraic Differential Evolution for Solving the Multidimensional Two-Way Number Partitioning Problem". *Expert Systems with Applications* 178 (September):114938. <https://doi.org/10.1016/j.eswa.2021.114938>.
- Shamir, Adi. 1982. "A Polynomial Time Algorithm for Breaking the Basic Merkle-Hellman Cryptosystem". In *23rd Annual Symposium on Foundations of Computer Science (Sfcs 1982)*, 145–52. <https://doi.org/10.1109/SFCS.1982.5>.
- Storer, Robert H., Seth W. Flanders, and S. David Wu. 1996. "Problem Space Local Search for Number Partitioning". *Annals of Operations Research* 63 (4): 463–87. <https://doi.org/10.1007/BF02156630>.
- Tsai, Li-Hui. 1992. "Asymptotic Analysis of an Algorithm for Balanced Parallel Processor Scheduling". *SIAM Journal on Computing* 21 (1): 59–64. <https://doi.org/10.1137/0221007>.
- Turner, Paxton, Raghu Meka, and Philippe Rigollet. 2020. "Balancing Gaussian Vectors in High Dimension". June 30, 2020. <https://doi.org/10.48550/arXiv.1910.13972>.
- Vafa, Neekon, and Vinod Vaikuntanathan. 2025. "Symmetric Perceptrons, Number Partitioning and Lattices". January 27, 2025. <https://doi.org/10.48550/arXiv.2501.16517>.
- Vershynin, Roman. 2018. *High-Dimensional Probability: An Introduction with Applications in Data Science*. 1st ed. Cambridge Series in Statistical and Probabilistic Mathematics. New York, NY: Cambridge University Press.
- Wainwright, Martin J. 2019. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press. <https://doi.org/10.1017/9781108627771>.
- Wein, Alexander S. 2020. "Optimal Low-Degree Hardness of Maximum Independent Set". November 12, 2020. <https://doi.org/10.48550/arXiv.2010.06563>.
- Wen, Jingwei, Zhenming Wang, Zhiguo Huang, Dunbo Cai, Bingjie Jia, Chongyu Cao, Yin Ma, Hai Wei, Kai Wen, and Ling Qian. 2023. "Optical Experimental Solution for the Multiway Number Partitioning Problem and Its Application to Computing Power Scheduling". *Science China Physics, Mechanics & Astronomy* 66 (9): 290313. <https://doi.org/10.1007/s11433-023-2147-3>.
- Yakir, Benjamin. 1996. "The Differencing Algorithm LDM for Partitioning: A Proof of a Conjecture of Karmarkar and Karp". *Mathematics of Operations Research* 21 (1): 85–99. <https://doi.org/10.1287/moor.21.1.85>.
- Zdeborová, Lenka, and Florent Krzakala. 2016. "Statistical Physics of Inference: Thresholds and Algorithms". *Advances in Physics* 65 (5): 453–552. <https://doi.org/10.1080/00018732.2016.1211393>.