

Breast Cancer Clinical Outcome Analysis

Age Group

- ☐ <40
- ☒ 40-50
- ☐ 50-60
- ☐ 60-70
- ☐ 70+

Tumor Stage

- ☐ 0
- ☐ 1
- ☐ 2
- ☐ 3
- ☐ 4


Cancer Type Detailed

- ☐ Breast Invasive Ductal Carcino...
- ☐ Breast Invasive Lobular Carcin...
- ☐ Breast Invasive Mixed Mucino...
- ☐ Breast Mixed Ductal and Lobu...
- ☐ Invasive Breast Carcinoma


Patient ID

- MTS-T2417
- MTS-T2392
- MTS-T2391
- MTS-T2320
- MTS-T2206
- MTS-T2190
- MTS-T2128
- MTS-T2087
- MTS-T2084
- MTS-T2076
- MTS-T2066
- MTS-T2064


Count of Patient ID by Tumor Stage




Count of Patient ID by Type of Breast Surgery



Count of Patient ID by Cancer Type Detailed



Count of Patient ID by Chemotherapy and Hormone Therapy



Breast Cancer Clinical Outcome Analysis

ER Status

- ☐ Negative
- ☐ Positive

Chemotherapy

- ☐
- ☐ NO
- ☒ YES

NPI Risk

- ☐ Good
- ☐ Moderate
- ☐ Poor


PR Status

- ☒ Negative
- ☐ Positive


Hormone Therapy

- ☐ NO
- ☐ YES


Average of Overall Survival (Months) by Overall Survival Status




Count of Patient ID by Overall Survival Status




Count of Patient ID by ER Status and HER2 Status



Sum of Tumor Size by Tumor Stage, Mutation Count and Overall Survival (Years)



Count of Patient ID by NPI Risk



Key Insights

- Stage II was the most common tumour stage.
- Invasive ductal carcinoma accounted for the majority of cases.
- Most patients underwent either mastectomy or breast-conserving surgery.
- Patients receiving chemotherapy showed significantly different survival outcomes (Log-rank test: $p = 0.00179$).
- Moderate NPI risk was the largest patient group.