1.5.5 Repeat the above excercise for the sides AB and AC.

Solution: : We know the value of \mathbf{I} is

$$\mathbf{I} = \frac{1}{\sqrt{37} + 4 + \sqrt{61}} \begin{pmatrix} \sqrt{61} - 16 - 3\sqrt{37} \\ -\sqrt{61} + 24 - 5\sqrt{37} \end{pmatrix}$$
 (1)

from the problem 1.5.2 . The equation of AB is:

$$\begin{pmatrix}
7 & 5
\end{pmatrix} \mathbf{x} - 2 = 0
\tag{2}$$

Let r_1 be the distance between **I** and AB, then

$$r_1 = \frac{\left| \begin{pmatrix} 7 & 5 \end{pmatrix} \mathbf{I} - 2 \right|}{\left\| \begin{pmatrix} 7 \\ 5 \end{pmatrix} \right\|} \tag{3}$$

$$= \frac{\begin{vmatrix} \sqrt{5} & \sqrt{5} & \sqrt{5} & \sqrt{61} - 16 - 3\sqrt{37} \\ \sqrt{37} + 4 + \sqrt{61} & \sqrt{7} & \sqrt{61} + 24 - 5\sqrt{37} \end{vmatrix} - 2 \end{vmatrix}}{\sqrt{7^2 + 5^2}}$$
(4)

$$=\frac{\frac{2\sqrt{61}-46\sqrt{37}+8}{\sqrt{37}+4+\sqrt{61}}-2}{\sqrt{74}}\tag{5}$$

$$=\frac{48\sqrt{37}}{\sqrt{74}(\sqrt{37}+4+\sqrt{61})}\tag{6}$$

$$=\frac{48}{\sqrt{2}(\sqrt{37}+4+\sqrt{61})}\tag{7}$$

$$=\frac{24\sqrt{2}}{\sqrt{37}+4+\sqrt{61}}\tag{8}$$

$$=1.8969$$
 (9)

Similarly, the equation of AC is

$$\begin{pmatrix}
4 & -4
\end{pmatrix} \mathbf{x} - 8 = 0
\tag{10}$$

Let r_2 be the distance between **I** and AC, then

$$r_{2} = \frac{\left| \begin{pmatrix} 4 & -4 \end{pmatrix} \mathbf{I} - 8 \right|}{\left\| \begin{pmatrix} 4 \\ -4 \end{pmatrix} \right\|} \tag{11}$$

$$= \frac{\begin{vmatrix} \frac{1}{\sqrt{37}+4+\sqrt{61}} \left(4 & -4\right) \left(\sqrt{61}-16-3\sqrt{37}\right) - 8 \\ -\sqrt{61}+24-5\sqrt{37} - 8 \end{vmatrix}}{\sqrt{4^2+(-4)^2}}$$
(12)

$$=\frac{\left|\frac{8\sqrt{61}+8\sqrt{37}-160}{\sqrt{37}+4+\sqrt{61}}-8\right|}{4\sqrt{2}}\tag{13}$$

$$=\frac{192}{4\sqrt{2}(\sqrt{37}+4+\sqrt{61})}\tag{14}$$

$$=\frac{48}{\sqrt{2}(\sqrt{37}+4+\sqrt{61})}\tag{15}$$

$$=\frac{24\sqrt{2}}{\sqrt{37}+4+\sqrt{61}}\tag{16}$$

$$=1.8969$$
 (17)